



# Effect of Anesthesia Type on Pain and Hemodynamic Parameters in Cesarean Section Surgery: Randomized Controlled Trial

Serveh Mohammadi <sup>1</sup>, Behzad Imani <sup>2,\*</sup>, Sina Ghasemi <sup>1</sup>, Behzad Shalmashi<sup>3</sup>

<sup>1</sup> Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>2</sup> Department of Operating Room, Faculty of Paramedicine, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>3</sup> Department of Anesthesiology, Faculty of Medicine, Urmia University of Medical Science, Mahabad, Iran

\*Corresponding Author: Department of Operating Room, Faculty of Paramedicine, Hamadan University of Medical Sciences, Hamadan, Iran. Email: behzadiman@yahoo.com

Received 2024 May 26; Revised 2024 September 7; Accepted 2024 September 10

## Abstract

**Background:** General and spinal anesthesia are commonly used for cesarean sections, each with implications for both the mother and the fetus. The choice of anesthesia technique depends on various factors. While spinal anesthesia is often preferred for cesarean surgeries due to its well-known benefits, general anesthesia may be chosen in emergency situations. The decision is ultimately based on the safety profile and benefits for both the mother and the fetus.

**Methods:** This clinical trial was conducted at Fatemieh Hospital in Hamadan and involved 60 patients undergoing elective cesarean section. The patients were randomly divided into two groups of 30: One group received spinal anesthesia, and the other group received general anesthesia. Variables such as intraoperative bleeding, the Apgar score of the newborn, postoperative pain, and postoperative systolic and diastolic blood pressure were analyzed and compared between the two groups using SPSS software version 16.

**Results:** The mean age of participants in the spinal and general anesthesia groups was  $32 \pm 3.5$  years and  $34 \pm 3.4$  years, respectively. The results showed that the amount of intraoperative bleeding in the general anesthesia group was higher than in the spinal anesthesia group, but this difference was not statistically significant ( $P = 0.85$ ,  $P < 0.05$ ). The Apgar scores at the first and fifth minutes in the spinal anesthesia group were higher than those in the general anesthesia group, but this difference was also not statistically significant ( $P = 0.32$ ,  $P < 0.05$ ). Postoperative pain in the spinal anesthesia group was significantly lower than in the general anesthesia group ( $P = 0.001$ ,  $P < 0.05$ ). Regarding hemodynamic parameters, systolic blood pressure after the operation was higher in the spinal anesthesia group compared to the general anesthesia group ( $P < 0.05$ ). Conversely, diastolic blood pressure in the general anesthesia group was higher than in the spinal anesthesia group at all measurement stages, though not statistically significant in some measurements ( $P > 0.05$ ).

**Conclusions:** The results of our study indicate that spinal anesthesia is a preferable and safer method compared to general anesthesia, based on the parameters examined, particularly in emergency cesarean sections.

**Keywords:** Cesarean Section, Regional Anesthesia, Spinal Anesthesia, General Anesthesia, Measurement Outcomes

## 1. Background

Cesarean section, commonly referred to as a C-section, is a surgical procedure in which a baby is delivered through an incision made in the mother's abdomen and uterus, typically when vaginal delivery is deemed unsafe or not feasible for both the mother and the baby (1). The choice between general anesthesia and regional anesthesia for a cesarean section depends on the patient's clinical condition and whether the surgery is elective or an emergency (2). While neuraxial

anesthesia (spinal and epidural anesthesia) is considered the gold standard for cesarean sections, general anesthesia is still used, particularly in cases where regional anesthesia is contraindicated or unsuccessful (1, 3). General anesthesia is often preferred in emergencies, such as in cases of fetal distress during pregnancy, as it allows for rapid induction (4).

Contraindications for regional anesthesia include conditions such as increased intracranial pressure, hypovolemic shock, sepsis, coagulation disorders, uterine atony, infection or inflammation at the injection

site, severe mitral valve stenosis, and a high risk of severe bleeding (2). Additionally, the hypotension associated with regional anesthesia may affect short-term neonatal outcomes (5). Cesarean sections under general anesthesia offer advantages, including rapid induction, optimal control of the airway, and a reduced risk of hypotension and cardiovascular instability, especially in cases such as preeclampsia with neurological complications (2).

The World Health Organization recommends an ideal cesarean section rate of 10 - 15% (5). However, the prevalence of cesarean sections, particularly in Iran, exceeds this recommended rate, posing significant challenges (6). Cesarean sections are considered life-saving surgical procedures for both the mother and child, with studies estimating their role in preventing a substantial number of maternal and neonatal deaths annually (7). Therefore, anesthesia management during cesarean sections is critical for ensuring the safety of both the newborn and the mother (7).

Due to emergency conditions and time constraints, general anesthesia is often more commonly used than regional anesthesia (8). The choice of anesthesia technique depends on several factors, including the patient's physiological condition, the physician's expertise, and the availability of drugs and medical equipment (7). Spinal anesthesia is frequently preferred for cesarean sections due to its advantages, such as reduced complications associated with general anesthesia and enhanced maternal-fetal bonding (9). However, spinal anesthesia can also lead to side effects like arterial hypotension and fetal distress (10).

General anesthesia involves the administration of intravenous or inhalation anesthetics, while spinal anesthesia requires the injection of a local anesthetic into the subarachnoid space (11). Both methods have associated risks and side effects. Given recent case reports and the preference for general anesthesia over regional anesthesia in certain situations, this study aims to conduct a randomized clinical trial to compare the hemodynamic parameters and fetal outcomes during cesarean sections performed under spinal versus general anesthesia (2, 3).

## 2. Objectives

This study aims to compare hemodynamic parameters, intraoperative bleeding, blood pressure, postoperative pain, and neonatal outcomes during and after cesarean section in term, singleton pregnancies without complications, performed under spinal and general anesthesia.

## 3. Methods

This study was a randomized controlled clinical trial with two groups: An intervention group that received spinal anesthesia and a control group that received general anesthesia for elective cesarean section. The study was conducted at Fatemeh Hospital in Hamadan, following ethical approval and informed consent from the participants. Ethical considerations outlined in the Declaration of Helsinki were strictly adhered to throughout the study.

### 3.1. Participants

The study sample consisted of 62 candidates scheduled for elective cesarean section. Two participants were excluded from the study, leaving 30 participants in each group. Data collection included fieldwork and the completion of a demographic information questionnaire, which gathered details such as age, hospitalization history, and education level.

### 3.2. Measurement Methods

Intraoperative bleeding was quantified by weighing blood-contaminated gauzes, with each gram of gauze weight considered equivalent to one milliliter of blood loss. Additionally, the volume of blood in the suction reservoir was measured after subtracting wash fluids and amniotic fluid (12). Pain intensity during the first 24 hours post-operation was evaluated using the Numeric Rating Scale (NRS) (13). Blood pressure was monitored using a manometer within the first 24 hours post-surgery. The Apgar scale was used to assess newborns' Apgar scores at the first and fifth minute after delivery. This scale evaluates five parameters: Appearance (skin color), pulse (heart rate), reflex response, activity (muscle tone), and respiration (respiratory rate and effort). Each parameter is scored from 0 to 2, yielding a total score between 0 and 10. A score of 7 or higher at 1 minute indicates good health, while a score below 7 may require immediate medical attention. A 5-minute score below 7 may indicate an increased risk of mortality or cerebral palsy (14, 15). Pain intensity was assessed on a scale of 0 to 10, with higher scores indicating greater pain (16). The average pain intensity was recorded using the NRS pain intensity scale at 0, 6, 12, 18, and 24 hours post-admission. Blood pressure was also monitored at various intervals following admission to the ward: Upon admission, and at 2, 4, 6, 12, and 24 hours thereafter.

### 3.3. Inclusion and Exclusion Criteria

### 3.3.1. Inclusion Criteria

- Pregnant women aged between 18 and 40 years.
- Absence of a history of hypertension.
- No history of coagulation disorders.
- Newborn birth weight between 2.5 to 3.8 kilograms, with full-term infants.
- Hemoglobin level of at least 90 grams per liter in patients.
- Categorization of patients according to the American Society of Anesthesiologists (ASA) as class 1 or class 2 in physical status (17).
- Informed consent obtained for participation in the study.

### 3.3.2. Exclusion Criteria

- Need for an emergency cesarean section.
- History of maternal stillbirth.
- Multiple pregnancies.
- Presence of chromosomal abnormalities.
- Need for a pain pump.
- Presence of polyhydramnios.

Considering a confidence level of 95% and a power of 8%, the sample size was calculated to be 28 participants in each group. Taking into account a 10% dropout probability, the final number of participants was determined to be 31 in each group (18).

$$N = \frac{2 \left( Z_{1-\frac{\alpha}{2}} + Z_{1-\beta} \right)^2 P(1-P)}{D^2}$$

$$= \frac{2(1.96 + 0.84)^2 0.16(1.016)}{(0.8)^2} = 31$$

### 3.4. Data Collection Tools

The data collection in this study comprises four parts:

**Numeric Rating Scale criteria:** In this method, individuals rate their pain on a scale from zero to ten. Zero represents no pain, 1 to 3 indicates mild pain, 4 to 6 represents moderate pain, and 7 to 10 indicates severe pain. According to previous studies, this scale demonstrates good validity and reliability.

**Measurement of bleeding:** The volume of blood loss was estimated by measuring the amount of blood-contaminated gauze and the volume of blood in the suction reservoir.

**Blood pressure monitoring:** Continuous monitoring of the patient's blood pressure was conducted for the first 24 hours after surgery using a monitor and manometer. Blood pressure measurements were taken upon admission to the ward and at 2, 4, 6, 12, and 24 hours after admission, and recorded.

**Apgar Scale:** The Apgar scale was used to assess the effects of anesthesia on the newborn (19). The Apgar score consists of five ordinal qualitative variables, each assigned a numerical value ranging from zero to two, which are then summed together to create a continuous quantitative variable (20).

In the general anesthesia group, after 5 minutes of preoxygenation with a mask, anesthesia induction was performed using thiopental sodium at a dose of 3 mg/kg and succinylcholine at a dose of 1.5 mg/kg. Endotracheal intubation was performed using a size 7.5 or 7 endotracheal tube. After clamping of the umbilical cord, intravenous fentanyl at a dose of 2 µg/kg and atracurium at a dose of 0.5 mg/kg were administered for intraoperative pain management and muscle relaxation, respectively. Neostigmine at a dose of 0.05 mg/kg and atropine at a dose of 0.01 mg/kg were administered to reverse the effects of atracurium.

In the spinal anesthesia group, after skin preparation and sterile draping in the sitting position, a 25-gauge needle was inserted into the L3-4 or L4-5 space. Upon observing cerebrospinal fluid, 2.2 mm of hyperbaric bupivacaine 0.5% at a speed of 0.2 mm/s was injected into the subarachnoid space. Ephedrine was used to manage hypotension during the procedure. Additionally, 3 mm of intravenous fentanyl were administered to the spinal anesthesia group to provide longer-lasting pain relief and reduce postoperative nausea and vomiting (21).

The duration of surgery in both groups was 30 minutes, and 50 units of oxytocin, equivalent to 5 mm, were administered intravenously in both groups.

### 3.5. Statistical Method

Data analysis was conducted using SPSS version 16 software, with a significance level set at 0.05. Initially, the assumption of normality for the data was assessed using the Kolmogorov-Smirnov and Shapiro-Wilk tests. In cases where the data did not meet the normality assumption, non-parametric equivalents were used. Descriptive statistics, including frequency, percentage, mean, and standard deviation, were calculated. Analytical statistics involved the use of chi-square tests and independent *t*-tests.

## 4. Results

**Table 1.** History of Hospitalization and Level of Education Between two Groups

Patient Profile	Cesarean Section with Spinal <sup>a</sup>	Cesarean Section with General Anesthesia <sup>a</sup>	Chi-Square Test Statistic	P-Value
<b>Hospitalization history</b>			0.693	0.405
Yes	19 (46.3)	22(53.7)		
No	11 (57.9)	8 (42.1)		
<b>Level of education</b>			0.125	0.939
High school	9 (52.9)	8 (42.1)		
Diploma	7(46.7)	8 (53.3)		
Bachelor degree	14 (50)	1(50)		

<sup>a</sup> Values are expressed as No. (%).

**Table 2.** Mother's age Between the Two Groups

Variable Group	Cesarean Section with Spinal <sup>a</sup>	Cesarean Section with General Anesthesia <sup>a</sup>	Independent T-Test	P-Value
Age	32 ± 5.03	34.5 ± 6.3	-1.69	0.095

<sup>a</sup> Values are expressed as mean ± standard deviation.

A total of 60 patients were included in the study and divided into two groups of 30 each, based on the method of anesthesia: General anesthesia and spinal anesthesia. The most common indication for cesarean section across both groups was a previous cesarean section, accounting for 55% of all patients. The demographic characteristics of the mothers in both groups are summarized in [Tables 1-3](#).

The mean age of participants in the spinal anesthesia group was 32 years [standard deviation (SD) = 3.5], while in the general anesthesia group, the mean age was 34 years (SD = 3.6). Results from the independent *t*-test indicated that the age difference between the two groups was not statistically significant ( $P > 0.05$ ). The frequency of patients with a history of hospitalization was 19 (63.3%) in the spinal anesthesia group and 22 (73.3%) in the general anesthesia group. According to the chi-square test, there was no significant difference in hospitalization history between the two groups ( $P > 0.05$ ).

In terms of educational level distribution, 14 individuals (46.7%) in both groups had a bachelor's degree. In the spinal anesthesia group, 7 individuals (23.3%) held a diploma, and 9 individuals (30%) had less than a diploma. In the general anesthesia group, 8 individuals (26.7%) held a diploma, and another 8 (26.7%) had less than a diploma. The chi-square test results showed no significant difference in educational level distribution between the two groups ( $P > 0.05$ ).

Using an independent *t*-test, the amount of intraoperative bleeding and the Apgar scores of

newborns at the first and fifth minutes after birth were compared between the spinal anesthesia and general anesthesia groups. The results are presented in [Table 3](#). It was observed that the amount of bleeding during surgery was slightly higher in the general anesthesia group compared to the spinal anesthesia group. However, the results of the independent *t*-test indicated that this difference was not statistically significant ( $P = 0.85$ ).

Regarding the Apgar scores at the first and fifth minutes after birth, the values in the spinal anesthesia group were slightly higher than those in the general anesthesia group. Nevertheless, the independent *t*-test results showed no statistically significant difference between the two groups in terms of Apgar scores at the first and fifth minutes ( $P = 0.32$ ).

These findings suggest that the use of spinal anesthesia, compared to general anesthesia, did not significantly affect the amount of intraoperative bleeding or the Apgar scores of newborns at the first and fifth minutes after birth.

Using an independent *t*-test, the pain levels were compared between the spinal anesthesia and general anesthesia groups at the time of admission, every 6 hours, at 12 hours, 18 hours, and 24 hours post-operation. The results are presented in [Table 4](#). As observed, the mean pain score in the general anesthesia group was consistently higher than in the spinal anesthesia group at all assessment stages. The results of

**Table 3.** Comparison of Intraoperative Bleeding and Newborn Apgar Scores Among Study Groups

Variables and Groups	Mean ± Standard Deviation	F	t	Mean Difference	P-Value
<b>Operation bloodless</b>		3.981	0.190	6.16	0.85
Spinal anesthesia	451.66 ± 146.90				
General anesthesia	445.50 ± 100.57				
<b>Apgar score in the first minute after birth</b>		4.391	1.000	0.03	0.85
Spinal anesthesia	9.00 ± 0.00				
General anesthesia	8.96 ± 0.18				
<b>Apgar scored in the fifth minute after the birth</b>		4.291	1.000	0.03	0.32
Spinal anesthesia	10.00 ± 0.00				
General anesthesia	9.96 ± 0.18				

this test demonstrated that the difference between the two groups was statistically significant ( $P = 0.001$ ).

As evident from Figures 1 and 2, blood pressure at various measured times did not follow a consistent trend. For instance, upon admission to the ward, the mean systolic and diastolic blood pressures were 117/76 millimeters of mercury (mmHg) and 60/76 mmHg, respectively. Two hours post-admission, these values increased slightly to 118/76 mmHg and 26/76 mmHg, respectively. However, at 4 hours post-admission, systolic and diastolic blood pressures decreased to 116/75 mmHg and 70/75 mmHg, respectively, compared to two hours earlier and upon admission. Similarly, at the final measurement (24 hours post-admission), the mean systolic and diastolic blood pressures were 118/77 mmHg. Overall, the highest and lowest systolic blood pressure values were recorded at 24 hours post-admission (118/40 mmHg) and 6 hours post-admission (114/05 mmHg), respectively. Regarding diastolic blood pressure, the highest and lowest values were recorded at the same time points (24 hours and 6 hours post-admission), at 77/66 mmHg and 74/55 mmHg, respectively.

Using an independent *t*-test, blood pressure levels at admission and at 2, 4, 6, 12, and 24 hours post-admission were compared between the spinal anesthesia and general anesthesia groups. The results are depicted in Figures 1 and 2. Overall, systolic blood pressure in the spinal anesthesia group was higher than in the general anesthesia group. Conversely, diastolic blood pressure in the general anesthesia group was higher than in the spinal anesthesia group at all measurement stages. The independent *t*-test results showed that the difference in systolic blood pressure between the two groups was significant for all measurements ( $P < 0.05$ ), while for diastolic blood pressure, the difference between the spinal anesthesia and general anesthesia groups was not significant at certain time points (2 hours, 6 hours, and 12 hours post-admission) ( $P > 0.05$ ).

## 5. Discussion

We conducted this study on selected cesarean sections to compare two types of anesthesia. No statistically significant differences were found between the two groups regarding age, previous hospitalization history, or maternal education level, which enhances the reliability of our study results. Regarding bleeding, our results indicate that cesarean sections performed under general anesthesia tend to have slightly more bleeding compared to those performed under spinal anesthesia. The mean blood loss during surgery in patients under general anesthesia was higher than in those under spinal anesthesia; however, this difference was not statistically significant.

In this study, blood loss was measured based on the volume of blood-contaminated gauzes and the contents of the suction canister, with deductions made for amniotic fluid and irrigation fluids, rather than using a specific method to estimate blood loss. Multiple studies have demonstrated that cesarean deliveries under general anesthesia result in greater blood loss compared to those under regional anesthesia (22). In a prospective randomized study on selected cesarean cases, the average hemoglobin level in women under general anesthesia decreased by 1.1 g/dL compared to spinal anesthesia, which is consistent with our findings (23). It has been suggested that the increased blood loss under general anesthesia may be due to the effects of uterine muscle relaxants, which are not used in spinal anesthesia, leading to more bleeding in general anesthesia cases (24).

In surgeries where uterine contractions are not a factor, such as pelvic surgeries and hysterectomies, lower blood loss has been reported in the regional anesthesia group compared to the general anesthesia group (25), likely due to the vasodilatory effects of the anesthetic gases used in general anesthesia. In contrast, a study by Elhassan et al. found that the estimated blood

**Table 4.** Comparison of Pain Level Among Study Groups

Variables and Groups	Mean ± Standard Deviation	F	t	Mean Difference	P-Value
<b>Pain level at the moment of entering the ward</b>		3.16	-6.179	1.36	0.001
Spinal anesthesia	6.73 ± 0.96				
General anesthesia	8.10 ± 0.75				
<b>Pain level 6 hours after the operation</b>		0.987	-6.406	1.23	0.001
Spinal anesthesia	5.20 ± 0.99				
General anesthesia	6.43 ± 1.16				
<b>Pain level 12 hours after the operation</b>		1.841	-3.420	1.16	0.001
Spinal anesthesia	4.26 ± 1.20				
General anesthesia	5.43 ± 1.43				
<b>Pain level 18 hours after the operation</b>		39.472	-4.702	1.43	0.001
Spinal anesthesia	2.93 ± 0.63				
General anesthesia	6.35 ± 1.54				
<b>Pain level 24 hours after the operation</b>		3.185	-5.869	1.50	0.001
Spinal anesthesia	1.66 ± 0.85				
General anesthesia	3.16 ± 1.14				

loss was lower in the general anesthesia group compared to spinal anesthesia, which contradicts our results. This difference may be due to the retrospective nature of their study and associated bias (26). However, in a study by Aksoy et al., which compared hemoglobin and hematocrit levels before and after surgery in a prospective study on selected cesareans, they concluded that blood loss was higher in the general anesthesia group compared to the spinal anesthesia group (23), a finding that aligns with our study due to the similarity of the study populations.

Regarding postoperative pain, our study shows that patients undergoing cesarean sections under general anesthesia experience more pain compared to those under spinal anesthesia. This is because neuraxial anesthesia (spinal) is superior to injectable and systemic analgesia for pain relief (22). Additionally, neuraxial analgesia is associated with earlier bowel function recovery, earlier mobilization, and shorter hospital stays compared to systemic analgesia (27). Studies have demonstrated that post-cesarean pain is greater in patients who underwent general anesthesia compared to spinal anesthesia, which aligns with our findings (22). This is likely due to the longer duration of action of intrathecal spinal medications compared to the intravenous opioids used in general anesthesia, as well as the use of intravenous fentanyl in the spinal anesthesia group in our study. Previous cesarean section experiences may have also influenced the results.

Postoperative blood pressure drops are among the risk factors for postoperative shivering and can increase the risk of wound infection, oxygen consumption, and

patient discomfort. Therefore, blood pressure management is a critical factor. Our study found that systolic blood pressure was higher in the spinal anesthesia group compared to the general anesthesia group, while diastolic blood pressure was higher in the general anesthesia group at some postoperative hours. This may be due to the effects of adrenaline administered during spinal anesthesia and the volume of intraoperative fluids in the general anesthesia group, findings that are consistent with the study by Karami et al. (28). Chen et al. concluded that, in terms of hemodynamic parameters, general anesthesia is superior to spinal anesthesia, which somewhat aligns with our results, possibly due to better blood pressure management during surgery and the performance of systemic and intravenous anesthesia drugs (17).

Moreover, there was no statistically significant difference in Apgar scores at 1 and 5 minutes between the two groups (29). Data analysis by Kakarin supports this finding, indicating that there is no significant difference not only in terms of the mean Apgar score at the first and fifth minutes but also in the need for neonatal oxygen between the two groups (30). In contrast, a study by Gwanzura et al. found that the Apgar score was higher in the spinal anesthesia group compared to the general anesthesia group, which contradicts our findings (31). This discrepancy may be attributed to various factors, such as the significantly larger sample size in their study compared to ours. Additionally, our study focused exclusively on elective cesarean sections. Similarly, in the study by Al-Husban et al., no significant difference in Apgar scores between the




Figure 1. Status of systolic blood pressure upon entering the ward and at different moments of hospitalization




Figure 2. Diastolic blood pressure status upon entering the ward and at different moments of hospitalization

spinal and general anesthesia groups was observed, which is consistent with our results (26).

One of the strengths of this study is the randomization process used to allocate participants to different treatment groups without bias, ensuring that patients were randomly assigned to receive either general or spinal anesthesia for their cesarean section. Additionally, the duration of surgery and the amount of

oxytocin administered were standardized across both groups. However, a limitation of our study is the lack of homogeneity in the samples concerning body mass index (BMI), as the study by Zandi et al. found a correlation between BMI and blood loss (32). Other limitations include the small sample size, partly due to maternal dissatisfaction with early maternal-neonatal communication in the spinal anesthesia method, as well

as concerns such as depression in infants born under general anesthesia and the risk of maternal aspiration (33, 34). Furthermore, most participants in this study had prior cesarean section experiences, which may have served as a confounding factor in pain perception. Lastly, only elective cesarean sections were investigated. It is hoped that future studies will conduct more comprehensive investigations into variables such as cognitive dysfunction in mothers post-surgery to provide more accurate comparisons between these two anesthesia techniques.

### 5.1. Conclusions

In this study, systolic blood pressure after childbirth was found to be higher in women undergoing spinal anesthesia, while diastolic blood pressure was higher in women receiving general anesthesia. Additionally, the amount of intraoperative bleeding was greater in the general anesthesia group compared to the spinal anesthesia group. Postoperative pain was reported to be less in the spinal anesthesia group than in the general anesthesia group. There was no statistically significant difference in Apgar scores at 1 and 5 minutes between the two groups. Therefore, regional anesthesia emerged as the preferable option for elective cesarean sections, based on both hemodynamic parameters and pain assessment. The advantages of regional anesthesia for maternal and fetal outcomes were found to be superior to those of general anesthesia.

### Acknowledgements

This study is the result of a research project approved by Hamadan University of Medical Sciences. We would like to thank the Vice Chancellor for Research and Technology of Hamadan University of Medical Sciences for the financial support of this study under project number 14020115213.

### Footnotes

**Authors' Contribution:** Study concept and design, acquisition of data, analysis and interpretation of data, and drafting of the manuscript: Serveh Mohammadi; administrative, technical, and material support and study supervision: Behzad Imani; acquisition of data and analysis and interpretation of data: Sina ghasemi; critical revision of the manuscript for important intellectual content and statistical analysis: Behzad Shalmashi.

**Clinical Trial Registration Code:** IRCT20230225057532N1.

**Conflict of Interests Statement:** This study has no conflicts of interest for the authors.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** From an ethical standpoint, the present research was conducted after obtaining approval from the Ethics Committee of Hamedan University of Medical Sciences (IR.UMSHA.REC.1401.1042).

**Funding/Support:** This research was financially supported by the Vice Chancellor for Research and Technology of Hamadan University of Medical Sciences.

**Informed Consent:** Informed consent was obtained from each patient after providing comprehensive explanations about the general purpose of the research. Participants were free to withdraw from the study at any time if desired.

### References

1. Neuman MD, Lim G, Dinh C, Clapp JT. Anesthesiologists' Attitudes Toward Randomization to General versus Neuraxial Anesthesia for Cesarean Delivery. *Anesthesiology*. 2023;140(1):170-2. <https://doi.org/10.1097/aln.0000000000004791>.
2. Achmad M, Kadek Intan J, I Nyoman Santa W, Pontisomaya P. General Anesthesia for the Gravid Patient in the Emergency Operating Room at Sanglah General Hospital. *Int J Anesthetic Anesthesiol*. 2022;9(1):134. <https://doi.org/10.23937/2377-4630/1410134>.
3. Mhyre JM, Sultan P. General Anesthesia for Cesarean Delivery: Occasionally Essential but Best Avoided. *Anesthesiology*. 2019;130(6):864-6. <https://doi.org/10.1097/aln.0000000000002708>.
4. Laudenbach V, Mercier FJ, Roze JC, Larroque B, Ancel PY, Kaminski M, et al. Anaesthesia mode for caesarean section and mortality in very preterm infants: an epidemiologic study in the EPIPAGE cohort. *Int J Obstet Anesth*. 2009;18(2):142-9. [PubMed ID: 19195873]. <https://doi.org/10.1016/j.ijoa.2008.11.005>.
5. Betran AP, Torloni MR, Zhang JJ, Gülmezoglu AM. WHO Statement on Caesarean Section Rates. *BJOG*. 2015;123(5):667-70. <https://doi.org/10.1111/1471-0528.13526>.
6. Rafiei M, Saei Ghare M, Akbari M, Kiani F, Sayehmiri F, Sayehmiri K, et al. Prevalence, causes, and complications of cesarean delivery in Iran: A systematic review and meta-analysis. *Int J Reprod Biomed*. 2018;16(4):221-34. [PubMed ID: 29942930]. [PubMed Central ID: PMC6004597].
7. Shoeibi G, Salehi M, Atef Yekta R. [Evaluation of the reasons of fear in spinal anesthesia during cesarean section]. *Tehran Univ Med J*. 2021;79(7):527-32. FA.
8. Devroe S, Van de Velde M, Rex S. General anesthesia for caesarean section. *Curr Opin Anaesthesiol*. 2015;28(3):240-6. <https://doi.org/10.1097/aco.0000000000000185>.



9. Kim WH, Hur M, Park SK, Yoo S, Lim T, Yoon HK, et al. Comparison between general, spinal, epidural, and combined spinal-epidural anesthesia for cesarean delivery: a network meta-analysis. *Int J Obstet Anesth.* 2019;**37**:5-15. [PubMed ID: 30415797]. <https://doi.org/10.1016/j.ijoa.2018.09.012>.
10. Uppal V, McKeen DM. Strategies for prevention of spinal-associated hypotension during Cesarean delivery: Are we paying attention? *Can J Anaesth.* 2017;**64**(10):991-6. [PubMed ID: 28702819]. <https://doi.org/10.1007/s12630-017-0930-0>.
11. Eshghizadeh M, Basiri Moghaddam M, Mohammadpour A, Banihashemi ZS. [The Effect of Coffee Consumption on Post Dural Puncture Headache due to Spinal Anesthesia in Cesarean Section: A Randomized Clinical Trial]. *Qom Univ Med Sci J.* 2016;**9**(12):8-15. FA.
12. Patel A, Goudar SS, Geller SE, Kodkany BS, Edlavitch SA, Wagh K, et al. Drape estimation vs. visual assessment for estimating postpartum hemorrhage. *Int J Gynaecol Obstet.* 2006;**93**(3):220-4. [PubMed ID: 16626718]. <https://doi.org/10.1016/j.ijgo.2006.02.014>.
13. Adarvishi S, Dastoorpur M, Yazdani Nejad H, Mohammadi P, Kurd Nejad M, Ranjbarha R, et al. [Comparison of neonatal and maternal outcomes after cesarean delivery with general anesthesia and spinal anesthesia]. *Anesthesiol Pain.* 2021;**12**(1):44-54. FA.
14. O'Donnell CP, Kamlin CO, Davis PG, Carlin JB, Morley CJ. Interobserver variability of the 5-minute Apgar score. *J Pediatr.* 2006;**149**(4):486-9. [PubMed ID: 1701319]. <https://doi.org/10.1016/j.jpeds.2006.05.040>.
15. Ehrenstein V. Association of Apgar scores with death and neurologic disability. *Clin Epidemiol.* 2009;**1**:45-53. [PubMed ID: 20865086]. [PubMed Central ID: PMC2943160]. <https://doi.org/10.2147/clep.s4782>.
16. Tandon M, Singh A, Saluja V, Dhankhar M, Pandey CK, Jain P. Validation of a New "Objective Pain Score" Vs. "Numeric Rating Scale" For the Evaluation of Acute Pain: A Comparative Study. *Anesth Pain Med.* 2016;**6**(1). e32101. [PubMed ID: 27110530]. [PubMed Central ID: PMC4834447]. <https://doi.org/10.5812/aapm.32101>.
17. Chen Y, Liu W, Gong X, Cheng Q. Comparison of Effects of General Anesthesia and Combined Spinal/Epidural Anesthesia for Cesarean Delivery on Umbilical Cord Blood Gas Values: A Double-Blind, Randomized, Controlled Study. *Med Sci Monit.* 2019;**25**:5272-9. <https://doi.org/10.12659/msm.914160>.
18. Shetabi H, Jabalameli M, Talakkoub R, Mansuri A. [Effect of Intravenous Administration of Atropine-Neostigmine in Prevention of Headache after Spinal Anesthesia in Cesarean Section: A Randomized Clinical Trial]. *J Mazandaran Univ Med Sci.* 2023;**32**(216):88-96. FA.
19. Akbarzadeh S, Farhoodi R, Lyu T, Awais M, Zhao X, Farooq Abbasi S, et al. Evaluation of Apgar Scores and Non-Nutritive Sucking Skills in Infants Using a Novel Sensitized Non-Nutritive Sucking System. *42nd Annual International Conference of the IEEE Engineering in Medicine & Biology Society (EMBC).* 20-24 July 2020; Montreal, QC, Canada. 2020. p. 4282-5.
20. Yassini M, Karamnia M, Taghipoor R. [The Effect of Spinal versus General Anesthesia on Apgar score of Neonates after Caesarian Delivery]. *Iran J Anaesthesiol Crit Care.* 2015;**37**(1):58-65. FA.
21. Shin DW, Kim Y, Hong B, Yoon SH, Lim CS, Youn S. Effect of fentanyl on nausea and vomiting in cesarean section under spinal anesthesia: a randomized controlled study. *J Int Med Res.* 2019;**47**(10):4798-807. [PubMed ID: 31452417]. [PubMed Central ID: PMC6833428]. <https://doi.org/10.1177/0300060519869515>.
22. Ring L, Landau R, Delgado C. The Current Role of General Anesthesia for Cesarean Delivery. *Curr Anesthesiol Rep.* 2021;**11**(1):18-27. [PubMed ID: 33642943]. [PubMed Central ID: PMC7902754]. <https://doi.org/10.1007/s40140-021-00437-6>.
23. Aksoy H, Aksoy U, Yucel B, Ozyurt SS, Acmaz G, Babayigit MA, et al. Blood loss in elective cesarean section: is there a difference related to the type of anesthesia? A randomized prospective study. *J Turk Ger Gynecol Assoc.* 2015;**16**(3):158-63. [PubMed ID: 26401109]. [PubMed Central ID: PMC4560473]. <https://doi.org/10.5152/jtgga.2015.15034>.
24. Wong CA. General anesthesia is unacceptable for elective cesarean section. *Int J Obstet Anesth.* 2010;**19**(2):209-12. <https://doi.org/10.1016/j.ijoa.2009.10.002>.
25. Guay J. The effect of neuraxial blocks on surgical blood loss and blood transfusion requirements: a meta-analysis. *J Clin Anesth.* 2006;**18**(2):124-8. [PubMed ID: 16563330]. <https://doi.org/10.1016/j.jclinane.2005.08.013>.
26. Al-Husban N, Elmuhaseb MS, Al-Husban H, Nabhan M, Abuhaleweh H, Alkhatib YM, et al. Anesthesia for Cesarean Section: Retrospective Comparative Study. *Int J Womens Health.* 2021;**13**:141-52. [PubMed ID: 33564269]. [PubMed Central ID: PMC7866905]. <https://doi.org/10.2147/IJWH.S292434>.
27. Ghaffari S, Dehghanpisheh L, Tavakkoli F, Mahmoudi H. The Effect of Spinal versus General Anesthesia on Quality of Life in Women Undergoing Cesarean Delivery on Maternal Request. *Cureus.* 2018;**10**(12). e3715. <https://doi.org/10.7759/cureus.3715>.
28. Karami A, Khademi S, Fattahi Saravi Z, Jouybar R, Esmaeilnezhad Z, Asadpour E. [Comparison of Maternal and Neonatal Outcomes between Vaginal Delivery and Cesarean Section under General or spinal anesthesia- retrospective study]. *Iran J Obstet Gynecol Infertil.* 2020;**23**(4):15-23. FA. <https://doi.org/10.22038/ijogi.2020.16284>.
29. Kavak ZN, Başgöl A, Ceyhan N. Short-term outcome of newborn infants: spinal versus general anesthesia for elective cesarean section: A prospective randomized study. *Eur J Obstet Gynecol Reprod Biol.* 2001;**100**(1):50-4. [https://doi.org/10.1016/s0303-0215\(01\)00417-1](https://doi.org/10.1016/s0303-0215(01)00417-1).
30. Afolabi BB, Lesi FEA. Regional versus general anaesthesia for caesarean section. *Cochrane Database Syst Rev.* 2012;(10). CD004350. <https://doi.org/10.1002/14651858.CD004350.pub3>.
31. Gwanzura C, Gavi S, Mangiza M, Moyo FV, Lohman MC, Nhemachena T, et al. Effect of anesthesia administration method on apgar scores of infants born to women undergoing elective cesarean section. *BMC Anesthesiol.* 2023;**23**(1):142. <https://doi.org/10.1186/s12871-023-02098-w>.
32. Zandi S, Imani B, Mostafayi M, Rabie S. [Prevalence of Early Maternal Complications of Cesarean Section and its Relationship with Body Mass Index in Fatemieh Hospital of Hamadan]. *Pajouhan Sci J.* 2020;**18**(2):52-7. FA. <https://doi.org/10.52547/psj.18.2.52>.
33. Jenkins JG, Khan MM. Anaesthesia for Caesarean section: a survey in a UK region from 1992 to 2002. *Anaesthesia.* 2003;**58**(11):1114-8. <https://doi.org/10.1046/j.1365-2044.2003.03446.x>.
34. Gogarten W, Vandermeulen E, Van Aken H, Kozek S, Llaou JV, Samama CM. Regional anaesthesia and antithrombotic agents: recommendations of the European Society of Anaesthesiology. *Eur J Anaesthesiol.* 2010;**27**(12):999-1015. <https://doi.org/10.1097/EJA.0b013e32833f6f6f>.