



Intensity or Duration: That Is the Main Question of the Exercise Training Management in Stroke-Induced Neurological Deficit - A Systematic and Meta-analysis Study

Sadegh Abbasian^{1,*}, Mahsa Kargar Moghaddam², Amin Azimkhani³, Fariborz Ramezani¹, Mohsen Ghotbi¹

¹ Assistant Professor, Department of Physical Education, Farhangian University, Tehran, Iran

² Ph.D. Student, Department of Exercise Physiology, Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran

³ Assistant Professor, Imam Reza International University, Mashhad, Iran

*Corresponding Author: Assistant Professor, Department of Physical Education, Farhangian University, P. O. Box.: 14665889, Tehran, Iran. Email: s.abbasian@cfu.ac.ir

Received: 19 May, 2024; Revised: 30 June, 2024; Accepted: 31 August, 2024

Abstract

Background: The neurological assessment plays a crucial role in evaluating stroke patients and is a skill that has been practiced by physicians for a long time, and further refined by practitioners. The purpose of the current study was to systematically review and synthesize the evidence-based neurological assessment of the different methods of treadmill-related training protocols in stroke patients.

Evidence Acquisition: In this study, the appropriate studies were documented (N = 1270) using PubMed as the sole electronic database. Moreover, subgroup studies were provided based on training intensity and duration, categorized as low intensity-low duration, low intensity-high duration, high intensity-low duration, and high intensity-high duration. Training intensities were standardized to meters per second (m/s), and the training duration was calculated in terms of minutes per session, the number of sessions per week, and the total number of training weeks, which were then converted into minutes. The assessment of neurological recovery following stroke was conducted using standard neurological assessment scales.

Results: This meta-analysis in humans found a notably positive standardized mean difference in protocols with high intensity and high duration (1.792; 95% CI = 0.566 to 3.02).

Conclusions: In conclusion, the current meta-analysis suggests that practitioners using the high intensity-high duration protocol should regularly monitor the neurological outcomes, as they may be negatively impacted by the intensity and duration of the protocol.

Keywords: Duration, Intensity, Neurological Deficits, Stroke, Training

1. Background

Stroke is a key contributor to patient mortality and is one of the main causes of disabilities globally (1, 2). In previous studies, it has been established that eye movement abnormalities, neurological deficits, arm and leg weakness, arm and leg paresis, gait ataxia, and subjective facial asymmetry are prominent indicators of post-stroke outcomes (3). Neurological assessment plays a crucial role in evaluating stroke patients and is a skill that has been practiced and refined by physicians for many years (4). Consequently, numerous post-stroke restorative modalities are under investigation,

including brain stimulation, growth factors, monoclonal antibodies, cell-based therapies, drugs, robotic and telehealth devices, and activity-based therapies, all of which have shown benefits in neural repair and cognitive function remapping after stroke (5).

Exercise training is one of the stroke-dependent modalities with evidence suggesting its potential to modulate stroke-associated impairments, particularly through enhanced cerebral blood flow in infarct areas of the brain (6). Recovery of gait is a primary objective in neurological rehabilitation (7), and treadmill exercise is a widely recognized and well-studied training method

in stroke recovery. Treadmill training has been shown to reduce post-stroke deficiencies (8). However, the effectiveness of neurological rehabilitation is strongly influenced by the intensity and duration of the training (9).

There is concern about the appropriate intensity and duration of treadmill exercise training, as higher intensity may lead to poorer outcomes following a stroke (10). While treadmill training is a standard rehabilitation practice, the optimal roles of intensity and duration in human studies remain unclear (8). Previous literature has underscored the importance of exercise intensity in stroke rehabilitation programs. Some studies suggest that gradually increasing exercise intensity may promote greater neurological recovery than low- or high-intensity regimens (11-13). Conversely, other studies indicate that high-intensity protocols result in superior neurological recovery compared to low-intensity protocols (12, 14). While low-intensity strategies may have a reduced effect on recovery, some evidence shows no significant differences in motor function recovery between low- and high-intensity training protocols (8).

In terms of duration, some studies report that shorter durations yield better neurological outcomes (14-16), while others find that longer durations are more beneficial (17-19). However, there is insufficient evidence regarding the effect of treadmill protocol intensity and duration on stroke-induced neurological deficits. The objective of this meta-analysis was to determine the effects of treadmill training on post-stroke neurological deficits in human studies. The findings may help guide physicians, healthcare professionals, and researchers in selecting the optimal intensity and duration of training protocols for stroke patients.

2. Evidence Acquisition

2.1. Definitions of Variables

In line with the criteria outlined by the World Health Organization (WHO), stroke is defined as the abrupt onset of clinical manifestations indicating focal (or global) dysfunction of cerebral function, lasting over 24 hours or leading to death, with no apparent cause other than of vascular origin (20). Additionally, a reduction of one or more points in the overall neurological scale score from entry to hospitalization was used as the criterion for defining neurological deficit (21).

In this meta-analysis, we included human neurological scales, such as assessments of eye movements, closed-eye balance, and the Modified Ashworth Scale (MAS). Therapeutic exercise or exercise

therapy refers to a prescribed and supervised program in which patients perform repeated voluntary dynamic movements or static muscle contractions. These exercises may target specific body regions or involve the whole body, either with or without external resistance. The goal is to relieve symptoms, control inflammation (if present), improve range of motion, muscular endurance, strength, and power, and ultimately assist patients in returning to their daily and recreational activities through training (22). For the present investigation, we included the use of treadmills with body-weight support as part of the therapeutic exercise.

According to the American College of Sports Medicine (ACSM) and the American Heart Association (AHA), exercise-related adaptations for stroke survivors are expected to manifest after two to six weeks, with three sessions per week and approximately 20 - 60 minutes per session (23, 24). Therefore, in line with previous investigations, the duration of treadmill training was calculated based on the total minutes allocated for therapeutic exercise in each session, week, and training week(s). The training protocol was considered as high duration (more than 500 minutes) if the total training time exceeded 500 minutes; otherwise, it was categorized as low duration (less than 500 minutes) (8).

The AHA practical guidelines on exercise recommendations for stroke survivors suggest that intensity training should be performed at a level of 40% - 70% of peak oxygen uptake or heart rate reserve, 50% - 80% of maximal heart rate, or a perceived exertion level of 11 - 14 on the 15-category Rated Perceived Exertion (RPE) scale (24). However, due to challenges in practical application, several studies have demonstrated that a training intensity of 2 km/h (or 0.6 m/s) is more feasible for stroke patients (8, 25, 26). In line with previous studies, we defined treadmill training intensity as the speed of treadmill training during exercise therapy. Training intensity was categorized as high intensity (more than 0.6 m/s) if the intensity exceeded 0.6 m/s, otherwise it was classified as low intensity (less than 0.6 m/s) (8). Training difference refers to the difference in intensity and duration between the experimental group and the control group, or the difference compared to pre-test values (8, 27).

2.2. Selection Process of Relevant Studies

We selected appropriate stroke-related articles for inclusion based on predefined criteria, such as study design, training intensity, duration conditions, and outcome measures, from January 1980 to March 2020 using the PubMed search engine. Headlines and

keywords related to medical topics were sought out from different platforms (8). Our previous data pool, comprising studies up to March 2020, was further updated (Box 1).

2.3. Research Entry Criteria

From our search results, we identified 87 studies that were potentially suitable, with six meeting our inclusion criteria (Table 1). Among these, four studies involved human participants. We also examined studies that included one or more independent variables (such as intensity and/or duration of treadmill training) and used a neurological scale as the dependent variable, as well as studies that featured multiple independent and dependent variables. Additionally, the amount of body-weight support (BWS) used during treadmill training (10% to 40% of body mass) was considered a crucial variable in the studies.

Our analysis focused exclusively on studies that clearly explained the training intensity using measurements in meters per minute (m/min), centimeters per second (cm/s), meters per second (m/s), or kilometers per hour (km/h), all of which were standardized to m/s for human experiments. The training duration was determined by calculating the minutes per session, the number of sessions per week, and the total number of training weeks, which were then converted into total minutes (30). Neurological recovery following a stroke was assessed using standard neurological assessment scales, though the specific names of these scales were not disclosed by the authors (Table 1) (30, 31).

2.4. Confounding Factors

In our study, potential confounding factors included the age difference at the time of stroke as well as the functional status prior to the occurrence of stroke. Additionally, the amount of BWS used during treadmill training (10% to 40% of body mass) was considered another confounding factor.

2.5. Research Biases

Similar to other meta-analyses, the present study was subject to several potential biases, such as training intensity, training duration (\geq two weeks), clarity of the training protocol, training safety (e.g., heart rate monitoring or BWS), training overload, and expression of neurological scales. Studies that addressed or covered at least half of the above-mentioned biases were included in the data pooling (32, 33).

2.6. Research Data Extraction

The author succinctly outlined the findings of human studies, including key details such as: (1) the first author's name and publication year; (2) applied data for every study, such as the number of subjects, sessions, days, and weeks of training; (3) a description of treatment specifics, including the intensity and duration of treadmill training during the entire therapeutic exercise session, which were then converted into meters per second (or per minute) and minutes, respectively; (4) classification of the therapeutic protocols into categories such as: (a) low intensity; (b) low duration, low intensity; (c) high duration, high intensity; (d) low duration, and high intensity - high duration, based on studies conducted on humans (34-37); (5) presentation of the study findings as mean \pm standard deviation (SD).

When data was unavailable for pooling, efforts were made to calculate the necessary information or to contact the authors for additional data; otherwise, studies were excluded due to lack of accessible data. Additionally, we contacted the corresponding authors of conference abstracts for full-paper publications, where possible, and consulted professionals in the relevant field for further insights.

2.7. Classification of Research Studies

Before commencing therapeutic exercise, a comprehensive appraisal is necessary to assess neurological deficits in human studies (7). In this meta-analysis, we predominantly employed neurological scales for subgroup categorizations. The neurological scales included assessments of eye movements, closed-eye balance, and the MAS for human studies. Subgroup classifications for human studies were determined based on previous research and are as follows (8, 34-37):

- (1) Low intensity (0.6 m/s or less) - low duration (500 minutes or less) training protocol
- (2) Low intensity (0.6 m/s or less) - high duration (more than 500 minutes) training protocol
- (3) High intensity (more than 0.6 m/s) - low duration (500 minutes or less) training protocol
- (4) High intensity (more than 0.6 m/s) - high duration (more than 500 minutes) training protocol

According to the neurological scales, we included four out of 28 studies in this meta-analysis. The subgroups of human studies included two subgroups categorized as low intensity-low duration protocol (14, 28), with one subgroup being excluded. Three subgroups were classified as low intensity-high

Box 1. Details of Search Strategy for Medical Topic Headlines of Stroke Studies

Details
 ("exercise"[MeSH Terms] OR "exercise"[All Fields]) AND ("stroke"[MeSH Terms] OR "stroke"[All Fields]) ("stroke"[MeSH Terms] OR "stroke"[All Fields]) AND ("cerebrum"[MeSH Terms] OR "cerebrum"[All Fields] OR "cerebral"[All Fields] OR "brain"[MeSH Terms] OR "brain"[All Fields]) AND ("exercise"[MeSH Terms] OR "exercise"[All Fields]) ("stroke"[Title] AND cerebral[Title]) AND exercise[Title] ("stroke[Abstract] AND cerebral[Abstract]) AND exercise[Abstract] ("stroke[Abstract] AND exercise[Abstract]) AND cerebral[Abstract] AND brain[Abstract]) AND physical[Abstract] AND training[Abstract]) ("Post-ischemic[All Fields] AND ("exercise"[MeSH Terms] OR "exercise"[All Fields]) ("exercise"[MeSH Terms] OR "exercise"[All Fields]) AND ("physical examination"[MeSH Terms] OR "physical"[All Fields] AND "examination"[All Fields]) OR "physical examination"[All Fields] OR "physical"[All Fields]) AND ("cerebrum"[MeSH Terms] OR "cerebrum"[All Fields] OR "cerebral"[All Fields] OR "brain"[MeSH Terms] OR "brain"[All Fields]) AND ("stroke"[MeSH Terms] OR "stroke"[All Fields]) AND ("brain"[MeSH Terms] OR "brain"[All Fields]) ("post-exercise[All Fields] AND ("cerebrum"[MeSH Terms] OR "cerebrum"[All Fields] OR "cerebral"[All Fields] OR "brain"[MeSH Terms] OR "brain"[All Fields]) AND ("ischaemia"[All Fields] OR "ischemia"[MeSH Terms] OR "ischemia"[All Fields]) "Hemiparetic[All Fields] AND treadmill[All Fields] AND ("1980/01/01"[PDAT] : "2018/03/01"[PDAT]).

Table 1. Characteristics and Biases of Stroke-Associated Neurological Deficit Studies on Stroke Patients

Study/Subgroup ID	Year	Speed	Intensity		Duration		Biases ^a						
			≤ 0.6 m.s ⁻¹	> 0.6 m.s ⁻¹	≤ 500 min	> 500 min	1	2	3	4	5	6	
Dumont et al. (28)	2015	√	√		√			+	+	-	-	-	+
Calabro et al. (29)	2015	×	√			√		-	+	+	+	+	+
Liu et al. (14)	2014	√	√	√	√	√		+	+	+	+	+	+
Freivogel et al. (17)	2008	√				√		+	+	+	+	+	+

Abbreviations: N/A, not applicable; BWS, body-weight support.

^a The meta-analysis biases were consisted of training intensity, duration, protocol expression, safety (heart rate monitoring or BWS), overload, and neurological scale expression, respectively.

duration protocol (14, 17, 29), and one subgroup was categorized as high intensity-high duration protocol (14). Notably, there was no study or subgroup on humans that was categorized as high intensity-low duration training protocol (Figure 1).

2.8. Statistical Analysis

Quantitative data were recorded as the mean ± SD following the specified criteria. Continuous data were then processed using the "Metan" command (38, 39) to calculate the mean, SD, and sample size for each group. Standard mean differences (SMD) with 95% confidence intervals (CI) and weight percentages for each study or subgroup were subsequently assessed using the "Metan" command, as described in prior research (8).

Egger's funnel plot and Egger's test, using the "Metabias" command, were applied to detect publication bias, following the method outlined by Egger et al. (40). The I-squared statistic (I²) was used to evaluate the level of heterogeneity, based on Higgins et al.'s recommendations (41). To account for variability among multiple studies, a random effects model was preferred over a fixed effects model (42).

All statistical analyses were performed using Stata 12 (StataCorp, USA), and study figures were generated using GraphPad Prism 6.01.

3. Results

Figure 2 illustrates the SMD with 95% CI for neurological rehabilitation in the dominant training protocol categories. Four studies (N = 112) examined the use of a motorized treadmill for walking, which included patients undergoing treadmill rehabilitation protocols (14, 17, 28, 29). One study (28) did not have a clear post-stroke rehabilitation SMD. The CIs for the SMD of both low intensity-low duration subgroups and high intensity-high duration subgroups were greater than zero. However, the CI for the SMD of the low intensity-high duration subgroups was significantly less than zero. Additionally, one subgroup from the low intensity-low duration protocol was excluded by Stata from the meta-analysis due to zero weight.

The six human studies included in the analysis exhibited a high level of heterogeneity (I² = 96.6%; P < 0.0001). While a negative SMD was observed, both the low intensity-low duration and low intensity-high duration protocols resulted in a reduction in neurological deficit, though the difference was not statistically significant (P = 0.277 and P = 0.615, respectively). Conversely, a positive SMD was observed in the high intensity-high duration protocol, indicating an increase in neurological deficit with higher intensity (SMD = 1.792; 95% CI = 0.566 to 3.02; P = 0.004). Nevertheless, there were no significant differences in SMD across the various subgroups in the study (P = 0.109; Figure 2). Furthermore, the Egger's funnel plot




Figure 1. The meta-analysis data extraction flowchart

and test with a 95% confidence interval indicated no publication biases in any of the study subgroups ($t = 2.09, P = 0.104$).

4. Discussion

Regarding recent literature, prior research has increasingly demonstrated that exercise training is being recommended as a critical component of stroke rehabilitation programs, given the growing body of evidence supporting its benefits in mitigating neurological and functional impairments following stroke (43). The outcomes of our earlier study strongly suggest that treadmill training, regardless of intensity or duration, results in significantly better motor function rehabilitation compared to the absence of training (8). However, the specific impact of exercise training intensity and duration on neurological recovery in stroke patients remains unclear.

To address this, we evaluated whether training duration, intensity, or a combination of both influenced the primary neurological outcomes in stroke-related human studies. The current meta-analysis revealed that approximately 79.4% of the combined studies did not

provide precise details on treadmill training protocols, thus failing to adequately address biases in the analysis. In contrast, only 20.6% of the studies accounted for all six biases considered in the meta-analysis. Furthermore, the meta-analysis highlighted a significant decline in the number of stroke-related treadmill training investigations over the last three years.

Our research findings indicate that 2 out of 7 interventions (28%) provided significant evidence of beneficial effects on neurological and functional recovery, based on the results of our meta-analyses.

A significant number of stroke-related training studies indicate that treadmill training yields comparable results to conventional therapy, suggesting that similar outcomes can be achieved with the control intervention, without conflicting findings. It is important to note that while adverse SMDs in neurological deficit were observed in the low intensity-low duration and low intensity-high duration protocols, these differences were not statistically significant. Only one study (29) reported a negative SMD, indicating that the low intensity-high duration protocol had a positive impact on neurological deficit. Despite this, there was no significant difference in the overall SMD across the




Figure 2. Recovery of neurological deficit following the treadmill training strategies in stroke patients. In this meta-analysis, we did not see a significant reduction in the neurological deficit after the treadmill training interventions, but there was nonsignificant decline in the neurological deficit following the low intensity-high duration and low intensity-low duration protocols ($P > 0.05$). However, the high intensity-high duration intervention significantly decreased the neurological deficit in stroke patients ($P = 0.0001$). Altogether, the current meta-analysis displayed that, regardless of training intensity and duration, treadmill training did not have a significant effect on recovery of the neurological deficit ($P > 0.05$).

subgroups. On the other hand, the high intensity-high duration protocol showed a positive SMD, suggesting

that as treadmill intensity increases, neurological deficit tends to worsen.

Furthermore, it has been shown that intensity plays a key role in inducing stress. Intense training leads to elevated stress levels, which are associated with reduced hippocampal brain-derived neurotrophic factor (BDNF) levels, a marker of neural activity (2). Previous studies have demonstrated that exercise training can enhance neurological recovery by boosting neural progenitor cell levels through the activation of the IGF-1/Akt and BDNF signaling pathways (2, 44, 45). Therefore, practitioners opting for the high intensity-high duration protocol should consistently monitor neurological outcomes that may be negatively impacted by the protocol.

Despite the findings, this meta-analysis indicated a lack of statistical power or summary effect sizes (SEs) for all neurological outcomes (SEs = 0.496), emphasizing the need for additional human studies. While a notably negative SMD was not detected in low intensity training protocols, these methods still showed positive impacts on motor functional recovery. Although the meta-analysis was carried out meticulously, it faced methodological limitations such as language restrictions and missing data (33, 46, 47). Additionally, the assessment of subgroups or sub-studies is limited by the observational nature of the included research, and caution is advised in interpreting these findings until stronger evidence is available (33, 39-41, 46, 47). Controlled variables differed between studies, and future human studies should further assess neurological outcomes following treadmill training.

Given the limited number of subgroups (six) that examined neurological recovery after treadmill training, further studies are needed to validate and expand upon these findings.

4.1. Conclusions

In conclusion, our findings deliver several key messages. First, this meta-analysis includes studies with the least biases, strengthening the reliability of the results. Furthermore, it confirms the significant impact of treadmill training protocols in enhancing neurological recovery and reducing the risk of disability in stroke patients. Additionally, the current meta-analysis suggests that practitioners using the high intensity-high duration protocol should regularly monitor neurological outcomes, as this protocol may negatively impact recovery in some cases. Lastly, to make it easier to achieve the functional and neurological improvements observed in these meta-analyses through regular exercise, it is essential to specifically identify the training intensity and duration

that show strong evidence of benefiting stroke patients. Furthermore, determining which type of study provides valid evidence for this is crucial for optimizing stroke rehabilitation protocols.

Footnotes

Authors' Contribution: Sadegh Abbasian and Mahsa Kargar Moghadam: Data collection and writing the first draft; Amin Azimkhani: Supervisor; Fariborz Ramezani: Consultant; Mohsen Ghotbi: Consultant.

Conflict of Interests Statement: The authors declared they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Funding/Support: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

References

- Agulla J, Brea D, Campos F, Sobrino T, Argibay B, Al-Soufi W, et al. In vivo theranostics at the peri-infarct region in cerebral ischemia. *Theranostics*. 2013;4(1):90-105. [PubMed ID: 24396517]. [PubMed Central ID: PMC3881229]. <https://doi.org/10.7150/thno.7088>.
- Sun J, Ke Z, Yip SP, Hu XL, Zheng XX, Tong KY. Gradually increased training intensity benefits rehabilitation outcome after stroke by BDNF upregulation and stress suppression. *Biomed Res Int*. 2014;2014:925762. [PubMed ID: 25045713]. [PubMed Central ID: PMC4090448]. <https://doi.org/10.1155/2014/925762>.
- Yew KS, Cheng EM. Diagnosis of acute stroke. *Am Fam Physician*. 2015;91(8):528-36. [PubMed ID: 25884860].
- Adams HP, Lyden P. Chapter 48 Assessment of a patient with stroke: neurological examination and clinical rating scales. In: Fisher M, editor. *Stroke Part III: Investigation and Management. Handbook of Clinical Neurology*. Vol. 94. Amsterdam: Elsevier; 2008. p. 971-1009. [https://doi.org/10.1016/S0072-9752\(08\)94048-3](https://doi.org/10.1016/S0072-9752(08)94048-3).
- Cramer SC. Treatments to Promote Neural Repair after Stroke. *J Stroke*. 2018;20(1):57-70. [PubMed ID: 29402069]. [PubMed Central ID: PMC5836581]. <https://doi.org/10.5853/jos.2017.02796>.
- Zhang P, Yu H, Zhou N, Zhang J, Wu Y, Zhang Y, et al. Early exercise improves cerebral blood flow through increased angiogenesis in experimental stroke rat model. *J Neuroeng Rehabil*. 2013;10:43. [PubMed ID: 23622352]. [PubMed Central ID: PMC3648391]. <https://doi.org/10.1186/1743-0003-10-43>.
- Mauritz KH. Gait training in hemiplegia. *Eur J Neurol*. 2002;9 Suppl 1:23-9. discussion 53-61. [PubMed ID: 11918646]. <https://doi.org/10.1046/j.1468-1331.2002.00905i023.x>.
- Abbasian S, Rastegar M. Is the Intensity or Duration of Treadmill Training Important for Stroke Patients? A Meta-Analysis. *J Stroke Cerebrovasc Dis*. 2018;27(1):32-43. [PubMed ID: 29108807]. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2017.09.061>.

9. Kwakkel G, Wagenaar RC, Twisk JW, Lankhorst GJ, Koetsier JC. Intensity of leg and arm training after primary middle-cerebral-artery stroke: a randomised trial. *Lancet*. 1999;**354**(9174):191-6. [PubMed ID: 10421300]. [https://doi.org/10.1016/S0140-6736\(98\)09477-X](https://doi.org/10.1016/S0140-6736(98)09477-X).
10. Dromerick AW, Lang CE, Powers WJ, Wagner JM, Sahrman SA, Videen TO, et al. Very early constraint-induced movement therapy (VECTORS): phase II trial results. *Stroke*. Philadelphia, PA: Lippincott Williams & Wilkins; 2007.
11. Matsuda F, Sakakima H, Yoshida Y. The effects of early exercise on brain damage and recovery after focal cerebral infarction in rats. *Acta Physiol (Oxf)*. 2011;**201**(2):275-87. [PubMed ID: 20726846]. [PubMed Central ID: PMC3045711]. <https://doi.org/10.1111/j.1748-1708.2010.02174.x>.
12. Rezaei R, Nourshahi M, Khodaghali F, Haghighparast A, Nasoohi S, Bigdeli M, et al. Differential impact of treadmill training on stroke-induced neurological disorders. *Brain Inj*. 2017;**31**(13-14):1910-7. [PubMed ID: 28898133]. <https://doi.org/10.1080/02699052.2017.1346287>.
13. Zhang A, Bai Y, Hu Y, Zhang F, Wu Y, Wang Y, et al. The effects of exercise intensity on p-NR2B expression in cerebral ischemic rats. *Can J Neurol Sci*. 2012;**39**(5):613-8. [PubMed ID: 22931702]. <https://doi.org/10.1017/s0317167100015341>.
14. Liu P, Wang Y, Hu H, Mao Y, Huang D, Li L. Change of muscle architecture following body weight support treadmill training for persons after subacute stroke: evidence from ultrasonography. *Biomed Res Int*. 2014;**2014**:270676. [PubMed ID: 24783198]. [PubMed Central ID: PMC3982255]. <https://doi.org/10.1155/2014/270676>.
15. Chen CC, Chang MW, Chang CP, Chan SC, Chang WY, Yang CL, et al. A forced running wheel system with a microcontroller that provides high-intensity exercise training in an animal ischemic stroke model. *Braz J Med Biol Res*. 2014;**47**(10):858-68. [PubMed ID: 25140816]. [PubMed Central ID: PMC4181221]. <https://doi.org/10.1590/1414-431x20143754>.
16. Curry A, Guo M, Patel R, Liebelt B, Sprague S, Lai Q, et al. Exercise pre-conditioning reduces brain inflammation in stroke via tumor necrosis factor- α , extracellular signal-regulated kinase 1/2 and matrix metalloproteinase-9 activity. *Neurol Res*. 2010;**32**(7):756-62. [PubMed ID: 19682410]. <https://doi.org/10.1179/174313209X459101>.
17. Freivogel S, Mehrholz J, Husak-Sotomayor T, Schmalohr D. Gait training with the newly developed 'LokoHelp'-system is feasible for non-ambulatory patients after stroke, spinal cord and brain injury. A feasibility study. *Brain Inj*. 2008;**22**(7-8):625-32. [PubMed ID: 18568717]. <https://doi.org/10.1080/02699050801941771>.
18. Wang X, Zhang M, Feng R, Li WB, Ren SQ, Zhang F. Exercise pre-conditioning alleviates brain damage via excitatory amino acid transporter 2 and extracellular signal-regulated kinase 1/2 following ischemic stroke in rats. *Mol Med Rep*. 2015;**11**(2):1523-7. [PubMed ID: 25370789]. <https://doi.org/10.3892/mmr.2014.2834>.
19. Wang X, Zhang M, Yang SD, Li WB, Ren SQ, Zhang J, et al. Pre-ischemic treadmill training alleviates brain damage via GLT-1-mediated signal pathway after ischemic stroke in rats. *Neuroscience*. 2014;**274**:393-402. [PubMed ID: 24907601]. <https://doi.org/10.1016/j.neuroscience.2014.05.053>.
20. Aho K, Harmsen P, Hatano S, Marquardsen J, Smirnov VE, Strasser T. Cerebrovascular disease in the community: results of a WHO collaborative study. *Bull World Health Organ*. 1980;**58**(1):113-30. [PubMed ID: 6966542]. [PubMed Central ID: PMC2395897].
21. Toni D, Fiorelli M, Gentile M, Bastianello S, Sacchetti ML, Argentino C, et al. Progressing neurological deficit secondary to acute ischemic stroke. A study on predictability, pathogenesis, and prognosis. *Arch Neurol*. 1995;**52**(7):670-5. [PubMed ID: 7619022]. <https://doi.org/10.1001/archneur.1995.00540310040014>.
22. Sebag JD. 49 - Therapeutic Exercises. In: Lennard TA, Walkowski S, Singla AK, Vivian DG, editors. *Pain Procedures in Clinical Practice (Third Edition)*. Saint Louis: Hanley & Belfus; 2011. p. 597-617. <https://doi.org/10.1016/B978-1-4160-3779-8.10049-1>.
23. Whaley MH, Brubaker PH, Otto RM, Armstrong LE; American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*. Philadelphia, PA: Lippincott Williams & Wilkins; 2006.
24. Gordon NF, Gulanic M, Costa F, Fletcher G, Franklin BA, Roth EJ, et al. Physical Activity and Exercise Recommendations for Stroke Survivors: An American Heart Association Scientific Statement From the Council on Clinical Cardiology, Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention; the Council on Cardiovascular Nursing; the Council on Nutrition, Physical Activity, and Metabolism; and the Stroke Council. 2004;**35**(5):1230-40. <https://doi.org/10.1161/01.str.0000127303.19261.19>.
25. McCain KJ, Pollo FE, Baum BS, Coleman SC, Baker S, Smith PS. Locomotor treadmill training with partial body-weight support before overground gait in adults with acute stroke: a pilot study. *Arch Phys Med Rehabil*. 2008;**89**(4):684-91. [PubMed ID: 18373999]. <https://doi.org/10.1016/j.apmr.2007.09.050>.
26. Yamada S, Tomida K, Tanino G, Suzuki A, Kawakami K, Kubota S, et al. How effective is the early fast treadmill gait speed training for stroke patients at the 2nd week after admission: comparison with comfortable gait speed at the 6th week. *J Phys Ther Sci*. 2015;**27**(4):1247-50. [PubMed ID: 25995599]. [PubMed Central ID: PMC4434020]. <https://doi.org/10.1589/jpts.27.1247>.
27. Kwakkel G, van Peppen R, Wagenaar RC, Wood Dauphinee S, Richards C, Ashburn A, et al. Effects of augmented exercise therapy time after stroke: a meta-analysis. *Stroke*. 2004;**35**(11):2529-39. [PubMed ID: 15472114]. <https://doi.org/10.1161/01.STR.00000143153.76460.7d>.
28. Dumont AJ, Araujo MC, Lazzari RD, Santos CA, Carvalho DB, Franco de Moura RC, et al. Effects of a single session of transcranial direct current stimulation on static balance in a patient with hemiparesis: a case study. *J Phys Ther Sci*. 2015;**27**(3):955-8. [PubMed ID: 25931768]. [PubMed Central ID: PMC4395752]. <https://doi.org/10.1589/jpts.27.955>.
29. Calabro RS, De Cola MC, Leo A, Reitano S, Balletta T, Trombetta G, et al. Robotic neurorehabilitation in patients with chronic stroke: psychological well-being beyond motor improvement. *Int J Rehabil Res*. 2015;**38**(3):219-25. [PubMed ID: 25816006]. <https://doi.org/10.1097/MRR.0000000000000114>.
30. Campos F, Sobrino R, Ramos-Cabrera P, Argibay B, Agulla J, Perez-Mato M, et al. Neuroprotection by glutamate oxaloacetate transaminase in ischemic stroke: an experimental study. *J Cereb Blood Flow Metab*. 2011;**31**(6):1378-86. [PubMed ID: 21266983]. [PubMed Central ID: PMC3130324]. <https://doi.org/10.1038/jcbfm.2011.3>.
31. Schaar KL, Breneman MM, Savitz SI. Functional assessments in the rodent stroke model. *Exp Transl Stroke Med*. 2010;**2**(1):13. [PubMed ID: 20642841]. [PubMed Central ID: PMC2915950]. <https://doi.org/10.1186/2040-7378-2-13>.
32. Van Peppen RP, Kwakkel G, Wood-Dauphinee S, Hendriks HJ, Van der Wees PJ, Dekker J. The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clin Rehabil*. 2004;**18**(8):833-62. [PubMed ID: 15609840]. <https://doi.org/10.1191/0269215504cr8430a>.
33. Veerbeek JM, van Wegen E, van Peppen R, van der Wees PJ, Hendriks E, Rietberg M, et al. What is the evidence for physical therapy poststroke? A systematic review and meta-analysis. *PLoS One*. 2014;**9**(2). e87987. [PubMed ID: 24505342]. [PubMed Central ID: PMC3913786]. <https://doi.org/10.1371/journal.pone.0087987>.
34. Westlake KP, Patten C. Pilot study of Lokomat versus manual-assisted treadmill training for locomotor recovery post-stroke. *J Neuroeng Rehabil*. 2009;**6**:18. [PubMed ID: 19523207]. [PubMed Central ID: PMC2708184]. <https://doi.org/10.1186/1743-0003-6-18>.

35. Moore JL, Roth EJ, Killian C, Hornby TG. Locomotor training improves daily stepping activity and gait efficiency in individuals poststroke who have reached a "plateau" in recovery. *Stroke*. 2010;**41**(1):129-35. [PubMed ID: 19910547]. <https://doi.org/10.1161/STROKEAHA.109.563247>.
36. Sullivan KJ, Knowlton BJ, Dobkin BH. Step training with body weight support: effect of treadmill speed and practice paradigms on poststroke locomotor recovery. *Arch Phys Med Rehabil*. 2002;**83**(5):683-91. [PubMed ID: 11994808]. <https://doi.org/10.1053/apmr.2002.32488>.
37. Hesse S, Bertelt C, Jahnke MT, Schaffrin A, Baake P, Malezic M, et al. Treadmill training with partial body weight support compared with physiotherapy in nonambulatory hemiparetic patients. *Stroke*. 1995;**26**(6):976-81. [PubMed ID: 7762049]. <https://doi.org/10.1161/01.str.26.6.976>.
38. Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health*. 2014;**72**(1):39. [PubMed ID: 25810908]. [PubMed Central ID: PMC4373114]. <https://doi.org/10.1186/2049-3258-72-39>.
39. Harris RJ, Deeks JJ, Altman DG, Bradburn MJ, Harbord RM, Sterne JA. Metan: Fixed- and Random-Effects Meta-Analysis. *Stata J*. 2008;**8**(1):3-28. <https://doi.org/10.1177/1536867x0800800102>.
40. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;**315**(7109):629-34. [PubMed ID: 9310563]. [PubMed Central ID: PMC2127453]. <https://doi.org/10.1136/bmj.315.7109.629>.
41. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;**327**(7414):557-60. [PubMed ID: 12958120]. [PubMed Central ID: PMC192859]. <https://doi.org/10.1136/bmj.327.7414.557>.
42. Wang WW, Xie CL, Lu L, Zheng GQ. A systematic review and meta-analysis of Baihui (GV20)-based scalp acupuncture in experimental ischemic stroke. *Sci Rep*. 2014;**4**:3981. [PubMed ID: 24496233]. [PubMed Central ID: PMC5379241]. <https://doi.org/10.1038/srep03981>.
43. Leon AS, Franklin BA, Costa F, Balady GJ, Berra KA, Stewart KJ, et al. Cardiac rehabilitation and secondary prevention of coronary heart disease: an American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. 2005;**111**(3):369-76. [PubMed ID: 15668354]. <https://doi.org/10.1161/01.CIR.0000151788.08740.5C>.
44. Chen BY, Wang X, Wang ZY, Wang YZ, Chen LW, Luo ZJ. Brain-derived neurotrophic factor stimulates proliferation and differentiation of neural stem cells, possibly by triggering the Wnt/beta-catenin signaling pathway. *J Neurosci Res*. 2013;**91**(1):30-41. [PubMed ID: 23023811]. <https://doi.org/10.1002/jnr.23138>.
45. Zheng HQ, Zhang LY, Luo J, Li LL, Li M, Zhang Q, et al. Physical exercise promotes recovery of neurological function after ischemic stroke in rats. *Int J Mol Sci*. 2014;**15**(6):10974-88. [PubMed ID: 24945308]. [PubMed Central ID: PMC4100192]. <https://doi.org/10.3390/ijms150610974>.
46. Lee CD, Folsom AR, Blair SN. Physical activity and stroke risk: a meta-analysis. *Stroke*. 2003;**34**(10):2475-81. [PubMed ID: 14500932]. <https://doi.org/10.1161/01.STR.0000091843.02517.9D>.
47. Borenstein M, Higgins JP. Meta-analysis and subgroups. *Prev Sci*. 2013;**14**(2):134-43. [PubMed ID: 23479191]. <https://doi.org/10.1007/s1121-013-0377-7>.