



# Investigation of the Antimicrobial Properties of Ethanolic Extracts of *Momordica charantia*, *Nerium oleander*, *Solanum nigrum*, and *Hyssopus officinalis* on Food Pathogens

Maryam Beigomi<sup>1,\*</sup>, Mohrram Valizadeh<sup>2</sup>

<sup>1</sup>Department of Food Science and Technology, Zahedan University of Medical Sciences, Zahedan, Iran

<sup>2</sup>Department of Physical Geography, University of Sistan and Baluchestan, Zahedan, Iran

\*Corresponding Author: Department of Food Science and Technology, Zahedan University of Medical Sciences, Zahedan, Iran. Email: beigomimaryam@gmail.com

Received: 16 February, 2025; Revised: 23 February, 2025; Accepted: 3 March, 2025

## Abstract

**Background:** Diseases caused by the consumption of food contaminated with pathogenic bacteria are significant public health issues. Given the substantial losses and health risks caused by pathogenic bacteria in humans, the use of medicinal plants as antimicrobial agents is suggested as a better method for controlling food safety.

**Methods:** The studied plants were collected from the Sistan region. Extraction was performed using a rotary device, and the minimum inhibitory concentration (MIC) and minimum lethal concentration (MLC) were determined using the microdilution method.

**Results:** The lowest inhibitory concentration of the ethanolic extract of *Momordica charantia* was 12.5 mg/mL, which inhibited *Staphylococcus aureus* and *Bacillus cereus* strains at this concentration. The highest inhibitory concentration was 50 mg/mL, which inhibited *Vibrio* bacteria at this concentration. The results of the antimicrobial properties of *Hyssopus officinalis* showed that the highest inhibitory concentration was 100 mg/mL, which inhibited *Listeria* and *Shigella* bacteria at this concentration, while the highest lethal concentration was 200 mg/mL.

**Conclusions:** The results of the study demonstrated that *M. charantia* extract exhibited good antimicrobial properties against foodborne pathogens.

**Keywords:** Medicinal Plants, Food Pathogens, Antimicrobial, Ethanol

## 1. Background

The demand for new antimicrobial agents is increasing due to the rise in multidrug-resistant bacteria. Consequently, researchers are exploring new therapeutic alternatives against these resistant strains (1). Plant derivatives have emerged as a potential new source, with these compounds acting in various ways to inactivate or block the growth of pathogens (2).

*Momordica charantia*, from the Cucurbitaceae family, features bright yellow flowers, and its fruit turns yellow or orange when ripe. It is rich in vitamin C (3). This species is notable for its antimicrobial (4), nutritional, and anti-inflammatory properties, and is used in developing countries for its healing properties for stomach ulcers (5), rheumatism, and more. *M. charantia* is employed in folk medicine to treat toothache,

diarrhea, boils, cancer, high blood pressure, obesity, bacterial and viral infections, diabetes, and pneumonia (6).

*Nerium oleander* is an evergreen shrub belonging to the Apocynaceae family, widely distributed worldwide. It possesses antibacterial (7), antifungal (8), antidiabetic (9), antioxidant (10), and antitumor properties (10, 11).

*Hyssopus officinalis* is a member of the Lamiaceae family and is widely grown in Iran. The active ingredients of this plant help control blood pressure, aid digestion, and reduce stomach bloating. The most important compounds in its essential oil include pinocamphe, alpha- and beta-pinene, camphene, and sesquiterpene alcohols, tannins, bitter substances, diosmin, hyssop, and mucilage compounds. Its essential oil is bitter, pungent, dry, and slightly warming (12). This plant is used for various purposes, such as anti-

inflammatory, antibacterial, antipyretic, antispasmodic, antihypertensive, and antilipidemic applications (13). *Hyssopus officinalis* has been shown to inhibit the growth of bacteria (14).

## 2. Objectives

The aim of this study was to investigate the antimicrobial properties of ethanolic extracts of *M. charantia*, *N. oleander*, *Solanum nigrum*, and *H. officinalis* on food pathogens.

## 3. Methods

The leaves of *M. charantia*, *N. oleander*, *S. nigrum*, and *H. officinalis* plants were collected from the plains of Sistan and Baluchestan province, Zabol city. The plant samples were dried in the shade and ground. Five grams of the ground powder was dissolved in 50 cc of ethanol and placed on a shaker for 24 hours. The extract was then filtered and concentrated using a rotary evaporator. The concentrated extracts were stored in the rotary evaporator until use.

### 3.1. Bacterial Strains and Culture Condition

The bacterial strains *Staphylococcus aureus* ATCC1189, *Shigella dysenteriae* ATCC1188, *Listeria monocytogenes* ATCC1298, *Vibrio cholerae* ATCC1611, and *Bacillus cereus* ATCC1015 were obtained from the Microbiology Laboratory of Zabol National University. The bacteria were cultured in a nutrient broth medium, and the microdilution method was used to examine the effect of the extracts on the bacteria. In the wells of a 96-well microplate, 10 microliters of nutrient broth medium were added. In the next step, 100 microliters of the desired plant extract, at different concentrations, were added, followed by 10 microliters of bacteria prepared at half McFarland dilution. The microplates were incubated for 24 hours, and the first clear well was considered as the minimum inhibitory concentration (MIC).

## 4. Results

The lowest inhibitory concentration of the ethanolic extract of *M. charantia* was 12.5 mg/mL, which inhibited *S. aureus* and *B. cereus* strains at this concentration. The highest inhibitory concentration was 50 mg/mL, which inhibited *Vibrio* bacteria at this concentration. The results of the minimum inhibitory concentration of *N. oleander* showed that the lowest inhibitory concentration against *B. cereus* bacteria was 25 mg/mL (Table 1). The antimicrobial properties of *H. officinalis* revealed that the highest inhibitory concentration was

100 mg/mL, inhibiting *Listeria* and *Shigella* bacteria at this concentration, while the highest lethal concentration was 200 mg/mL (Table 1).

## 5. Discussion

Although the antibacterial properties of effective plant compounds, including essential oils and extracts, have been studied in the past, their mechanism of action in reducing or eliminating microbial load requires further investigation. While many chemical compounds in plants are similar, they do not have a specific mechanism for their effect on microorganisms; rather, each compound targets a specific site in the cell. The main factor in the antibacterial effect of plant extracts and essential oils is the chemical compounds that constitute them.

Sabzali investigated the antibacterial properties of the hydroalcoholic extract of *N. oleander* on pathogenic bacteria. The results showed that the most effective hydroalcoholic extract of oleander was at a concentration of 76 mg/mL. The largest diameter of the growth inhibition zone at this concentration was related to *Enterococcus faecalis* bacteria, and the smallest diameter was related to *Pseudomonas aeruginosa*. The results indicated that the lowest MIC was for *S. aureus* at a concentration of 5 mg/mL, and the highest MIC was for *Escherichia coli* and *P. aeruginosa* at a concentration of 76 mg/mL (15).

In the Hamoonnavard study, the antimicrobial effect of *N. oleander* on *S. aureus* and *Staphylococcus epidermidis* showed sensitivity to 40 and 80 µL of 25 mg/mL concentration, all concentrations of 50 mg/mL leaf extract, and all concentrations of 50 mg/mL flower extract (16). Additionally, the antibacterial properties of three ethanolic and aqueous petroleum extracts of *Nerium oleander* on four bacteria (*B. subtilis*, *S. aureus*, *M. luteus*, and *P. aeruginosa*) showed that the largest diameter of the growth inhibition zone was observed at a concentration of 100 mg/mL. The ethanolic extract had a greater effect on *S. aureus* and *M. luteus* with growth inhibition zone diameters of 18 and 14 mm, respectively. *Pseudomonas aeruginosa* and *Bacillus subtilis* were more sensitive to the aqueous extract, with inhibition zone diameters of 15 and 17 mm, respectively, at a concentration of 100 mg/mL (17).

In a study by Mouhcine, which investigated the antimicrobial activity of aqueous and ethanolic extracts of *N. oleander*, the results showed that the diameter of the inhibitory zone of the aqueous extract against *Enterococcus faecalis* was  $10.0 \pm 1.2$  mm and against *L. monocytogenes* was  $4.0 \pm 1.0$  mm, while the diameter of the inhibitory zone of the ethanolic extract against

**Table 1.** Minimum Inhibitory Concentration and Minimum Lethal Concentration of Medicinal Plant Extracts Against Foodborne Pathogens

Variables	MIC/MBC			
	<i>Momordica charantia</i>	<i>Nerium oleander</i>	<i>Solanum nigrum</i>	<i>Hyssopus officinalis</i>
<i>Staphylococcus aureus</i>	12.5 - 25	50 - 100	25 - 50	25 - 50
<i>Listeria</i>	25 - 50	100 - 200	12.5 - 25	100 - 200
<i>Vibrio</i>	50 - 100	50 - 100	50 - 100	50 - 100
<i>Bacillus cereus</i>	12.5 - 25	25 - 50	12.5 - 25	25 - 50
<i>Shigella</i>	25 - 50	50 - 100	50 - 100	100 - 200

Abbreviations: MIC, minimum inhibitory concentration; MBC, minimum bactericidal concentration.

*Enterococcus faecalis* was  $5.3 \pm 0.6$  mm (18). Another study showed that a concentration of 200 µg/mL of *N. oleander* essential oil inhibited the formation of *P. aeruginosa* biofilm (19).

Rajendra et al. found that the benzene extract had a higher inhibitory diameter (14 mm) than the ethanolic extract of *N. oleander* (11 mm) against *B. subtilis* (20). Saedian et al. reported that the minimum inhibitory concentration and minimum bactericidal concentration of *E. coli* showed the highest sensitivity to the alcoholic extract of the leaves, with averages of 62.5 mg/mL and 125 mg/mL, respectively (21). Valizadeh et al. studied the anti-biofilm effect of ethanolic and acetone extracts of *Karla* extract, showing that the lowest inhibitory and lethal concentrations were 1.25 mg/mL and 2.5 mg/mL, respectively (22). Masithoh et al. found that the diameter of the inhibitory zone of *Karla* extract against *Aeromonas hydrophila* was 12.3 mm (23).

In a study of the effect of methanolic extract of *S. nigrum* leaves conducted by Zhao et al. on *E. coli*, *S. aureus*, *B. subtilis*, and *Pasteurella multocida*, they showed that this extract has a relatively moderate effect on these microbes (24).

### 5.1. Conclusions

The results of this study showed that the medicinal plant *Karla* is a better inhibitor than other plants for eliminating foodborne pathogens.

### Acknowledgements

The authors of this article would like to thank all the professors who helped collect and write this article.

### Footnotes

**Authors' Contribution:** Study concept and design: M. B. and M. V.; All authors read and approved the final

manuscript.

**Conflict of Interests Statement:** The authors declare no conflict of interests.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** This study has been approved by Ethic Committee of Zahedan University of Medical Sciences (ethical code number: IR.ZAUMS.REC.1402.378).

**Funding/Support:** The authors declared no financial support to write this manuscript.

### References

- Ali Mirza S, Afzaal M, Begum S, Arooj T, Almas M, Ahmed S, et al. Uptake mechanism of antibiotics in plants. *Antibiotics and Antimicrobial Resistance Genes in the Environment*. Vol. 1. Amsterdam, The Netherlands: Elsevier; 2020. p. 183-8. <https://doi.org/10.1016/b978-0-12-818882-8.00011-5>.
- Michelin DC, Moersch PE, Lima AC, Nascimento GGF, Paganelli MO, Chaud MV. [Evaluation of the antimicrobial activity of vegetal extracts]. *Rev Bras Farmacogn*. 2005;15(4). Portuguese. <https://doi.org/10.1590/s0102-695x2005000400010>.
- Huang HJ, Chen SL, Chang YT, Chyuan JH, Hsieh-Li HM. Administration of *Momordica charantia* Enhances the Neuroprotection and Reduces the Side Effects of LiCl in the Treatment of Alzheimer's Disease. *Nutrients*. 2018;10(12). [PubMed ID: 30513908]. [PubMed Central ID: PMC631675]. <https://doi.org/10.3390/nu10121888>.
- Guarniz WAS, Canuto KM, Ribeiro PRV, Dodou HV, Magalhaes KN, Miranda Sa K, et al. *Momordica charantia* L. Variety from Northeastern Brazil: Analysis of Antimicrobial Activity and Phytochemical Components. *Pharmacog J*. 2019;11(6):1312-24. <https://doi.org/10.5530/pj.2019.11.203>.
- Gurdal B, Kultur S. An ethnobotanical study of medicinal plants in Marmaris (Mugla, Turkey). *J Ethnopharmacol*. 2013;146(1):113-26. [PubMed ID: 23261486]. <https://doi.org/10.1016/j.jep.2012.12.012>.
- Shibib BA, Khan LA, Rahman R. Hypoglycaemic activity of *Coccinia indica* and *Momordica charantia* in diabetic rats: depression of the hepatic gluconeogenic enzymes glucose-6-phosphatase and fructose-1,6-bisphosphatase and elevation of both liver and red-cell shunt enzyme glucose-6-phosphate dehydrogenase. *Biochem J*. 1993;292(1):267-70. <https://doi.org/10.1042/bj2920267>.

7. El Sawi NM, Geweely NS, Qusti S, Mohamed M, Kamel A. Cytotoxicity and Antimicrobial Activity of Nerium oleander Extracts. *J Applied Animal Res.* 2010;**37**(1):25-31. <https://doi.org/10.1080/09712119.2010.9707089>.
8. Hadizadeh I, Peivastegan B, Kolahi M. Antifungal activity of nettle (*Urtica dioica* L.), colocynth (*Citrullus colocynthis* L. Schrad), oleander (*Nerium oleander* L.) and konar (*Ziziphus spina-christi* L.) extracts on plants pathogenic fungi. *Pak J Biol Sci.* 2009;**12**(1):58-63. [PubMed ID: 19579919]. <https://doi.org/10.3923/pjbs.2009.58.63>.
9. Dey P, Saha MR, Chowdhuri SR, Sen A, Sarkar MP, Haldar B, et al. Assessment of anti-diabetic activity of an ethnopharmacological plant Nerium oleander through alloxan induced diabetes in mice. *J Ethnopharmacol.* 2015;**161**:128-37. [PubMed ID: 25498854]. <https://doi.org/10.1016/j.jep.2014.12.012>.
10. Singhal KG, Gupta GD. Hepatoprotective and antioxidant activity of methanolic extract of flowers of Nerium oleander against CCl4-induced liver injury in rats. *Asian Pac J Trop Med.* 2012;**5**(9):677-85. [PubMed ID: 22805717]. [https://doi.org/10.1016/S1995-7645\(12\)60106-0](https://doi.org/10.1016/S1995-7645(12)60106-0).
11. Rashan LJ, Franke K, Khine MM, Kelter G, Fiebig HH, Neumann J, et al. Characterization of the anticancer properties of monoglycosidic cardenolides isolated from Nerium oleander and Streptocaulon tomentosum. *J Ethnopharmacol.* 2011;**134**(3):781-8. [PubMed ID: 21291990]. <https://doi.org/10.1016/j.jep.2011.01.038>.
12. Ghfir B, Fonvieille JL, Koulali Y, Ecalle R, Dargent R. Effect of essential oil of Hyssopus officinalis on the lipid composition of Aspergillus fumigatus. *Mycopathologia.* 1994;**126**(3):163-7. [PubMed ID: 7935731]. <https://doi.org/10.1007/BF01103770>.
13. Ardestani A, Yazdanparast R. Inhibitory effects of ethyl acetate extract of Teucrium polium on in vitro protein glycosidation. *Food Chem Toxicol.* 2007;**45**(12):2402-11. [PubMed ID: 17673348]. <https://doi.org/10.1016/j.fct.2007.06.020>.
14. Nedorostova L, Kloucek P, Kokoska L, Stolicova M, Pulkrabek J. Antimicrobial properties of selected essential oils in vapour phase against foodborne bacteria. *Food Control.* 2009;**20**(2):157-60. <https://doi.org/10.1016/j.foodcont.2008.03.007>.
15. Sabzali S, Bakhtiyari S, Rostamzad A, Haghani K. [A study of the anti-bacterial activities of Nerium oleander's hydroalcoholic extract]. *Sci Magazine Yafte.* 2013;**15**(2):53-9. FA.
16. Asadi-Samani M. Evaluation of Nerium oleander aqueous extract effect on Staphylococcus aureus and Staphylococcus epidermis. *J Shahrekord Univ Med Sci.* 2013;**15**:46-56.
17. Tannu G, Gupta AK, Kumar S, Singh K. anti-microbial activity of Nerium oleander stem extract. *Int J Pharma Professional's Res.* 2011;**2**(1):210-1.
18. Mohammed EM, Mouhcine M, Amin L, Saaid A, Khalil H, Laila B. Cytotoxic, antioxidant and antimicrobial activities of Nerium oleander collected in Morocco. *Asian Pacific J Tropical Med.* 2019;**12**(1). <https://doi.org/10.4103/1995-7645.250342>.
19. Almana TN, Alharbi NS, Ramachandran G, Kanisha Chelliah C, Rajivgandhi G, Manoharan N, et al. Anti-biofilm effect of Nerium oleander essential oils against biofilm forming Pseudomonas aeruginosa on urinary tract infections. *J King Saud Univ-Sci.* 2021;**33**(2). <https://doi.org/10.1016/j.jksus.2021.101340>.
20. Rajendra DC, Asma PB, Jayprakash SS, Meena JK, Pournima SS. Phytochemical screening and antibacterial activity of Nerium indicum leaves. *Int J Pharm Teach Practi.* 2013;**4**(3):1-4.
21. Saeidian S, Eslami M, Dashipoor A. Antibacterial Effect of Alcoholic and Aqueous Extract of Carla on Escherichia coli and Staphylococcus aureus. *Pajouhan Sci J.* 2019;**17**(4):15-24. <https://doi.org/10.52547/psj.17.4.15>.
22. Valizadeh M, ur Rehman F, Hassanzadeh MA, Beigomi M, Fazeli-Nasab B. Investigating the Antimicrobial Effects of Glycyrrhiza glabra and Salvia officinalis Ethanolic Extract Against Helicobacter pylori. *Int J Infect.* 2021;**8**(4). <https://doi.org/10.5812/iji.114563>.
23. Masithoh DA, Kusdarwati R, Handijatno D. Antibacterial activity of bitter gourd (Momordica charantia L.) leaf extract against Aeromonas hydrophila. *IOP Conference Series: Earth Environmental Sci.* 2019;**236**. <https://doi.org/10.1088/1755-1315/236/1/012096>.
24. Zhao WD, Liu DX, Wei JY, Miao ZW, Zhang K, Su ZK, et al. Caspr1 is a host receptor for meningitis-causing Escherichia coli. *Nat Commun.* 2018;**9**(1):2296. [PubMed ID: 29895952]. [PubMed Central ID: PMC5997682]. <https://doi.org/10.1038/s41467-018-04637-3>.