

## Effect of *Pistacia Atlantica* Extract on Glutathione Peroxidase Tissue Levels and Total Oxidative Capacity of Liver and Plasma Lipid Profile of Rats

Parvin Farzanegi,\*<sup>1</sup> Monireh Mousavi,<sup>1</sup> Abbas Ghanbari-Niaki<sup>2</sup>

1. Department of Exercise Physiology, Faculty of Humanities, Islamic Azad University Sari Branch, Sari, Iran
2. Department of Exercise Physiology, Faculty of Physical Education and Sports Sciences, University of Mazandaran, Babolsar, Iran

Article information	Abstract
<p>Article history: Received: 25 Jan 2012 Accepted: 20 May 2012 Available online: 10 Feb 2013 ZJRMS 2013; 15(11): 59-63</p> <p>Keywords: Glutathione peroxidase Total oxidative capacity Lipid profile <i>Pistacia atlantica</i> Endurance exercise</p> <p>*Corresponding author at: Department of Hematology, School of Medical Sciences, Tarbiat Modares University, Tehran, Iran. E-mail: parvin.farzanegi@gmail.com</p>	<p><b>Background:</b> Exercise causes increased oxygen consumption, leaving cells exposed to oxidative stress. Antioxidants may have a protective effect by inhibiting lipid peroxidation. Thus, this study aims to examine the effect of <i>Pistacia atlantica</i> extract on glutathione peroxidase levels and total oxidative capacity of liver and plasma lipid profile of rats.</p> <p><b>Materials and Methods:</b> In this experimental study, 28 female rats' weight <math>155.8 \pm 2.7</math> grams were randomly and equally divided into 4 groups of exercise-saline, control-saline, exercise-mastic, and control-mastic. The exercise groups exercised for 8 weeks (5 days per week, 60 minutes daily, 25 meters per minute, on a zero degree slope). The rats received equal volumes of mastic and saline orally for 4 weeks. Blood and tissue samples were taken 72 hours after the last exercise session. Data were analyzed using one-way variance analysis (ANOVA).</p> <p><b>Results:</b> Consumption of <i>Pistacia atlantica</i> extract together with endurance exercising for 8 weeks did not significantly affect glutathione peroxidase concentration, total oxidative capacity, LDL, triglyceride, or cholesterol, but significantly reduced HDL (<math>p=0.002</math>).</p> <p><b>Conclusion:</b> Results showed that antioxidant and lipid profile levels were not affected by consumption of supplements and endurance exercising. However, further studies are required to assess the long term effects of this herbal extract.</p> <p>Copyright © 2013 Zahedan University of Medical Sciences. All rights reserved.</p>

### Introduction

Free radicals are atoms or molecules with a single electron that makes them highly reactive. Formation of free radicals in living organisms damages macro-molecules such as DNA, proteins, and lipids. To defend against free radicals, body has an antioxidant defense mechanism that includes enzymatic antioxidants like super-oxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX), and non-enzymatic antioxidants such as vitamins A, C, and E, flavonoids, uric acid, bilirubin, ferritin, and thiols like glutathione (GSH), ubiquinone (coenzyme Q<sub>10</sub>), and micronutrients such as, iron, copper, zinc, selenium, manganese [1]. The imbalance between the production of free radicals and antioxidant defense system causes oxidative stress that is involved in pathogenesis of various diseases [2]. Oxygen consumption and metabolism of the whole body is elevated by exercise, and consequently, cells are exposed to oxidative stress and reactive oxygen species [3, 4].

Da Cunha et al. have reported that by increasing anti-oxidative activity of enzymes such as superoxide dismutase, catalase, and glutathione peroxidase in rat's cerebral cortex, exercise can have a protective role against oxidative imbalance and brain damage induced by homocysteine [5]. It is hypothesized that long-term aerobic exercise can significantly improve liver function and lipid profile [6, 7]. To date, many investigations have been conducted, in search of natural antioxidants from

plant sources. Bing et al. point out that consumption of the antioxidant *Kingobiloba* extract increases endurance capacity, delays fatigue, improves athletic ability, and promotes the recovery process after strenuous exercise in rats. This improvement was the result of increased superoxide dismutase activity and lowering of liver malondialdehyde (MDA), causing reduced liver tissue lipid peroxidation damage [8]. Akil et al. declared selenium consumption increases brain tissue GSH concentration in rats following swimming exercise [9]. *Pistacia atlantica* or Mastic is a pistachio species of the Anacardiaceae family [10]. Different concentrations of phenolic compounds in the mastic skin are perfectly able to slow down the oxidation process [11].

In a study, consumption of edible mastic oil and powder increased HDL and decreased blood serum LDL in male and female rabbits [12]. It has also been suggested that dried mixed-nuts may have desirable effects beyond lipid reduction. Sari et al. in their prescription for the participants in a Mediterranean diet substituted saturated fat with pistachio (comprising about 20% of daily caloric intake) for a period of 4 weeks. The pistachio diet led to lower glucose, lipids, interleukin 6, total oxidant status, and malon dialdehyde, and increased superoxide dismutase [13]. As mentioned, most previous studies focused on consumption of antioxidant supplements and exercise separately, without producing comparable results.

Therefore, as no research has been carried out so far into the simultaneous effects of *Pistacia atlantica* extract (mastic) consumption and exercise on some liver tissue antioxidant indices and plasma lipid profile, this study intends to examine the effect of endurance exercise on glutathione peroxidase tissue level and total oxidative capacity of liver and plasma lipid profile of rats, with and without Pistacia plant extract.

## Materials and Methods

For the purposes of this experimental study, 28, 6-8 week-old female rats weight 150-200 grams were procured from the laboratory research and animal reproduction center at Amol Pasteur Institute and kept in polyethylene cages at 22-24°C temperature conditions and 12:12 hour light-dark cycle with no food or water restrictions. They were then randomly divided into 4 equal groups of exercise-saline, control-saline, exercise-mastic, and control-mastic.

The mastic preparation procedure was as follows [14]: Ten grams of mastic fruit powder in 150 ml water was simmered for 45 minutes, and then cooled down to room temperature, and twice filtered through Wattman Number-4 paper. For every 200 grams weight of rats, 1.5 ml (7.5 µl for every gram weight) of the filtered solution was fed to the rats in control-mastic and exercise-mastic groups, 5 days per week, for 4 weeks after the end of the second week, 3 hours after exercise. Simultaneously, equal volume of saline was fed to the rats in control-saline and exercise-saline groups in the same manner.

The exercise program for the two exercise groups included 8 weeks of endurance running exercises (5 days per week, at specified intensity and duration) on a specially designed treadmill. After a 5-day induction, the exercise program began as follows: First week: first session; rats exercised on the treadmill at speed of 20 m/min for 20 minutes. By the fifth session, speed was increased to 25 m/min and duration to 35 minutes. Second week: speed was increased to 25 m/min and duration to 60 minutes for each session. These conditions were maintained through to the end of the eighth week.

On each day after completion of the exercise course, 2 rats were randomly selected from each of the exercise and control groups. Seventy-two hours after the last exercise session, and after 4 hours of fasting, these rats were anesthetized by intra-peritoneal injection of ketamine (30-50 mg/kg) and xylazine (3-5 mg/kg). Then, immediately, blood samples were taken from the right ventricle with an EDTA-coated syringe, and transferred to EDTA-coated test tubes.

Blood samples were then centrifuged at 1500 rpm for 10 minutes. Plasma and liver tissue were frozen in liquid nitrogen and kept at -80°C for subsequent measurements. To avoid the nightly rhythm effect, sampling began at 8 o'clock in the morning and finished by 11.30 am. To examine glutathione peroxidase and total oxidative capacity of liver, first, the liver tissue was homogenized in a phosphate buffer (17 mMol and 7.4 pH) at the speed

of 3000×g and temperature of 4°C for 15 minutes using a mechanical homogenizer (Polytron, Germany).

The top liquid from homogenized centrifuged samples was used for tests. Glutathione peroxidase level and total oxidative capacity were analyzed using ELISA method in accordance with the guidelines of German manufactured kit, Cayman (Sensitivity: 0.09, Intra-Assay CV%: 4.5), and values of 8.7 mg/ml and 11.4 U/ml were obtained for Glutathione peroxidase and total oxidative capacity, respectively. These results were read using ELISA reader kit (Sunrise model, Takken Corporation, Vienna, Austria).

Cholesterol was measured using enzymatic colorimeter method in accordance with kit manufacturer's guidelines (Greiner Bahlingen Company, Germany, with 4 mg/dl sensitivity). HDL was measured using the same method and with Randox Company kit (sensitivity: 3 mg/dl) of Antrim, U.K. Triglyceride was measured by the German, Greiner Bahlingen kit with 1.5 mg/dl sensitivity. LDL was measured using the values of HDL-c, TG, and TC. Tests results were analyzed by ELISA- reader kit (Ststfax, U.S.).

Normal distribution of data was assured by Kolmogorov-Smirnov test, and significant differences between groups were determined using one-way variance analysis (ANOVA). Tukey test was used for assessment of changes between groups. Data were processed by SPSS-16 software, with significance level of  $p \leq 0.05$ .

## Results

After 8 weeks of endurance exercise and consumption of mastic, the following results were obtained:

Glutathione peroxidase concentration reduced in the exercise-mastic group compared with the three other groups. But this reduction was not significant. The difference between exercise-mastic and exercise-saline groups in terms of changes in glutathione peroxidase was not significant, either. In the exercise-mastic group, compared with the three other groups, only an insignificant reduction in total oxidative capacity concentration was observed. Also, the changes in total oxidative capacity in exercise-mastic and exercise-saline groups were insignificant.

HDL concentration in exercise-mastic group, compared with other groups reduced significantly ( $p=0.002$ ), but changes in HDL in exercise-mastic and exercise-saline groups was not significant.

LDL concentration increased insignificantly in exercise-mastic group, compared with three other groups. In addition, changes in LDL between exercise-mastic and exercise-saline groups were insignificant.

Triglyceride concentration in exercise-mastic group showed an insignificant reduction compared with three other groups. Changes in triglyceride in exercise-mastic and exercise-saline groups were insignificant. Cholesterol concentration in exercise-mastic group reduced insignificantly compared with three other groups. Changes in cholesterol in exercise-mastic and exercise-saline groups were not significant, either (Table 1).

**Table 1.** Descriptive indices of study main variables (mean±SD)

Variable	Control-saline group	Exercise-saline group	Control-mastic group	Exercise mastic group	p-Value
Glutathione peroxidase (nmol/ml)	0.10±0.001	0.11±0.0009	0.105±0.0007	0.105±0.002	0.898
Total oxidative capacity (U/ml)	0.037±0.006	0.04±0.006	0.036±0.003	0.035±0.003	0.242
HDL(mg/dl)	49.13±4.81	53.35±6.85	47.086±2.87	45.12±1.351	0.002*
LDL(mg/dl)	40.065±19.75	28.84±15.52	26.29±23.29	44.61±20.58	0.259
Triglyceride (mg/dl)	92.14±17.72	91.71±23.16	97.71±16.97	105.29±14.90	0.493
Cholesterol (mg/dl)	108.29±20.33	100.43±16.98	93.00±23.88	110.71±22.33	0.4

\* $p \leq 0.05$  significance level

## Discussion

Results of the study showed, after 8 weeks consumption of *Pistacia atlantica* extract in combination with endurance exercises, the desired effects were not observed in glutathione peroxidase concentration, liver total oxidative capacity, and LDL, triglyceride, or cholesterol concentrations. However, there was a significant reduction in plasma HDL.

Evidence shows a change in enzymatic antioxidants due to exercise leads to oxidative stress [5]. Studies in the past have provided evidence showing that adaptation of antioxidant activity with exercise is induced by a change in mRNA gene expression and regulation of protein enzyme levels. This adaptation is effectively influenced by physiological and environmental factors such as gender, age, and diet [16, 17]. It is hypothesized that consumption of mastic as an antioxidant, increases activity of anti-oxidative enzymes and reduces cholesterol levels in exercised people, and consequently, exercise-induced oxidative stress is effectively reduced [11, 12]. However, in this study, glutathione peroxidase level was lower in exercise-mastic group compared with other groups, but statistically, this was insignificant. Also, between exercise-mastic and exercise-saline groups, the difference in glutathione peroxidase changes was not significant. This meant that, consumption of mastic was not able to increase glutathione peroxidase activity, and could not help improve antioxidant defense system. Glutathione peroxidase reduction have also been shown in other studies [18, 19]. There is evidence that shows intense physical activity disrupts glutathione peroxidase balance through reducing its level in tissues, changing redox status of cells (cell regeneration), and interfering in its production and transfer [20]. Gul et al. described reduced glutathione peroxidase activity level in heart tissue following endurance exercise as exhaustive [21]. Unlike results of this study, consumption of ginkgo boliba [8] and selenium [9] caused an increase in glutathione peroxidase. Along the same lines, Leelarungrayub et al. study showed reduced malondialdehyde and nitrite oxide, and increased Q<sub>10</sub> and glutathione peroxidase following 9 days of exercise and 300 mg of Q<sub>10</sub> intake, 12 days after that, in young swimmers [22]. But, Tiidus et al. reported no significant change in glutathione peroxidase in men and women following 8 weeks of cycling exercise [19]. It has been reported that animals lacking antioxidants such as vitamin E and selenium in their food, have higher antioxidant and exercise compatibility compared with

those having normal diet [23, 24]. Also, it has been suggested that antioxidant supplements should not be taken prior to exercise, as they impede adaption of muscle cells. In other words, consumption of antioxidants just before competition, when probably feeling exhausted, leads to production of Reactive Oxygen Species (ROS), and may be ineffective [25]. In this study, subjects were animals that freely received supplements during the 8 weeks of experiment.

Increased oxygen intake and tissue damage during exercise, may stimulate the activity of inflammatory cells like neutrophils, which leads to ROS production by NADPH oxidase (Nicotinamide Adenine Dinucleotide Phosphate-oxidase). Also, increasing concentration of catecholamines and temperature during exercise causes oxidative stress. Hence, increasing ROS requires more antioxidants activity [4].

In this study, liver total oxidative capacity reduced in exercise-mastic group, with no significant difference between groups. Also, liver total oxidative capacity increased in exercise group. This increase was also observed in other similar studies [25]. In Aguilo et al. study, increased levels of thiobarbituric acid (total oxidative damage indicator) was observed in aerobic exercise group after a tiring practice session [26]. A possible reason for this result could be short exercise period or non-consumption of supplements during exercise. The general public is of the opinion that regular physical exercise and supplement intake leads to adaptations that elevate total oxidative capacity for defending against oxidative stress. Sacke et al. in their study observed a significant reduction in oxidative stress indicators after exercise in experimental groups, and a significant increase in total oxidative capacity compared with control group [27]. The reason for contradicting results of this study and Sacke's may be non-consumption of plant extract.

Other results of this study revealed significant reduction in HDL concentration after 8 weeks of exercise and mastic intake in exercise-mastic group compared with others. However, no significant change was seen in LDL concentration, triglyceride, or cholesterol. This may be the very first study that showed an unexpected reduction in HDL level in exercise-mastic group. Along these lines, Young Choi et al. in an investigation into the effect of coffee on lipid profile and antioxidants in fit rats observed a reduction in HDL level and an increase in total oxidative capacity [28]. In another study by George et al. on the effects of progressive resistance exercise on lipids

and lipoproteins in adults, a reduction in HDL and cholesterol was reported [29].

In confirmation of previous studies [12, 13], and the desirable effects of pistachio on lipid profile, Edwards et al. [30] and Kocyigi et al. [31] reported that 3 weeks of pistachio diet significantly reduced cholesterol, TC/HDL, and LDL/HDL and significantly increased HDL in patients and volunteers with normal fat. Sheridan et al. reported, 4 weeks of consumption of pistachio caused a significant reduction in TC/HDL and LDL/HDL, and significantly increased HDL in patients with moderate hypercholesterolemia. But, cholesterol, LDL, and triglyceride levels did not change [32]. In Edward's study, pistachio replaced recommended daily fat intake (20% daily calorie). Therefore, positive effects of pistachio on lipid profile may have been due to reduced daily calorie from fat intake [30]. Previously, pistachio was regarded as a rich diet of MUFA and effective in reducing CHD risk in diabetic patients [33]. High fiber content of pistachio may explain its desirable effects on lipid parameters. Most studies have recommended pistachio consumption of 15-20 percent of daily calorie intake. In a recent study by Aksoy, pistachio consumption level equal to 20% of daily calorie intake in laboratory rats showed positive effects on lipid profile and LDL oxidation. But, when consumption was doubled (equal to 40%), these effects were reversed [35].

Given that pistachio is a fatty nutrient, could increased daily calorie (by 20%) and the subsequent weight gain be detrimental? Results showed that addition of varying amounts of pistachio to the daily calorie did not tangibly increase subject's weight. As rats are genetically similar and had the same diet, therefore, consumption of pistachio as 20% of daily calorie appears rational. Even though, cholesterol reducing mechanisms of pistachio are unknown, but the following may have an impact: 1- improved endothelial function, 2- A compound of

pistachio called Arginine (nitric oxide precursor) may enhance EDV, 3- Pistachio contains fiber, vitamins, micronutrients, and fitsterol, and with a significant reduction in total oxidant status, it may play an important role in improving the pathogenesis and progression of atherosclerosis [33, 34]. Oxidation and antioxidant components may be influenced by different factors, but total oxidant status is more practical to measure than other variables. Therefore, reduction in TOS after a pistachio diet could provide extra data about positive effects of pistachio on lipid reduction.

Generally, this study showed that consumption of *Pistacia atlantica* plant extract together with endurance exercises did not have significant effects on liver tissue antioxidant indicators, or on plasma lipid profile (LDL, triglyceride, and cholesterol) of exercised rats, but caused a reduction in HDL. Therefore, not only the results of other studies about positive effects of pistachio diet with exercise on lipid profile was not confirmed, but also, the idea that pistachio intake with exercise has effects beyond lipid reduction must be more prospectively investigated.

### Acknowledgements

This paper is the result of thesis belonging to Mis. Monireh Mosavi under code No. We sincerely appreciate authorities of Islamic Azad University, Sari Science and Research Branch for helping us conduct this research.

### Authors' Contributions

All authors had equal role in design, work, statistical analysis and manuscript writing.

### Conflict of Interest

The authors declare no conflict of interest.

### Funding/Support

Islamic Azad University Sari Branch.

### References

1. Filaire E, Rouveix M, Massart A, et al. Lipid peroxidation and antioxidant status in rat: Effect of food restriction and wheel running. *Eur J Appl Physiol* 2009; 107(2): 243-50.
2. Rowlands DS, Pearce E, Aboud A. Oxidative stress, inflammation, and muscle soreness in an 894-km relay trail run. *Eur J Appl Physiol* 2011; 112(5): 1839-48.
3. Sun L, Shen W, Liu Z, et al. Endurance exercise causes mitochondrial and oxidative stress in rat liver: Effects of a combination of mitochondrial targeting nutrients. *Life Sci* 2010; 86(1-2): 39-44.
4. Hamdi P. The effects of gender and exercise on malondialdehyde, nitric oxide and total glutathione levels in rat liver. *Afr J Pharm Pharmacol* 2011; 5(4): 515-21.
5. da Cunha MJ, da Cunha AA, Ferreira AG, et al. Physical exercise reverses glutamate uptake and oxidative stress effects of chronic homocysteine administration in the rat. *Int J Dev Neuro Sci* 2012; 30(2): 69-74.
6. Farzanegi P, Mohammadi-Rishsefid N, Habibian M and Jafari H. [The effect of omega-3 on oxidative stress in elite karate athletes] *Persian [Abstract]. J Mazand Univ Med Sci* 2012; 22(91): 70-80.
7. Sastre J, Asensi M, Gasc E, et al. Exhaustive physical exercise causes oxidation of glutathione status in blood, prevention by antioxidant administration. *Am J Physiol* 1992; 263: 992-5.
8. Bing Y, Zhaobao W. Effects of Ginkgo biloba extract on free radical metabolism of liver in mice during endurance exercise. *Afr J Tradit Complement Altern Med* 2010; 7(4): 291-5.
9. Akil M, Bicer M, Menevse E, et al. Selenium supplementation prevents lipid peroxidation caused by arduous exercise in rat brain tissue. *Bratisl Lek Listy* 2011; 112(6): 314-7.
10. Saffar-Zadeh A. Determination of the chemical composition of acorn (*Quercus branti*) *Pistacia atlantica* and *Pistacia Khinjck* seeds as non-conventional feedstuffs. *Acta Agr Kapos* 1999; 3(3): 59-69.
11. Saeb M, Nazifi S, Mirzaei A. [Studies on the effects of Turpentine oil on the serum concentration of lipids and Lipoproteins of female rabbits] *Persian [Abstract]. J Shahid Sadoughi Uni Med Sci Health Serv* 2004; 3(12): 42-50.

12. Nazifi S, Saeb M, Yavari M and Jalae J. [Studies on the effects of turpentine powder on the serum concentration of lipids and lipoproteins of male rabbits] Persian [Abstract]. Iran J Endocrinol Metab 2005; 7(1): 73-8.
13. Sari I, Baltaci Y, Bagci C, et al. Effect of pistachio diet on lipid parameters, endothelial function, inflammation, and oxidative status: A prospective study. Nutrition 2010; 26(4): 399-404.
14. Hamdan II, Afifi FU. Studies on the in vitro and in vivo hypoglycemic activities of some medicinal plants used in treatment of diabetes in Jordanian traditional medicine. J Ethnopharmacol 2004; 93: 117-21.
15. Ghanbari-Niaki A, Abednazari H, Tayebi SM, et al. Treadmill training enhances rat agouti-related protein in plasma and reduces ghrelin levels in plasma and soleus muscle. Metabolism 2009; 58(12): 1747-52.
16. Shao JP. Anti fatigue effect of aqueous extract of salvia in endurance training rats' skeletal muscle. Int J Phy Scis 2011; 6(11): 2697-700.
17. Shaarawy SM, Tohamy AA, Elgendy SM, et al. Protective effects of garlic and silymarin on NDEA-induced rats hepatotoxicity. Int J Biol Sci 2009; 5(6): 549-57.
18. Leeuwenburgh C, Ji LL. Glutathione and Glutathione Ethyl Ester supplementation of Mice Alter Glutathione homeostasis during exercise. J Nutr 1998; 128: 2420-6.
19. Tiidus PM, Pushkarenko J, Houston ME. Lack of antioxidant adaptation to short-term aerobic training in human muscle. Am J Physiol 1996; 271: 832-836.
20. Smarsh DN, Liburt N, Streltsova J, et al. Oxidative stress and antioxidant status in intensely exercising horses administered nutraceutical extracts. Equine Vet J 2010; 42(Suppl 38): 317-22.
21. Gul M, Demirecan B, Taysi S. Effects of endurance training and acute exhaustive exercise on oxidant defense mechanisms in rat heart. Comp Biochem Physiol A Mol Integr Physiol 2006; 143(2): 239-45.
22. Leelarungrayub D, Sawattikanon N, Klaphajone J and Pothongsunan P. Coenzyme Q10 supplementation decreases oxidative stress and improves physical performance in young swimmers: A pilot study. The Open Sports Med J 2010; 4: 1-8.
23. Deminice R, Jordao AA. Creatine supplementation reduces oxidative stress biomarkers after acute exercise in rats. Amino Acids 2012; 43(2): 709-15.
24. Ji LL, Stratman FW, Lardy HA. Antioxidant enzyme systems in rat liver, and skeletal muscle: Influences of selenium deficiency, chronic training and acute exercise. Arch Biochem Biophys 1998; 263: 150-60.
25. Lafay S, Jan C and Nardon K. Grape extract improves antioxidant status and physical performance in elite male athletes. J Sports Sci Med 2009; 8: 468-80.
26. Aguilo A, Tauler P, Guix MP. Effect of exercise intensity and training on antioxidants and cholesterol profile in cyclists. J Nutr Biochem 2003; 14(6): 319-25.
27. Sackeck J, Blumberg JB. Role of vitamin and oxidative stress in exercise. Nutrition 2009; 17(2): 809-14.
28. Young Choi E, Young Jang J, Ok Cho Y. Coffee intake can promote activity of antioxidant enzymes with increasing MDA level and decreasing HDL-cholesterol in physically trained rats. Nutr Res Pract 2010; 4(4): 283-9.
29. Goerge A, Kiristis S. Impact of progressive resistance training on lipids and lipoproteins in adults. Prev Med 2009; 48(1): 9-19.
30. Edwards K, Kwaw I, Matud J and Kurtz I. Effect of pistachio nuts on serum lipid levels in patients with moderate hyper cholesterolemia. J Am Coll Nutr 1999; 18: 229-32.
31. Kocyigit A, Koylu AA, Keles H. Effects of pistachio nuts consumption on plasma lipid profile and oxidative status in healthy volunteers. Nutr Metab Cardiovasc Dis 2006; 16: 202-9.
32. Sheridan MJ, Cooper JN, Erario M and Cheifetz CE. Pistachio nut consumption and serum lipid levels. J Am Coll Nutr 2007; 26: 141-8.
33. Kris-Etherton PM, Pearson TA, Wan Y, et al. High-monounsaturated fatty acid diets lower both plasma cholesterol and triacylglycerol concentrations. Am J Clin Nutr 1999; 70: 1009-15.
34. Anderson JW, Smith BM, Gustafson N. Health benefit and practical aspects of high fiber diets. Am J Clin Nutr 1994; 59(suppl): 1242S-7.
35. Aksoy N, Aksoy M, Bagci C, et al. Pistachio intake increases high density lipoprotein levels and inhibits low-density lipoprotein oxidation in rats. Tohoku J Exp Med 2007; 212: 43-8.

**Please cite this article as:** Farzanegi P, Mousavi M, Ghanbari-Niaki A. Effect of Pistacia atlantica extract on glutathione peroxidase tissue levels and total oxidative capacity of liver and plasma lipid profile of rats. Zahedan J Res Med Sci (ZJRMS) 2013; 15(11): 59-63.