

Association of Maternal Body Mass Index with Adverse Maternal and Prenatal Outcomes

Rahele Alijahan,¹ Babak Nakhostin,² Sousan Salimi,¹ Sadegh Hazrati³*

1. Ardabil District Health Center, Ardabil University of Medical Sciences, Ardabil, Iran
2. General Physician, Ardabil District Health Center, Ardabil University of Medical Sciences, Ardabil, Iran
3. Department of Environmental Health, Faculty of Public Health, Ardabil University of Medical Sciences, Ardabil, Iran

Article information	Abstract
<p>Article history: Received: 12 May 2012 Accepted: 20 Sep 2012 Available online: 3 Mar 2013 ZJRMS 2013;15(9): 56-62</p> <p>Keywords: Maternal body mass index Pregnancy Prenatal outcomes Maternal outcomes</p> <p>*Corresponding author at: Department of Environmental Health, Faculty of Public Health, Ardabil University of Medical Science, Ardabil, Iran E-mail: S.Hazrati@Arums.ac.ir</p>	<p>Background: The present study aimed to determine association between abnormal maternal body mass index and adverse maternal/prenatal outcomes</p> <p>Materials and Methods: In this descriptive-correlation study 8270 pregnant women referred to rural and urban health centers of Ardabil district (from Mar 2009 to Dec 2010) were studied. Data were collected from prenatal healthcare records using a self designed questionnaire. Women with twin pregnancy, less than 18 and above 35 of age, and women with systemic or chronic disease were excluded from the study. The variables examined in this study include, demographic information (e.g. age, social and economy status, and literacy), present pregnancy information (e.g. parity, hemoglobin level, gestational diabetes, preeclampsia) and prenatal information (e.g. preterm delivery, low birth weight, and congenital malformation). Data were analyzed through Kruskal wallis, chi-square, and logistic regression tests using SPSS-16.</p> <p>Results: Eight point two, 25 and 15.4% pregnant of women were underweight, overweight, and obese, respectively. Obese women were at increased risk for macrosomia (OR=1.820, CI: 1.345-2.447, $p=0.001$), unwanted pregnancy (OR= 1.436, CI: 1.198-1.720, $p=0.001$), pregnancy induced hypertension (OR= 1.633, CI: 1.072-2.486, $p=0.022$), preeclampsia (OR= 4.666, CI: 2.353-9.2550, $p=0.001$), and still birth (OR= 2.602, CI: 1.306-5.184, $p=0.007$). However, the risk of low birth weight delivery in underweight women were 1.6 times higher than the normal cases (OR= 1.674, CI: 0.962-2.912, $p=0.068$).</p> <p>Conclusion: Considering high prevalence of abnormal maternal body mass index and its associated adverse maternal and prenatal outcomes; consultation before pregnancy is recommended in order to achieve normal body mass index and reduce the relevant complications.</p>

Copyright © 2013 Zahedan University of Medical Sciences. All rights reserved.

Introduction

Body mass index is a determinant of excess body fat status, which is calculated as body weight in kilograms divided by height in meters squared [1, 2]. WHO and the national institutes of health define underweight as a body mass index ≤ 18.5 kg/m², normal weight as a BMI of 18.5-24.9 kg/m², overweight as a BMI of 25-29.9 kg/m², and obesity as a BMI of ≥ 30 kg/m² [2, 3].

Women's overall health is influenced by body weight [1]; a women's risk of disease rises in proportion to the increase in body weight [4]. During pregnancy and childbirth, greater body weight is associated with an elevated risk of hypertensive disorders, gestational diabetes, cesarean birth, fetal macrosomia, thromboembolic disease, stillbirth, perinatal death, preterm delivery and fetal birth defect [5, 6].

The effect of being underweight in pregnancy on obstetric performance is less clear. While some have found increased incidences of preterm delivery, low birth weight, and increased prenatal loss in these women, others have reported a protective effect of maternal

underweight on certain pregnancy complications and interventions [7]. It is clear that women with abnormal body mass index require more healthcare resources during pregnancy than women of normal body weight and hence increase pressure on our health care system. Therefore, it is of public health importance to study the impact of maternal body mass index on adverse pregnancy and birth outcomes.

There are earlier studies that have investigated the impact of maternal body mass index on selected obstetric outcomes in Islamic Republic of Iran, but most of them were limited to one or two maternal BMI classes or were performed in a small sample size [8-11].

The objective of the present study was to thoroughly assess, in a large retrospective data, adverse obstetric and neonatal outcomes over the maternal BMI strata.

Materials and Methods

In this descriptive correlational study, the population included all the women having pregnancy from March

2009 to December 2010 coming to urban and rural health centers in Ardabil. All in all, 8270 women who ended their pregnancy due to abortion, preterm delivery, still birth and delivery at term were included in the study. The data were collected through examining the recorded personal profiles and filling out a questionnaire.

Those women whose heights and weights were not recorded in their personal profiles and the ones who did not receive pregnancy care were excluded from the study. Individuals aged less than 18 and above 35 and those having diseases such as coronary, pulmonary, thyroid, asthma and multiple pregnancies were also excluded from the study. Data were collected through a researcher-made questionnaire by 10 trained midwives.

The questionnaire included 95 questions, of which 8 were on maternal demographic information (maternal age, socioeconomic status, educational level). 70 questions were about the women's current pregnancy characteristic (high blood pressure, preeclampsia, anemia, the way of delivery, pregnancy diabetes, urinary infection, unwanted pregnancy, hyperemesis gravidarum), and finally 17 questions were about newborn characteristic (birth weight, age of birth, having jaundice, hospitalization). The validity of the questionnaire was verified through content validity approach by three professors of Ardabil University of Medical Sciences.

All the pregnant women coming to Ardabil health centers for prenatal care, generally will receive maternal integrated care with an interval of one to two months based on the guidelines provided by Iranian Ministry of Health. Pregnancy tests of hemoglobin, hematocrit, complete urine analysis, glucose tolerance tests (GCT, GTT) will be taken during 26-30 weeks of pregnancy and the results are recorded in their profiles. Diagnosis of anemia, pregnancy diabetes, preeclampsia, unwanted pregnancy, and high blood pressure were done according to ministerial guidelines of integrated care. Hemoglobin levels of under 11 g/dl during 6-10 weeks of pregnancy and under 10.5 g/dl during 26-30 weeks were considered as anemia. Having urinary infection was taken in the form of bacterial growth more than 100,000 organisms per ml in the urine culture.

The lack of tendency of one or both of the parents to have a baby and unwanted pregnancy were recorded in the health profiles. Systolic blood pressure of 140 and/or higher and diastolic blood pressure of 90 and higher in a mother who had previously normal blood pressure were considered as an increase in the rate of general blood pressure. The same phenomenon plus the positive test of urinary protein was taken as an indicator of preeclampsia. If the mother had previous hospitalization due to serious pregnancy vomiting, it was also accounted for hyperemesis gravidarum. If the result of GCT test was 140 mg/dl or higher, fasting blood sugar test and one-hour, two-hour and three-hour GTT was requested. If the results of two of those four tests were unusual, having diabetes was put forward [12].

In order to calculate maternal BMI, the height and weight of the first three months of pregnancy were used. As in the health system of Iran, classification of BMI

during pregnancy care is performed according to guidelines of Ministry of Health, the classification of BMI in line with the above-mentioned guidelines was done in the form of thin BMI ($<19.8 \text{ kg/m}^2$), normal weight (the BMI from 19.8 to 26 kg/m^2), overweight (the BMI between 26.1 and 29.9) and obese ($\geq 30 \text{ kg/m}^2$) [13]. This classification is a little bit different from the one proposed by WHO.

Due to large sample size of the present study and the recorded data in the profiles, our access to each subject to have a written or oral agreement was not possible. The researchers were committed to keep confidential the personal profiles of the subjects. The data after collection and classification were analyzed through inferential and descriptive statistical tests of chi square, Kruskal-Wallis and logistic multiple regression using SPSS-16. The *p*-value was set at lesser or equal to 0.05.

Results

Based on the classification of BMI, 675 (8.2%) of the subjects were thin, 4252 (51.4%) normal, 2069 (25%) slightly overweight, and 1274 (15.4%) were obese. Different groups of BMI had statistically significant relationship with education, socioeconomic class, age and place of residence. The prevalence of obese BMI was more in women with low education, higher socioeconomic class, the age over 25 and those living in urban areas (Table 1).

Different classifications of BMI had statistically significant relationship with pregnancy diabetes, preeclampsia, high blood pressure, caesarian delivery, abortion, still birth, and unwanted pregnancy. The rate of nulliparity, prim gravidity and anemia in women with thin BMI was higher. In addition, the frequency of multiparity, diabetes, high blood pressure, preeclampsia, caesarian, abortion, still birth and unwanted pregnancy was high in women with the overweight and obese conditions. There was not any statistically significant relationships between maternal BMI and urinary infection, hyperemesis gravidarum, spotting and bleeding, leakage, polyhydramnion, blood type and the maternal Rh (Table 2).

Different categories of BMI did have statistically significant relationships with low birth weight, macrosomia and newborn's gender. The rate of preterm delivery and low birth weight in women with thin BMI was more frequent.

The frequency of macrosomia, congenital malformations and non-successful breastfeeding was high in women with overweight and obese body mass index. High rate of female and male genders were observed in newborns' of women with thin BMI and overweight and obese women, respectively (Table 3). Based on the results of logistic regression, the probability of fetal macrosomia in overweight women was 1.5 times and 2 times in fat ones. The odds of caesarian delivery were 1.7 fold in overweight women and 2 fold in obese ones.

The chance of unwanted pregnancy in fat women was 1.6 times, having high blood pressure in overweight and

obese groups were 2.5-3 times, respectively. The likelihood of fetus being male in women who were slightly overweight and obese was 1.2 times. The risk of preeclampsia was 3 fold in overweight and 4.5 times in obese women, the probability of still birth was 2.6 times

in obese women and the odds of low birth weight was 1.6 fold higher in thin ones. The probability of having anemia decreased with increasing BMI and declined with decreasing BMI regarding fetal macrosomia and caesarian delivery (Table 4).

Table 1. Maternal demographic characteristics for the four BMI categories

General characteristics	Maternal body mass index	Under weight N(%)	Normal N(%)	Overweight N(%)	Obese N(%)	p-Value
Maternal education	Secondary \geq	343(50.9)	2175(51.6)	1051(51.4)	736(58.2)	0.001
	High school \leq	330(49.1)	2040(48.4)	994(48.6)	529(41.8)	
Social class	Middle and low	552(83)	3334(79.9)	1557(76.7)	948(75.9)	0.001
	High	113(17.0)	840(20.1)	473(23.3)	301(24.1)	
Maternal age	18-25	520(77.4)	2514(59.4)	885(43.0)	411(32.4)	0.001
	26-30	121(18.0)	1194(28.2)	704(34.3)	483(38.0)	
	30-35	31(4.6)	525(12.4)	469(22.8)	376(29.6)	
Area of resident	Urban	564(83.7)	3528(83.0)	1820(88.0)	1132(88.9)	0.001
	Rural	110(16.3)	724(17.0)	249(12.0)	142(11.1)	

Table 2. Present pregnancy characteristics based on BMI categories

Body Mass Index maternal variables		Under weight		Normal		Over weight		Obese		p-Value
		N	%	N	%	N	%	N	%	
Gravidity	1	466	70.8	2336	56.1	765	37.9	336	26.9	0.001
	2	144	21.9	1274	30.6	788	39.0	492	39.4	
	3	37	5.6	384	9.2	311	15.4	274	21.9	
	4	9	1.4	108	2.6	109	5.4	102	8.2	
	5 \leq	2	0.3	62	1.5	45	2.2	45	3.6	
Parity	0	111	38.9	611	26.1	225	16.0	108	11.1	0.001
	1	140	49.1	1307	55.8	831	59.2	571	58.7	
Urinary tract infection	2 \leq	34	11.9	425	18.1	347	24.7	293	30.1	0.921
	yes	213	32.7	1313	32.1	653	32.6	387	31.5	
Hemoglobin level in 6-10 week of gestation	no	439	67.3	2778	67.9	1352	67.4	842	68.5	0.052
	11>	32	4.8	216	5.2	75	3.7	52	4.2	
Hemoglobin level in 26-30 week of gestation	11 \leq	630	95.2	3903	94.8	1944	96.3	1176	95.8	0.001
	10.5>	84	17.0	476	15.5	179	13.2	105	12.0	
Gestational diabetes	10.5 \leq	410	83.0	2588	84.5	1289	87.8	771	88.0	0.050
	Yes	6	0.9	35	0.8	27	1.3	21	1.7	
Preeclampsia	No	659	99.1	4134	99.2	1998	98.7	1224	98.3	0.001
	Yes	1	0.2	14	0.3	20	1.0	21	1.7	
Hypertension	No	665	99.8	4181	99.7	2024	99.0	1243	98.3	0.005
	Yes	5	0.7	27	0.6	27	1.3	20	1.6	
Hyperemesis Gravidarum	No	665	99.3	4183	99.4	2018	98.7	1243	98.4	0.585
	Yes	56	8.4	350	8.3	155	7.6	93	7.4	
Cesarean	No	614	91.6	3859	91.7	1896	92.4	1171	92.6	0.001
	Yes	184	38.6	1459	49.1	871	60.7	590	67.4	
Bleeding or spotting	No	293	61.4	1512	50.9	563	39.3	285	32.6	0.496
	Yes	19	2.8	130	3.1	64	3.1	29	2.3	
Abortion	No	648	97.2	4074	96.9	1970	96.9	1227	97.7	0.023
	Yes	4	0.6	54	1.3	31	1.5	28	2.2	
Still birth	No	665	99.4	4158	98.7	2023	98.5	1238	97.8	0.006
	Yes	1	0.3	18	0.4	13	0.6	16	1.3	
Leakage	No	669	99.7	4190	99.6	2039	99.4	1249	98.7	0.081
	Yes	15	2.3	76	1.8	35	1.7	11	0.9	
Oligohydramnios	No	650	97.7	4124	98.2	2007	98.3	1247	99.1	0.979
	Yes	2	0.5	15	0.6	6	0.5	4	0.6	
Unwanted pregnancy	No	396	99.5	2369	99.4	1130	99.5	664	99.4	0.003
	Yes	27	4.0	156	3.7	94	4.6	77	6.1	
Blood group	A	245	36.7	1527	36.9	713	35.2	471	37.8	0.226
	B	141	21.1	765	18.5	399	19.7	222	17.8	
	AB	32	4.8	289	7.0	158	7.8	88	7.1	
	O	249	37.3	1555	37.6	754	37.3	466	37.4	
Rh	Positive	593	90.0	3656	89.2	1781	89.1	1067	86.7	0.060
	Negative	66	10.0	442	10.8	218	10.9	164	13.3	

Table 3. Neonatal characteristics based on BMI categories

Maternal variables	Body mass index	Under weight		Normal		Over weight		Obese		p-Value
		N	%	N	%	N	%	N	%	
Gestational age	Term	381	93.4	2407	95.2	1174	95.8	692	96.2	0.141
	Preterm	27	6.6	122	4.8	52	4.2	27	3.8	
Low birth weight	2500>	21	5.6	83	3.5	25	2.2	20	3.0	0.010
	2500≤	354	94.4	2298	96.5	1122	97.8	644	97.0	
Macrosomia	Yes	20	5.3	204	8.6	151	13.2	101	15.2	0.001
	No	355	94.7	2177	91.4	996	86.8	563	84.8	
Jaundice	Yes	230	56.7	1474	58.4	704	57.9	432	59.5	0.808
	No	176	43.3	1048	41.6	511	42.1	294	40.5	
Hospitalization	Yes	31	7.6	183	7.1	73	6.0	49	6.7	0.521
	No	377	92.4	2378	92.9	1152	94.0	680	93.3	
Cause of hospitalization	Jaundice	10	35.7	79	43.4	33	42.3	23	51.1	0.730
	Prematurity	8	28.6	3	10.7	11	6.0	5	6.4	
	Breathing distress	3	10.7	11	6.0	5	6.4	1	2.2	
Congenital malformation	Others	7	25.0	49	26.9	21	26.9	15	33.3	0.580
	Yes	3	0.7	22	0.9	16	1.3	8	1.1	
Breast feeding	No	402	99.3	2547	99.1	1221	98.7	725	98.9	0.560
	Yes	409	98.8	2554	98.6	1228	98.6	724	98.0	
Neonatal sex	Female	5	1.2	35	1.4	18	1.4	15	2.0	0.004
	Male	210	51.6	1254	49.2	537	43.9	329	45.4	
		197	48.4	1294	50.8	658	56.1	396	54.6	

Table 4. Demonstrates the relative risks of maternal and neonatal outcome for the BMI categories

Variables	Under weight	Normal	Over weight	Obese
Macrosomia	<i>p</i> =0.019 OR=0.498 CI=(0.890-0.278)	1	<i>p</i> =0.008 OR=1.427 CI=(11.099-1.854)	<i>p</i> =0.001 OR=1.820 CI=(1.354-2.447)
Cesarean	<i>p</i> =0.011 OR=0.724 CI=(0.564-0.930)	1	<i>p</i> =0.001 OR=1.739 CI=(1.473-2.053)	<i>p</i> =0.001 OR=2.141 CI=(1.737-2.638)
Gestational diabetes	<i>p</i> =0.195 OR=2.107 CI=(0.683-6.498)	1	<i>p</i> =0.065 OR=2.011 CI=(0.956-4.231)	<i>p</i> =0.139 OR=1.916 CI=(0.810-4.529)
Unwanted pregnancy	<i>p</i> =0.872 OR=0.950 CI=(0.510- 1.771)	1	<i>p</i> =0.474 OR=1.151 CI=(0.783-1.693)	<i>p</i> =0.022 OR=1.633 CI=(1.072-2.486)
Hypertension	<i>p</i> =0.764 OR=1.211 CI=(0.347- 4.222)	1	<i>p</i> =0.011 OR=2.499 CI=(1.231-5.072)	<i>p</i> =0.006 OR=3.011 CI=(1.372-6.604)
Hemoglobin < 10.5	<i>p</i> =0.842 OR=0.966 CI=(0.687-1.357)	1	<i>p</i> =0.002 OR=0.678 CI=(0.532-0.864)	<i>p</i> =0.007 OR=0.661 CI=(0.490-0.893)
Abortion	<i>p</i> =0.019 OR=0.498 CI=(0.890-0.278)	1	<i>p</i> =0.008 OR=1.427 CI=(1.099-1.854)	<i>p</i> =0.001 OR=1.820 CI=(1.354-2.447)
Male sex	<i>p</i> =0.011 OR=0.724 CI=(0.564-0.930)	1	<i>p</i> =0.001 OR=1.739 CI=(1.473-2.053)	<i>p</i> =0.001 Or=2.141 CI=(1.737-2.638)
Low birth weight	<i>p</i> =0.195 OR=2.107 CI=(0.683-6.498)	1	<i>p</i> =0.065 OR=2.011 CI=(0.956-4.231)	<i>p</i> =0.139 OR=1.916 CI=(0.810-4.529)
Preeclampsia	<i>p</i> =0.872 OR=0.950 CI=(0.510-1.771)	1	<i>p</i> =0.474 OR=1.151 CI=(0.783-1.693)	<i>p</i> =0.022 OR=1.633 CI=(1.072-2.486)
Still birth	<i>p</i> =0.764 OR=1.211 CI=(0.347-4.222)	1	<i>p</i> =0.011 OR=2.499 CI=(1.231-5.072)	<i>p</i> =0.006 OR=3.011 CI=(1.372-6.604)

Discussion

In this study, the thin, overweight and obese BMI in subjects were 8.2, 25 and 15.4%, respectively. In total, 48.6% of those women had abnormal BMI which shows high prevalence of abnormal BMI in the subjects of the study. Overweighed and obese women had low level of education and belonged to higher socioeconomic class and aged group. The results of our study are in line with

the findings of El-Gilany et al. and Chen et al. [6, 14]. In the present study, an increase in the number of pregnancy and delivery correlated with an increase in BMI (*p*=0.001) being in agreement with the results of some other studies [14-16].

The maximum frequency of anemia observed in women with thin BMI. The more BMI, the less frequency of anemia, as 17% of women with thin BMI and 12% with fat BMI had anemia during 26-30 weeks of pregnancy.

The results of logistic regression indicated that the odds of developing anemia in women with overweight and fat BMI was less than 1 and it is even less compared with normal women. Since all the women with high BMI belonged to higher socioeconomic class, access to good diet might be the reason for their decrease in anemia. These results concur with the findings of Sebire et al. but are in contrast to El-Gilany et al. findings [6, 16].

Obesity has been recognized as a risk factor of insulin resistance and type 2 diabetes [17-19]. On the other hand, normal pregnancy develops another form of insulin resistance in the body. Therefore, it increases 40-50% during normal pregnancy, which in turn, increases the probability of having pregnancy diabetes in overweight and fat women [2]. Although the differences were not statistically significant, however, the frequency of pregnancy diabetes increased with increasing the BMI (i.e. 0.9% in thin, 1.3% in overweight and 1.7% in fat women) in present study. In the Heinrich et al. study, the chance of developing pregnancy diabetes in fat women was as four times high as that of normal ones [4]. In a study by Sakamoto et al., the frequency of gestational diabetes did not have any significant relationship with different levels of BMI and the probability of gestational diabetes was the same for both thin and fat women [20].

Obesity is accompanied by dyslipidemia and chronic oxidative inflammatory which may cause endothelial damage and then preeclampsia [20]. In a number of studies, the odds of having preeclampsia and high blood pressure in women with slightly overweight and fat BMI was significantly high [2, 6, 14, 16]. In our study based on the results of logistic regression, fat and overweight BMI were among the important factors causing high blood pressure and preeclampsia so much so that in fat and overweight women the odds of developing high blood pressure were 2.5 and 3 and preeclampsia were 3 and 4.6 times, respectively.

Different studies confirmed the relationship between high levels of BMI and caesarian. These studies have put forward the accumulation of fat tissues in abdomen and hip of fat women and fetal macrosomia as the causing factors of delivery progress and the increase in caesarian delivery [4, 6, 19, 21, 22]. Similar to these findings, our study showed that fat and overweight BMI is an important risk factor for caesarian. The chance of caesarian in overweight women was 1.7 and in fat ones it was 2 times. Also, the probability of fetal macrosomia in women with overweight and fat BMI was 1.4 and 1.8 times, respectively.

In this study, frequency of abortion and still birth increased with increasing the BMI. The frequency of abortion in thin women was 0.6%, in overweight and fat ones 1.5% and 2.2% respectively. In terms of still birth, the respective values were 0.3, 0.6 and 1.3% respectively. The results of the logistic regression showed that thin and slightly overweight BMI were not important risk factors for still birth. However, the probability of still birth in fat women increased to 2.6 times. The main cause of still birth in women with higher BMI is not yet known. Researchers believe in such factors as high blood

pressure, diabetes, atherosclerosis and maternal apnea while sleeping for still birth in fat women [5]. In of Mantakas et al., thin and slightly overweight BMI did not have any effect on still birth delivery. However, it was 2.6 times for fat women being in line with the findings of our study [7]. There was not a significant relationship between maternal BMI and still birth in El-Gilany et al.'s study [6].

In the present study, the frequency of unwanted pregnancy in women with overweight and fat BMI was significantly high so long as the probability of unwanted pregnancy was 1.6 times in fat ones. In Holt et al.'s study, it was also 1.6 for fat women. In a study by Zieman et al., there was a direct relationship between a rise in BMI and failure in contraceptive methods. There was not an increase in the frequency of unwanted pregnancy in overweight women in Kaneshiro et al.'s study [20].

In El-Gilany et al.'s study, the probability of low birth weight in women with thin BMI was 2.3 and with an increase in BMI level, there was a decrease in low birth weight risk [6]. Cedergren et al., Sakamoto et al. and Heinich et al. came up with similar findings [4, 6, 18, 21] which are in agreement with our findings in this study. In our study, the most frequent preterm delivery was observed in women with thin BMI and as BMI increased we observed a decrease in the number of preterm delivery, although the differences were not statistically significant. In Jensen et al., and El-Gilany et al.'s study, there was not statistically significant relationship between BMI and preterm delivery, nevertheless, the frequency of preterm delivery was high in women with thin BMI [6, 23]. Recently, there have been some studies on the relationship between high BMI and preterm delivery. In the studies of Chen et al. and Heinrich et al., the probability of preterm delivery in fat and overweight women was 1.5 fold [4, 14]. Researchers believe that high BMI is a preventative factor for preterm delivery but such disorders as diabetes, preeclampsia and high blood pressure can be accompany with high body mass index leading to preterm delivery [5]. In the present study, the frequency of congenital abnormalities and failure in breastfeeding was higher in women with overweight and fat BMI but it was not statistically significant. In some studies, the relationship between high BMI and inborn irregularity and also nervous disorders has been addressed [24, 25]. In Hazel et al.'s study, unsuccessful breastfeeding was rampant among fat women [26]. The mentioned case studies have been done with large samples and the small number of newborns with congenital malformations in our study might be the cause of this weak relationship.

We found that jaundice, hospitalization and breathing distress in newborns of women with thin BMI to be high but it was not significant. As preterm delivery and low birth weight are more prevalent in women with thin BMI, the findings of the present study are verifiable.

In the present study, for any unknown reason, male gender in women with overweight and fat BMI was significantly higher (odds ratio=1.2). We could not find any research on the relationship between BMI and the

gender of the newborns. Further research is needed in this area.

The results of our study indicated a high prevalence of abnormal BMI in pregnant women and an increase in the risk of maternal and newborn disorders. Although, all the subjects in our study had received adequate pregnancy care, there was an increase in the risk of fetal macrosomia, caesarian delivery, unwanted pregnancy, high blood pressure, preeclampsia and still birth in overweight and obese women and low birth weight in thin ones. The mentioned disorders have a major impact on maternal and newborn mortality rate. Diminishing this process is the main challenge of national health system. Seeking guidance and accessing normal BMI before pregnancy can significantly reduce the mentioned disorders and improve maternal and newborn health.

Some studies have concluded that gaining ideal weight on the part of pregnant women leads to the least amount of maternal and newborn disorders especially in obese women [27]. American Medical Organization has determined the ideal weight gain as 5-9 kg in fat women [2, 28]. But, recent studies have come to the conclusion that such amount of weight gain has changed these days [27]. In the present study, due to lack of access to mothers

after delivery, examining the effect of pregnancy weight gain on maternal and newborn was not possible. Further studies on the maternal and fetal disorders and the range of weight variation can help in determining the range of weight gain with little maternal and newborn disorders.

Acknowledgements

The present research was conducted after receiving authorization No. 10239 in Feb 2009 from research council of Ardabil University Medical Sciences. We thank all those people working in Ardabil health centers who helped us throughout this research.

Authors' Contributions

All authors had equal role in design, work, statistical analysis and manuscript writing.

Conflict of Interest

The authors declare no conflict of interest.

Funding/Support

Ardabil University of Medical Science.

References

- Albers LL, Greulich B, Peralta P. Body mass index, midwifery intra partum care, and childbirth laceration. *J Midwifery Women Health* 2006; 51(4): 249-53.
- Yogev Y, Visser G. Obesity, gestational diabetes and pregnancy outcome. *Semin Fetal Neonatal Med* 2009; 14(2): 77-84.
- Byers BD, Betancourt A, Lu F, et al. The effect of pregnancy obesity and sFlt-1-induced preeclampsia-link syndrome on fetal programming of adult vascular function in a mouse model. *Am J Obstet Gynecol* 2009; 200(4): 432.
- Demont-Heinrich C, Hansen M, McCullon A and Archer L. The association of pregnancy body mass index and adverse maternal and prenatal outcomes. *Colorado Dept Public Health Environ* 2009; 69: 1-7.
- Rowlands I, Grave N, Jersey S, et al. Obesity in pregnancy: Outcome and economics. *Semin Fetal Neonatal Med* 2010; 15(2): 94-99
- el-Gilany A, Hammad S. Body mass index and obstetric outcomes in Saudi Arabia: A prospective cohort study. *Ann Saudi Med* 2010; 30(5): 376-380.
- Mantakas A, Farrell T. The influence of increasing BMI in nulliparous women on pregnancy outcome. *Eur J Obstet Gynecol Reprod Biol* 2010; 153(1): 43-6.
- Tabandeh A, Kashani E. [The association of maternal body mass index and increasing weight gain during pregnancy with maternal or neonatal adverse outcome] Persian. *J Gorgan Univ Med Sci* 2007; 9(1): 20-24.
- Mostafazadeh F, Kahnemoie F, Mohammadi R, et al. [Prevalence of obesity in pregnant women] Persian. *J Ardabil Univ Med Sci* 2010; 12(1): 39-43.
- Nematollahzadeh M, Ziaei S, Kazemnejad A. [The association of pre pregnancy body mass index and pregnancy weight gain with preterm delivery] Persian. *Zahedan J Res Med Sci* 2010; 12(5): 89-94
- Garshasbi A, Solbi Z, Fagihzadeh S, et al. [Effect of body mass index on pregnancy outcome] Persian. *Daneshvar Med J* 2007; 16(77): 33-39.
- Valafard SH. [Integration of maternal health care] Persian. 6th ed. Tehran: Ministry of Health and Medical Education Press; 2010.
- Torabi P, Sheikholeslam R, Minaei M, editors. [National guidelines of nutrition during pregnancy and lactation for physicians and bachelors of midwifery] Persian. 1th ed. Tehran: Ministry of Health and Medical Education Press; 2005.
- Chen Z, Du J, Shao L, et al. Pregnancy body mass index, gestational weight gain and pregnancy outcome in China. *Int J Gynecol Obstet* 2009; 109(1): 41-44.
- Dietz P, Callaghan W, Sharma A. High pregnancy weight gain and risk of excessive fetal growth. *Am J Obstet Gynecol* 2009; 201(1): 51.
- Sebire NJ, Jolly M, Harris JP, et al. Maternal obesity and pregnancy outcome: A study of 287213 pregnancies in London. *Int J Obes Relat Metab Disord* 2001; 25(8) :1175-82.
- Martin RR, Hyde MJ, Modi N. Maternal obesity and infant outcome. *Early Hum Dev* 2010; 86(11): 715-22.
- Tsakamoto H, Fukuoka H, Inoue K, et al. Restricting weight gain during pregnancy in Japan: A controversial factor in reducing perinatal complications. *Eur J Obstet Gynecol Reprod Biol* 2007; 133(1): 53-9.
- Bergholt T, Lim L, Jorgensen J and Roboson M. Maternal body mass index in the first trimester and risk of cesarean delivery in nulliparous women in spontaneous labor. *AM J Obstetric Gynecol* 2007; 196(1): 163.
- Cunningham FG, Leveno KJ, Hauth JC, editors. *Williams obstetrics*. 23th ed. New York: McGraw Hill; 2010.

21. Cedergren M. Effects of gestational weight gain and body mass index on obstetric outcome in Sweden. *Int J Gynaecol Obstet* 2006; 93(3): 269-274.
22. Vahratian A, Siega-Riz A, Savitz DA and Zhang J. Maternal pre-pregnancy overweight and obesity and the risk of cesarean delivery in nulliparous women. *Ann Epidemiol* 2005; 15(1): 467-474
23. Jensen DM, Sorensen B, Molsted-Pedersen L, et al. Pregnancy outcome and pregnancy body mass index in 2458 glucose-tolerant Danish women. *AM J Obstetric Gynecol* 2003; 189(1): 239-44.
24. Rasmussen SA, Chu S, Kim S, et al. Maternal obesity and risk of neural tube defects: A meta-analysis. *AM J Obstetrics Gynecol* 2008; 198(6): 611-619.
25. Gilboa S, Correa A, Botto LD, et al. Association between pre pregnancy body mass index and congenital heart defects. *AM J Obstetrics Gynecol* 2010; 202(1): 51.
26. Oddy WH, Jianghong L, Landsborough L, et al. The association of maternal overweight and obesity with breast feeding duration. *J Pediatr* 2006; 149(2): 185-191.
27. Oken E. Excess gestational weight gain amplifies risks among obese mothers. *Epidemiology* 2009; 20(1): 82-83.
28. Zilko CM, Rehkopf D, Abrams B. Association of maternal gestational weight gain with short and long term maternal and child health outcomes. *Am J Obstetrics Gynecol* 2010; 202(6): 574-578.

Please cite this article as: Alijahan R, Nakhostin B, Salimi S, Hazrati S. Association of maternal body mass index with adverse maternal and prenatal outcomes. *Zahedan J Res Med Sci (ZJRMS)* 2013; 15(9): 56-62.