Published online 2019 August 19.

**Research Article** 

# The Apoptotic Pathway Induced by Vincristine in Mouse Proliferating and Resting Normal Lymphocytes and Lymphoma Cell Line

Mohaddese Shahhosseini<sup>1,2</sup>, Susan Kaboudanian Ardestani<sup>2</sup> and Roya Yaraee <sup>[0],\*</sup>

<sup>1</sup>Department of Immunology, Medical Faculty, Shahed University, Tehran, Iran <sup>2</sup>Institute of Biochemistry and Biophysics. University of Tehran. Tehran. Iran

Corresponding author: Department of Immunology, Medical Faculty, Shahed University, Persian Gulf Highway, In Front of Imam Khomeini Holy Shrine, Tehran, Iran. Tel: +98-2151212652, Email: ryaraee@yahoo.com

Received 2016 December 05; Revised 2019 April 28; Accepted 2019 July 22.

# Abstract

**Background:** Most anti-cancer drugs target mitosis and induce apoptosis in cancerous cells. In the immune system, proliferation and apoptosis of lymphocytes are indeed essential modulating elements.

**Objectives:** In this study, we have investigated the effect of vincristine on normal resting and normal proliferating lymphocytes comparing with cancerous cells.

**Methods:** Resting and proliferating splenocytes from mice and BCL1 (mouse lymphoma cell line) were cultured with different concentrations of vincristine for 48 hours, and cell lysates were prepared. The activity of caspases 3, 8, and 9 in cell lysates was measured using specific chromogenic substrates DEVD-pNA for caspase 3, IETD-pNA for caspase 8, and LEHD-pNA for caspase 9, the activity calculated as  $\mu$ mol/min/mg protein.

**Results:** In the BCL1 cell line, the activity of both caspases 8 and 9 and caspase 3 increased at the presence of vincristine (5  $\mu$ g/mL). In resting splenocytes, however, only a mild increase in caspase 9 activity was observed without any change in the activity of caspases 8 or 3. In the same situation, the activity of caspase 3 and 9 (but not caspase 8) elevated in proliferating cells exposed to vincristine. Nearly similar results were obtained with higher concentrations of vincristine (up 20  $\mu$ g/mL).

**Conclusions:** The results suggesting that vincristine may induce internal pathways of apoptosis in normal and cancerous cells while extrinsic pathway was induced in cancerous cells. On the other hand, the effects are highly dependent on the activation status of normal cells, and affirms that responding immune cells should be more seriously noticed when side effects of anticancer drugs are estimated.

Keywords: Vincristine, Tumor Cell Line, Lymphocytes, Caspase

## 1. Background

Vincristine, as a key drug for the treatment lymphoid malignancies. belongs to Vinca alkaloids - the secondmost-used class of cancer drugs. It was widely used as a potent chemotherapeutic agent in the treatment of various cancers such as lymphoma; however, it has common side effects including nausea, vomiting, weight loss, diarrhea, bloating, mouth sores, dizziness, headache, hair loss, and also neutropenia and peripheral neuropathy (1-3).

Many anti-cancer chemotherapy drugs target mitosis, hence induce apoptosis in cancerous cells, such as vincristine, which induces mitotic arrest as a microtubule inhibitor (4, 5). Therefore, cells undergo more apoptosis when they are more dividing. However, many studies regarding the effects of vincristine may notice only normal resting lymphocytes e.g. peripheral blood mononuclear cells, so the effect on normal activated lymphocytes remains unseen whereas these normal proliferating cells may be more susceptible to cytotoxic effects of the drug. It is noteworthy to consider that proliferation and apoptosis of lymphocytes are essential parts of the immune modulation and apoptosis induction by chemotherapy drugs include the normal proliferating lymphocytes responsive to malignancies or other invaders.

Caspases (cysteine-dependent asparagine-oriented protease) are the main players in apoptosis. Two distinct yet interconnected signaling pathways control apoptosis by activating caspases. The intrinsic apoptotic pathway engages caspase 9 via members of the BCL-2 protein family and the mitochondria in reaction to severe cellular damage or stress and is mediated by a multimeric adaptor complex known as the Apaf-1 apoptosome (6). The extrinsic pathway however usually activates caspase 8 via

Copyright © 2019, Zahedan Journal of Research in Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

cell-surface death receptors (7). The activation of caspase 8 may lead to caspase 9 activation as well (8, 9). Caspase 8 has anti-tumor roles, and increased levels of caspase 3 in tumor cells induce apoptosis (10, 11).

The detailed mechanism involved in the induction of apoptosis by vincristine not yet well defined, but it is caspase-dependent (12). Its apoptosis induction or caspase activation in normal proliferating lymphocytes is also unclear. We have previously shown that 10  $\mu$ g/mL (and more) of vincristine caused cell death in both resting and proliferating lymphocytes in vitro and the cells undergo apoptosis (double staining with acridine orange and ethidium bromide) (13). However, the apoptotic pathway induced by vincristine in proliferating and resting normal lymphocytes and lymphoma cell lines is not clearly compared.

# 2. Objectives

The aim of this study is to evaluate the effect of vincristine on the activity of main apoptotic caspases (3, 8 and 9) in resting and proliferating lymphocytes and cancerous lymphoma cell line (B cell lymphoma BCL1).

# 3. Methods

## 3.1. Cells and Culture

## 3.1.1. Splenocytes

The spleen of 8 - 10-weeks BALB/c mice was removed under sterile conditions and the splenocytes separated as cell suspension and centrifuged, then resuspended in 2 mL Tris-buffered (0.2%) ammonium chloride (0.83%) and incubated for 2 minutes at room temperature to remove red blood cells. Immediately after that, 2 mL of fetal bovine serum (FCS) (Gibco) was added, centrifuged at 300 g for 10 minutes, and washed twice with RPMI medium (Gibco). Cell viability, as determined by the Trypan blue exclusion method, exceeded 95%. The cells were resuspended in 10% FCS RPMI medium and cultured in 96-well microplates at 2  $\times 10^5$  cells per well.

Vincristine, at various concentrations (3.7, 5, 10 and 20  $\mu$ g/mL depending on work design), were added with or without 25  $\mu$ g/mL concanavalin A (Con A) as a lymphocyte stimulator (5 wells were assigned for each condition) and incubated for 48 hours at 37°C, 5% CO<sub>2</sub>.

# 3.1.2. Cell Line

BCL1 cells were purchased from the Pasteur Institute Cell Bank (Tehran, Iran, code C551). The cells were cultured and passaged at 37°C, 5% CO<sub>2</sub> in 10% FCS supplemented RPMI1640 Medium.  $2 \times 10^4$  cells were added to each well of the 96-well microplate and cultured overnight, then, the supernatants were removed and the desired concentrations of vincristine were added (5 wells for each condition) and incubated for 48 hours at 37°C, 5% CO<sub>2</sub>.

#### 3.2. Cell Lysates

Briefly, after treatment with desired doses of vincristine for a specific time period (48 hours), the cells were washed and then lysed in a lysis buffer composed of 20 mM PIPES (piperazine-N, N'-bis (2-ethanesulfonic acid)), 10 mM KCl (potassium chloride), 2 mM MgCl<sub>2</sub> (magnesium chloride), 4 mM DTT (Dithiothreitol), pH 7.4 and 20  $\mu$ L antiprotease PMSF (phenylmethylsulfonyl fluoride, and 1  $\mu$ L leupeptin were added per 1 mL of lysing buffer). To obtain total protein of each sample, 100  $\mu$ L of this cocktail was added to each 10<sup>6</sup> splenocytes or 10<sup>5</sup> BCL1 and incubated on ice for 30 minutes. Then, the cells were passed from an insulin syringe 10 times and the Lysates were clarified by centrifugation at 3500 g (4°C) for 15 minutes. The supernatants were collected and froze at -70°C and the total protein of each sample was determined by the Bradford Method (14).

#### 3.3. Caspase Activity

The caspase activity was assessed using specific colorimetric substrates labeled with the chromophore p-nitroaniline (pNA). Fifty  $\mu$ L of the cell lysates was mixed with 50  $\mu$ L of caspase assay buffer composed of 100 mM NaCl (sodium chloride), 50 mM Hepes (N-2hydroxyethylpiperazine-N'-2-ethanesulfonic acid), 0.1% CHAPS (3- ((3-Cholamidopropyl) dimethylammonio)-1propanesulfonic acid), 0.1 mM EDTA (ethylene diamine tetraacetic acid), 10 mM DTT, in 10% Glycer, pH = 7.40l), 5  $\mu$ L of caspase 3 substrate DEVD-pNA (N-Acetyl-Asp-Glu-Val-Asp p-nitroanilide) was added and incubated at 37°C for 4 hours. Hydrolysis of DEVD-pNA by caspase 3 released free pNA that produces a yellow color detectable at 405 nm. The cleavage of the peptide substrate was monitored at 405 nm using a 96-well microplate reader at time intervals of 10 minutes. The same protocol performed for Caspase -9 but the specific substrate replaced by LEHD-pNA (N-Acetyl-Leu-Glu-His-Asp p-nitroanilide). caspase assay buffer for caspase 8 was a little different (100 mM NaCl, 50 mM Hepes, 0.1% CHAPS, 0.5 mM EDTA, BSA 0.05 %, 1 mM DTT, in 10% Glycer, pH = 7.4ol) and IETD-pNA (N-Acetyl-Ile-Glu-Thr-Asp-p-nitroanilide) was used as a specific substrate (15). Supernatants of untreated cells were considered as a negative control. Control 1 (without substrate) and control 2 (without supernatant) were included as well. Caspase activity was measured using a standard curve with known concentrations of specific substrate, and the

activity according to  $\mu$ mol/min/mg protein was calculated whenever necessary. The experiments were conducted on three independent samples of cultured cells.

#### 3.4. Statistical Analysis

Statistical differences were assessed by the ANOVA (analysis of variance) and Tukey's post hoc test and P < 0.05 was considered statistically significant.

# 4. Results

#### 4.1. BCL1 (Cancerous Cell): Caspase 8, 9 and 3 Activation

As shown in Figure 1A, in vincristine treated cells (5  $\mu$ g/mL, 48-hour incubation), the activity of caspase 3 was significantly augmented as assayed in cell lysates. Vincristine also causes an increase in the activity of caspases 8 and 9 detectable 20 minutes after adding substrate to cell lysate (Figure 1B and C).

The activity of caspases was calculated using a standard curve with known concentrations of substrates and presented as  $\mu$ mol/min/mg protein (table of Figure 1). In the normal culture of BCL1 cells, no caspase 3 activity was observed, however, at the presence of vincristine, the activity of caspase 3 was  $4.87 \pm 0.88 \ \mu$ mol/min/mg protein; more than each of caspases 8 or 9 ( $3.29 \pm 0.06$  and  $3.53 \pm 0.09 \ \mu$ mol/min/mg protein, respectively, P <). In lower concentration of vincristine ( $3.7 \ \mu$ g/mL), the activity of both caspases 8 and 9 were less but detectable ( $2.18 \pm 0.87$  and  $1.58 \pm 0.07 \ \mu$ mol/min/mg protein respectively) and no caspase 3 activation was observed.

# 4.2. Normal Resting Splenocytes: Low-Level Activation of Caspase 9

As shown in Figure 2, vincristine at 5 µg/mL induced a low-level activation of Caspase 9 in normal resting lymphocytes (from mouse spleen), but no activity of caspase 8 and capase-3 was observed.

Higher concentrations of vincristine were also examined and the caspase activity was calculated as  $\mu$ mol/min/mg protein presented in table of Figure 2. As shown, at the absence of vincristine (normal culture), resting lymphocytes have no caspase 3 activity, however, at the presence of vincristine, the activity of caspase 3 was observed at 10 and 20  $\mu$ g/mL concentration (5.95  $\pm$  0.50 and 6.73  $\pm$  0.28  $\mu$ mol/min/mg protein respectively), but not at 5  $\mu$ g/mL of vincristine. There was no increase in caspase 8 of normal resting cells in any concentration of vincristine used, but caspase 9 activity was elevated dose-dependently i.e. 1.99  $\pm$  0.29, 3.93  $\pm$  0.12 and 13.49  $\pm$  0.63  $\mu$ mol/min/mg protein at 5, 10 and 20  $\mu$ g/mL of vincristine respectively.

4.3. Normal Proliferating Splenocytes: Activation of Caspase 9 and 3

As shown in Figure 3, in normal proliferating lymphocytes, vincristine at 5  $\mu$ g/mL induced both caspase 3 and caspase 9 activity, but not caspase 8.

Vincristine at higher concentrations (8.5 and 20  $\mu$ g/mL) had no observable effect on caspase 8 activity, however, it significantly augmented the activity of caspase 9 and caspase 3 (table of Figure 3). The highest level of caspase activity has been observed in caspase 9 activity when vincristine was administered at 20  $\mu$ g/mL (23.81  $\pm$  0.02  $\mu$ mol/min/mg protein).

#### 4.4. Caspase Activities

A comparison between the effect of vincristine (5  $\mu$ g/mL) on various caspases (3, 8, and 9) and various cells (cancerous BCL1, normal resting and activated lymphocytes) is presented in Figure 4A. The activity of caspase 3 and 9 in activated lymphocytes is clearly comparable with BCL1 cells, although resting cells have no caspase 3 activity and caspase 8 activity is seen only in BCL1. As shown in Figure 4B, caspases 3 and 9 are activated at higher concentrations (20  $\mu$ g/mL) both in resting and stimulated cells, although the activity is about twice in activated cells.

#### 5. Discussion

Apoptosis induction is one of the important mechanisms regarding cancer therapy (16). Vincristine bind to  $\beta$ tubulin close to the guanosine triphosphate (GTP)-binding sites (the vinca domain) at the  $\beta$ - $\alpha$ -tubulin heterodimers (17). Prolonged mitotic arrest leads to phosphorylationmediated inactivation of BCL-2 and BCL-XL. Inactivation of antiapoptotic BCL-2 proteins promotes activation of BAX (BCL2-associated X protein) and BAK (Bcl-2 homologous antagonist killer), cleavage of caspase-9 and -3 and caspasedependent apoptosis (18). It induces distinct death programs in primary ALL cells depending on cell-cycle phase, and cells in G1 are particularly susceptible to perturbation of interphase microtubules (19). Pharmacokinetic studies of vincristine in patients with cancer have shown variable characteristics (few minutes to hundred hours in serum) and also neurotoxic side effects. It is able to accumulate a few hundred times more than extracellular concentrations (20-23).

Apoptosis and proliferation in lymphocytes are among the most important features of the immune system (24, 25). Lymphocytes in response to invaders (including malignancies) start to proliferate before any effector function (26, 27). Therefore, anti-tumor drugs may affect normal



**Figure 1.**  $2 \times 10^4$  BCL1 cells were incubated with vincristine at 5  $\mu$ g/mL for 48 hours, then, cell lysates were prepared and colorimetric assay of caspase activity was performed using DEVD-pNA for caspase 3 (A), IETD-pNAfor caspase 8 (B), and LEHD-pNA for caspase 9 (C) as a substrate. The absorbance was read at 10 min intervals. Supernatants of untreated cells were (without Vincristine) considered as a negative control, the sample without substrate was designed as "control 1" and without supernatant as "control 2". The results are presented as mean  $\pm$  SD (very low SD in some experiments makes the bars hardly visible) and the activity of caspases 3, 8, and 9 in BCL1 lymphoma cell line is shown as  $\mu$ mol/min/mg protein (table).

proliferating lymphocytes much more than normal resting lymphocytes, however, these normal resting cells from the peripheral blood are usually used to study the adverse effect of vincristine on lymphocytes (28, 29). Thus, evaluating the effect of an anti-tumor drug (especially antilymphoma) on activated normal lymphocytes could be informative.

We have previously reported that different concentrations of vincristine induce cell death and apoptosis in resting and proliferating spleen lymphocytes. The cytotoxicity of vincristine was determined by the MTT assay and the percentage of apoptotic cells were also determined using double staining acridine orange and ethidium bromide. The toxic effect of vincristine on normal cells was highly dependent on the time and activation status of the cells. The IC50-according to MTT test - was about 3.6  $\mu$ g/mL for BCL1 and 10.6 and 8.5  $\mu$ g/mL for resting and proliferating lymphocytes respectively (13). In this study, we compared the effect of vincristine on the activity of main apoptotic caspases (3, 8, and 9) in resting and proliferating lymphocytes in addition to cancerous lymphoma cell line (BCL1).

According to our results, vincristine (5  $\mu$ g/mL) induces the activity of both caspase 8 and 9 (and caspase 3) in BCL1 cell line. These results are in agreement with various studies reporting that vincristine induces caspase 3 (or 7) and caspase 8 activation in various cell lines (30-33) and promoted the expression of caspase-9 (and caspase-3) in human neuroblastoma cells (34).

It is noteworthy that lower concentration causes gently increase in the activity of caspases 8 and 9 in BCL1 cell line but did not results in caspase 3 activation immediately. It may be due to weak activation of caspases unable to lead to reach the threshold to activate caspase 3 (35). There is an in vitro study that shows that increased cytotoxicity with alterations in G1 and S cell cycle phases may occur without detectable differences in apoptosis (36).

In resting splenocytes only a mild increase in caspase 9 activity was observed without any change in the activity of caspase 8 or 3. It can be attributed to the low toxicity of vincristine on normal lymphocytes. At the same situation, the activity of caspase 3 and 9 elevated in proliferating cells exposed to vincristine (1.5 fold). The results indicate that the activated lymphocytes are more susceptible to apoptosis. The same results were obtained with higher concentrations of vincristine (up 20  $\mu$ g/mL) i.e. no caspase 8 activity in non-cancerous splenocytes (resting or proliferating) and augmented activity of caspase 9 and 3 in both. It can be concluded likewise that the caspase 8 pathway was not easily activated in normal lymphocytes by vincristine.

Regarding the importance of lymphocytes (and their



**Figure 2.** Vincristine at 5  $\mu$ g/mL were added to 2  $\times$  10<sup>5</sup> splenocytes and cultured for 48 hours, cell lysates prepared and caspase activity was assayed colorimetric using DEVD-pNA for caspase 3 (A), IETD-pNA for caspase 8 (B), and LEHD-pNA for caspase 9 (C) as a substrate. The absorbance was read at 10 min intervals. Supernatants of untreated cells (without Vincristine) were considered as a negative control, the sample without substrate designed as "control 1" and without supernatant as "control 2". The results are presented as mean  $\pm$  SD (very low SD in some experiments makes the bars hardly visible) and the activity of caspases 3, 8, and 9 in BCL1 lymphoma cell line is shown as a  $\mu$ mol/min/mg protein (table).

proliferation), in response to various pathogens and also tumor cells, paying attention to the activation status of these cells leads to more admissible results.

# Acknowledgments

We thank our colleagues from Institute of Biochemistry and Biophysics, University of Tehran for their assistance especially with caspase activity assays.

#### Footnotes

Authors' Contribution: Study concept and design: Roya Yaraee, Susan Kaboudanian and Mohaddese Shahhosseini. Acquisition of data: Mohaddese Shahhosseini. Analysis and interpretation of data: Mohaddese Shahhosseini. Drafting of the manuscript: Roya Yaraee, Susan Kaboudanian and Mohaddese Shahhosseini. Critical revision of the manuscript for important intellectual content: Roya Yaraee. Statistical analysis: Roya Yaraee, Susan Kaboudanian and Mohaddese Shahhosseini. Administrative, technical, and material support: Roya Yaraee and Susan Kaboudanian. Study supervision: Roya Yaraee.

**Conflict of Interests:** The authors declared no conflict of interests.

**Ethical Approval:** This article has a license from the Research Ethics Committee of Shahed University, code IR.Shahed.REC.1395,23.

**Funding/Support:** This article is part of M.Sc. thesis approved at and supported by Medical Faculty of Shahed University.





**Figure 3.** Vincristine at 5  $\mu$ g/mL added to 2 × 10<sup>5</sup> ConA-stimulated splenocytes and cultured for 48 hours, cell lysates prepared and caspase activity was assayed colorimetric using DEVD-pNA for caspase 3 (A), IETD-pNA for caspase 8 (B), and LEHD-pNA for caspase 9 (C) as a substrate. The absorbance was read at 10 min intervals. Supernatants of untreated cells (without vincristine) was considered as a negative control, the sample without substrate designed as "control 1" and without supernatant as "control 2". The results are presented as mean  $\pm$  SD (very low SD in some experiments makes the bars hardly visible) and the activity of caspases 3, 8, and 9 in BCL1 lymphoma cell line is shown as  $\mu$ mol/min/mg protein (table).



**Figure 4.** Comparing the activation of caspases with vincristine at 5  $\mu$ g/mL (A) or 20  $\mu$ g/mL (B) administered in vitro. Briefly 2 × 10<sup>4</sup> BCL1 or 2 × 10<sup>5</sup> splenocytes resting or stimulated (with ConA) were cultured at the presence or absence of vincristine for 48 hours; cell lysates were prepared and caspase activity was assayed colorimetrically and the caspase activity was calculated using standard curves (fold increase according to resting lymphocytes at 5  $\mu$ g/mL vincristine).

#### References

- Moudi M, Go R, Yien CY, Nazre M. Vinca alkaloids. Int J Prev Med. 2013;4(11):1231–5. [PubMed: 24404355]. [PubMed Central: PMC3883245].
- Han Y, Smith MT. Pathobiology of cancer chemotherapy-induced peripheral neuropathy (CIPN). *Front Pharmacol.* 2013;4:156. doi: 10.3389/fphar.2013.00156. [PubMed: 24385965]. [PubMed Central: PMC3866393].
- Wilson WH. Treatment strategies for aggressive lymphomas: what works? *Hematology Am Soc Hematol Educ Program*. 2013;2013:584–90. doi: 10.1182/asheducation-2013.1.584. [PubMed: 24319235].
- Chumduri C, Gillissen B, Richter A, Richter A, Milojkovic A, Overkamp T, et al. Apoptosis resistance, mitotic catastrophe, and loss of ploidy control in Burkitt lymphoma. *J Mol Med (Berl)*. 2015;93(5):559–72. doi: 10.1007/s00109-014-1242-2. [PubMed: 25548804].
- Hensley P, Mishra M, Kyprianou N. Targeting caspases in cancer therapeutics. *Biol Chem*. 2013;**394**(7):831-43. doi: 10.1515/hsz-2013-0128. [PubMed: 23509217]. [PubMed Central: PMC3721733].
- Hu Q, Wu D, Chen W, Yan Z, Yan C, He T, et al. Molecular determinants of caspase-9 activation by the Apaf-1 apoptosome. *Proc Natl Acad Sci U S A*. 2014;111(46):16254–61. doi: 10.1073/pnas.1418000111. [PubMed: 25313070]. [PubMed Central: PMC4246342].
- Ashkenazi A. Targeting the extrinsic apoptotic pathway in cancer: Lessons learned and future directions. J Clin Invest. 2015;125(2):487-

9. doi: 10.1172/JCI80420. [PubMed: 25642709]. [PubMed Central: PMC4319431].

- 8. Ola MS, Nawaz M, Ahsan H. Role of BCl-2 family proteins and caspases in the regulation of apoptosis. *Mol Cell Biochem*. 2011;**351**(1-2):41–58. doi: 10.1007/s11010-010-0709-x. [PubMed: 21210296].
- Creagh EM. Caspase crosstalk: Integration of apoptotic and innate immune signalling pathways. *Trends Immunol*. 2014;**35**(12):631–40. doi: 10.1016/j.it.2014.10.004. [PubMed: 25457353].
- Shalini S, Dorstyn L, Dawar S, Kumar S. Old, new and emerging functions of caspases. *Cell Death Differ*. 2015;22(4):526–39. doi: 10.1038/cdd.2014.216. [PubMed: 25526085]. [PubMed Central: PMC4356345].
- MacKenzie SH, Clark AC. Death by caspase dimerization. *Adv Exp Med Biol.* 2012;**747**:55-73. doi: 10.1007/978-1-4614-3229-6\_4. [PubMed: 22949111]. [PubMed Central: PMC3877935].
- Zhu BK, Wang P, Zhang XD, Jiang CC, Chen LH, Avery-Kiejda KA, et al. Activation of Jun N-terminal kinase is a mediator of vincristine-induced apoptosis of melanoma cells. *Anticancer Drugs*. 2008;**19**(2):189–200. doi: 10.1097/CAD.0b013e3282f3138a. [PubMed: 18176116].
- 13. Shahhosseini M, Kabudanian Ardestani S, Yaraee R. [Comparing the efffect of vincristine on proliferating normal lymphocytes and BCL1 cell line]. *Daneshvar Med*. 2012;**11**:1-11. Persian.
- Rahmani-Nezhad S, Safavi M, Pordeli M, Ardestani SK, Khosravani L, Pourshojaei Y, et al. Synthesis, in vitro cytotoxicity and apoptosis inducing study of 2-aryl-3-nitro-2H-chromene derivatives as potent anti-breast cancer agents. *Eur J Med Chem.* 2014;86:562–9. doi: 10.1016/j.ejmech.2014.09.017. [PubMed: 25216378].
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72:248–54. doi: 10.1006/abio.1976.9999. [PubMed: 942051].
- Goldar S, Khaniani MS, Derakhshan SM, Baradaran B. Molecular mechanisms of apoptosis and roles in cancer development and treatment. *Asian Pac J Cancer Prev.* 2015;16(6):2129–44. doi: 10.7314/apjcp.2015.16.6.2129. [PubMed: 25824729].
- van Vuuren RJ, Visagie MH, Theron AE, Joubert AM. Antimitotic drugs in the treatment of cancer. *Cancer Chemother Pharmacol.* 2015;**76**(6):1101-12. doi: 10.1007/s00280-015-2903-8. [PubMed: 26563258]. [PubMed Central: PMC4648954].
- Czaplinski S, Hugle M, Stiehl V, Fulda S. Polo-like kinase 1 inhibition sensitizes neuroblastoma cells for vinca alkaloid-induced apoptosis. *Oncotarget*. 2016;7(8):8700–11. doi: 10.18632/oncotarget.3901. [PubMed: 26046302]. [PubMed Central: PMC4890998].
- Kothari A, Hittelman WN, Chambers TC. Cell cycle-dependent mechanisms underlie vincristine-induced death of primary acute lymphoblastic leukemia cells. *Cancer Res.* 2016;**76**(12):3553–61. doi: 10.1158/0008-5472.CAN-15-2104. [PubMed: 27197148]. [PubMed Central: PMC4911277].
- Lengsfeld AM, Dietrich J, Schultze-Maurer B. Accumulation and release of vinblastine and vincristine by HeLa cells: Light microscopic, cinematographic, and biochemical study. *Cancer Res.* 1982;**42**(9):3798– 805. [PubMed: 7105045].
- Said R, Tsimberidou AM. Pharmacokinetic evaluation of vincristine for the treatment of lymphoid malignancies. *Expert Opin Drug Metab Toxicol*. 2014;10(3):483–94. doi: 10.1517/17425255.2014.885016. [PubMed: 24512004].
- Mora E, Smith EM, Donohoe C, Hertz DL. Vincristine-induced peripheral neuropathy in pediatric cancer patients. *Am J Cancer Res.* 2016;6(11):2416-30. [PubMed: 27904761]. [PubMed Central:

PMC5126263].

- Visconti R, Grieco D. Fighting tubulin-targeting anticancer drug toxicity and resistance. *Endocr Relat Cancer*. 2017;24(9):T107–17. doi: 10.1530/ERC-17-0120. [PubMed: 28808045].
- Malemud CJ, Haque A, Louis NA, Wang J. Immune response and apoptosis introduction. J Clin Cell Immunol. 2012;1(S3). doi: 10.4172/2155-9899.S3-e001.
- Beug ST, Cheung HH, LaCasse EC, Korneluk RG. Modulation of immune signalling by inhibitors of apoptosis. *Trends Immunol.* 2012;33(11):535–45. doi: 10.1016/j.it.2012.06.004. [PubMed: 22836014].
- Xu G, Shi Y. Apoptosis signaling pathways and lymphocyte homeostasis. *Cell Res.* 2007;**17**(9):759–71. doi: 10.1038/cr.2007.52. [PubMed: 17576411].
- Mognol GP, Carneiro FR, Robbs BK, Faget DV, Viola JP. Cell cycle and apoptosis regulation by NFAT transcription factors: New roles for an old player. *Cell Death Dis.* 2016;7. e2199. doi: 10.1038/cddis.2016.97. [PubMed: 27100893]. [PubMed Central: PMC4855676].
- Zwaan CM, Kaspers GJ, Pieters R, Hahlen K, Janka-Schaub GE, van Zantwijk CH, et al. Different drug sensitivity profiles of acute myeloid and lymphoblastic leukemia and normal peripheral blood mononuclear cells in children with and without Down syndrome. *Blood.* 2002;**99**(1):245–51. doi: 10.1182/blood.v99.1.245. [PubMed: 11756178].
- Vilpo JA, Koski T, Vilpo LM. Selective toxicity of vincristine against chronic lymphocytic leukemia cells in vitro. *Eur J Haematol.* 2000;65(6):370-8. doi: 10.1034/j.1600-0609.2000.065006370.x. [PubMed: 11168494].
- Delgado ME, Olsson M, Lincoln FA, Zhivotovsky B, Rehm M. Determining the contributions of caspase-2, caspase-8 and effector caspases to intracellular VDVADase activities during apoptosis initiation and execution. *Biochim Biophys Acta*. 2013;**1833**(10):2279–92. doi: 10.1016/j.bbamcr.2013.05.025. [PubMed: 23747563].
- Cavaliere V, Papademetrio DL, Lombardo T, Costantino SN, Blanco GA, Alvarez EM. Caffeic acid phenylethyl ester and MG132, two novel nonconventional chemotherapeutic agents, induce apoptosis of human leukemic cells by disrupting mitochondrial function. *Target Oncol.* 2014;9(1):25–42. doi: 10.1007/s11523-013-0256-y. [PubMed: 23430344].
- 32. Shimizu H, Ohgoh M, Momose Y, Nishizawa Y, Ogura H. Massive cell death of cerebellar granule neurons accompanied with caspase-3-like protease activation and subsequent motor discoordination after intracerebroventricular injection of vincristine in mice. *Neuroscience*. 2002;**115**(1):55–65. doi: 10.1016/s0306-4522(02)00403-7. [PubMed: 12401321].
- Tabal S, Elbanani A, Deyab M, Abulayha A. Rituximab increases the cytotoxicities of vincristine and hydroxyurea through caspasedependent and caspase-independent cell death, respectively. *Cancer Biother Radiopharm*. 2015;**30**(3):125–31. doi: 10.1089/cbr.2014.1769. [PubMed: 25714921].
- Tu Y, Cheng S, Zhang S, Sun H, Xu Z. Vincristine induces cell cycle arrest and apoptosis in SH-SY5Y human neuroblastoma cells. *Int J Mol Med.* 2013;**31**(1):113–9. doi: 10.3892/ijmm.2012.1167. [PubMed: 23129065].
- Thomsen ND, Koerber JT, Wells JA. Structural snapshots reveal distinct mechanisms of procaspase-3 and -7 activation. *Proc Natl Acad Sci U S A*. 2013;**110**(21):8477-82. doi: 10.1073/pnas.1306759110. [PubMed: 23650375]. [PubMed Central: PMC3666719].
- 36. Florez MM, Feo HB, da Silva GN, Yamatogi RS, Aguiar AJ, Araujo JP Jr, et al. Cell cycle kinetics, apoptosis rates and gene expressions of MDR-1, TP53, BCL-2 and BAX in transmissible venereal tumour cells and their association with therapy response. *Vet Comp Oncol*. 2017;**15**(3):793–807. doi: 10.1111/vco.12220. [PubMed: 26879698].