



Comparative Analysis of Corticosteroid Therapy in Pregnant Women with COVID-19: Evaluating Glycemic Control and Transient Hyperglycemia

Mohammad Reza Salehi¹, Marjan Ghaemi  ², Sahar Masoumi², Sina Azadnajafabadi  ^{3,4}, Amir Hossein Norooznezhad⁵, Fahimeh Ghotbzadeh Vahdani², Zahra Panahi², Nafiseh Hivechi², Arefeh Eshghinejad², Saina Nassiri², Nasim Eshraghi², Arezoo Behzadian², Fatemeh Asadi², Zeinab Mansouri², Shahrazad Hadavand⁶, Rosa Bahari², Elham Sadat Banihashemi⁶, Seyed Jafar Razavi⁷, Soroush Faramarzi², Mohadese Dashtkoohi  ² and Sedigheh Hantoushzadeh  ^{2,*}

¹Department of Infectious Diseases, Imam Khomeini Complex, Tehran University of Medical Sciences, Tehran, Iran

²Vali-E-Asr Reproductive Health Research Center, Family Health Research Institute, Tehran University of Medical Sciences, Tehran, Iran

³Non-Communicable Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran

⁴Department of Surgery, Tehran University of Medical Sciences, Tehran, Iran

⁵Medical Biology Research Center, Health Technology Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁶Obstetricians and gynecologists, Department of Obstetrics and Gynecology, Faculty of Medicine, Shahed University, Tehran, Iran

⁷Telemedicine Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

*Corresponding author: Vali-E-Asr Reproductive Health Research Center, Family Health Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Email: hantoushzadeh@tums.ac.ir

Received 2023 October 31; Accepted 2023 November 09.

Abstract

Background: Corticosteroid administration may impair blood sugar control and cause other adverse effects in pregnant women with moderate to severe COVID-19. To our knowledge, there have been no studies on the effect of corticosteroid therapy on pregnant women with COVID-19 infection in terms of hyperglycemia or insulin needs.

Objectives: The purpose of this study was to compare 2 different regimens of corticosteroid therapy, specifically dexamethasone, and methylprednisolone, in terms of their impact on newly diagnosed hyperglycemia in pregnant women infected with COVID-19, as well as the duration of this condition.

Methods: The current cohort study was conducted from August to November 2021 on hospitalized pregnant women with severe COVID-19. They received either the first protocol, which involved daily administration of 2 mg/kg of methylprednisolone intravenously, or the second protocol, which included daily intravenous administration of 6 mg of dexamethasone. The study aimed to compare the incidence and duration of hyperglycemia until delivery between these 2 groups.

Results: A total of 59 participants were recruited after meeting the inclusion criteria. There were no significant differences in the demographic and clinical characteristics of patients between the 2 groups. Among the 59 pregnant women included in the study, 24 (40.7%) developed hyperglycemia that required insulin therapy. The incidence of hyperglycemia did not significantly differ between the 2 treatment regimens (P-value = 0.069). In the follow-up period, hyperglycemia resolved in both groups within up to 4 weeks after treatment, and they no longer required insulin.

Conclusions: Corticosteroid-treated pregnant patients are associated with transient hyperglycemia, the incidence of which is unrelated to the type of corticosteroid used. Therefore, corticosteroid administration should be initiated with caution and monitored appropriately in pregnant women with COVID-19 infection.

Keywords: COVID-19, Corticosteroids, Pregnancy, Dexamethasone, Methylprednisolone, Gestational Diabetes Mellitus, Transient Hyperglycemia

1. Background

The coronavirus disease 2019 (COVID-19) pandemic has had a significant impact on maternal health, leading to clinical and psychological complications and changes in healthcare provision and accessibility to healthcare services (1). Pregnant women with COVID-19 infection

experience higher rates of hospitalization and admission to the intensive care unit (ICU) (2). This could be attributed to pregnancy-induced alterations in the immune system and the body's response to viral infections during fetal development (3).

In non-pregnant patients with COVID-19 infection,

a massive release of pro-inflammatory cytokines and chemokines can lead to a cytokine storm, triggering severe inflammatory responses and causing tissue/organ damage or even death (4). Consequently, a short course of glucocorticoids has been shown to improve clinical outcomes in patients with severe COVID-19 (5). Additionally, there is evidence that COVID-19 infection significantly affects the cytokine profile in pregnant women, with cytokine levels correlating with disease severity and varying across trimesters (6).

Although COVID-19 primarily impacts the respiratory system, there have been reports of metabolic disturbances, hyperglycemia, and new-onset diabetes shortly after the pandemic's outbreak (7, 8). One possible explanation is that the SARS-CoV-2 virus may affect pancreatic β -cells, reducing insulin secretion. Furthermore, the significant production of cytokines can induce insulin resistance. Both reduced insulin secretion and increased insulin resistance can contribute to hyperglycemia, regardless of previous diabetes status (9, 10).

Pregnancy is associated with relative glucose intolerance and insulin resistance. Therefore, the combination of pregnancy and corticosteroids may lead to impaired glycemic control in both diabetics and non-diabetics. The duration and severity of hyperglycemia were significantly associated with the initial glucose tolerance test before corticosteroid therapy and other maternal risk factors for gestational diabetes mellitus (GDM) (11).

Long-term use of corticosteroids during pregnancy is linked to adverse effects for both the fetus and the mother, including hyperglycemia (12), preeclampsia, preterm premature rupture of membranes, pyelonephritis, and thromboembolisms. Neonates born to mothers who have received corticosteroid therapy during pregnancy are more susceptible to hypoglycemia (13). Additionally, short-term antenatal corticosteroid therapy is commonly administered to women at risk of preterm delivery to promote fetal pulmonary maturation. Previous studies have recommended monitoring glycemic levels for at least 5 days after initiating corticosteroid therapy to detect any potential hyperglycemia (14, 15).

As a result, corticosteroids may lead to impairments in blood sugar control and other adverse effects in pregnant women with moderate to severe COVID-19. Several studies have investigated corticosteroid therapy in pregnant women with COVID-19 disease (16, 17). However, we did not find any studies specifically addressing the impact of corticosteroid therapy on the incidence of hyperglycemia and insulin requirements in pregnant women with COVID-19 infection.

2. Objectives

The objective of this study was to compare 2 different corticosteroid therapies, namely systemic dexamethasone, and methylprednisolone, in terms of their effects on glycemic status during pregnancy and the duration of hyperglycemia until the postpartum period.

3. Methods

3.1. Setting

This cohort study was conducted from August 2021 to November 2021 at an academic center. The inclusion criteria comprised pregnant women with COVID-19 (singleton pregnancies) confirmed by real-time polymerase chain reaction (PCR) who required hospitalization. Admission criteria were defined as moderate to severe illness based on the National Institutes of Health (NIH) criteria, including $\text{SpO}_2 < 94\%$ on room air at sea level, a ratio of arterial partial pressure of oxygen to fraction of inspired oxygen ($\text{PaO}_2/\text{FiO}_2$) < 300 mm Hg, a respiratory rate > 30 breaths/min, or lung infiltrates $> 50\%$ (18).

Exclusion criteria encompassed gestational or overt diabetes mellitus, hypertension, prior corticosteroid use for any reason, contraindications to steroid therapy, immunodeficiency disorders, and unwillingness to participate in the study. Additionally, the diagnostic criteria for GDM included at least 1 abnormal value (≥ 92 , 180, and 153 mg/dL for fasting, 1-hour, and 2-hour plasma glucose concentration, respectively) following a 75 g oral glucose tolerance test (OGTT) (19), which was conducted for all participants.

3.2. Study Design

The admitted patients received treatment and monitoring from a team of healthcare providers, including infectious disease specialists and obstetricians. Each participant was treated according to 1 of the 2 hospital protocols. Protocol 1 involved administering 2 mg/kg of methylprednisolone intravenously (IV) daily (infused over 60 minutes), which was tapered to half the dosage every 5 days for a total of 10 days. Protocol 2 consisted of 6 mg of dexamethasone intravenously daily for 10 days. Both protocols included the administration of remdesivir, with 200 mg IV on the first day followed by 100 mg IV daily for 5 days, and anticoagulation therapy with enoxaparin or heparin, which was common to both protocols.

Fasting and postprandial plasma glucose levels at 2 hours after each meal were evaluated 24 hours after

corticosteroid therapy. Fasting sugar levels above 92 and postprandial levels above 120 were considered indicative of hyperglycemia. If 30% of the glucose levels were above normal within 48 hours, insulin treatment and a diabetic diet were initiated for the patients. Blood glucose levels were monitored until hyperglycemia was under control. Fetal monitoring was conducted using the fetal biophysical profile (BPP) twice weekly, and fetal growth was assessed by ultrasound every 3 weeks as long as hyperglycemia persisted. After discharge, glucose control was continued, and insulin administration was adjusted if necessary until normal glucose levels were achieved.

Demographic features, as well as baseline clinical and laboratory characteristics, oxygen saturation, type of oxygen supplementation, and respiratory rate, were evaluated in all patients. All participants were followed up until 6 weeks after delivery, and their glycemic status, gestational age at delivery, mode of delivery, and duration of insulin requirement were assessed and compared between the 2 groups.

3.3. Statistical Analysis

All analyses were performed using SPSS v. 26.0 (IBM Corp., Armonk, NY, USA). Numerical data were described as mean \pm standard deviation (SD), and categorical data were presented as numbers (%). Statistical differences were assessed using an independent *t*-test or Mann-Whitney U test as appropriate. The chi-square test or Fisher's exact test was used for comparing categorical variables. The paired-sample *t*-test was employed to evaluate changes in clinical indices before and after the intervention, with a *P*-value < 0.05 considered statistically significant.

3.4. Ethical Considerations

The study received approval from the ethics committee of Tehran University of Medical Sciences with an institutional review board (IRB) number IR.TUMS.IKHC.REC.1400.351. Written informed consent was obtained from all patients or their legally authorized representatives after providing an explanation of the study's objectives and methods to the best of their understanding.

4. Results

A total of 75 pregnant women were assessed, of which 59 met the inclusion criteria. Among these, 26 (44.1%) patients received methylprednisolone, while 33 (55.9%) patients received dexamethasone. As shown in Table 1, there were no significant differences (*P* > 0.05) in

demographic and clinical characteristics between the 2 groups at the time of admission.

The laboratory indices at admission for each group are displayed in Table 2. There were no statistically significant differences in laboratory indices between the groups (*P*-value > 0.05), except for D-dimer, which was significantly higher in the methylprednisolone group (*P*-value = 0.033). However, it is important to note that there was no evidence of thromboembolism reported in the patients.

In total, 24 (40.7%) cases developed hyperglycemia during hospitalization, requiring insulin control. The incidence of increased glucose levels in the methylprednisolone and dexamethasone groups was 8 (30.8%) and 16 (48.5%), respectively, showing no significant difference (*P*-value = 0.169). A comparison of patient characteristics between the hyperglycemic and normal groups before starting treatment is presented in Table 3. Additionally, a comparison of the patients' laboratory indices between the hyperglycemic and normal groups is shown in Table 4. As observed, there were no differences in patient characteristics and laboratory indices, except for higher fasting blood glucose (*P*-value < 0.000) in the hyperglycemic group.

During follow-up telephone calls, 47 out of 59 (79.6%) of the patients responded. Preterm delivery (*P*-value = 1.00) and Cesarean section (*P*-value = 0.617) in patients with and without hyperglycemia did not show significant differences. This result was consistent across both treatment groups, including methylprednisolone (*P*-value = 0.638) and dexamethasone (*P*-value = 0.404). In the hyperglycemia group, the duration of insulin requirement lasted up to 4 weeks and did not differ significantly between the 2 treatment groups (Table 5).

5. Discussion

This study revealed that the incidence of increased blood glucose after treatment with corticosteroids in pregnant women with moderate to severe COVID-19 infection was 40.7%, requiring insulin control. Although serum glucose levels returned to normal within 4 weeks, no significant difference was observed between the 2 corticosteroid therapy regimens.

COVID-19 has been shown to potentially disrupt metabolism in the long term, leading to increased blood sugar levels and the onset of diabetes. This effect may be attributed to the virus's binding to angiotensin-converting enzyme 2 (ACE2), which is present in pancreatic beta cells and adipose tissue. It is likely that SARS-CoV-2 can directly impair glucose metabolism (20). Additionally, the inflammatory response triggered by the infection can

Table 1. Comparison of the Patients' Characteristics in Each Group^{a, b}

Characteristics (Mean \pm SD)	Methylprednisolone	Dexamethasone	Total	P-Value ^c
Age (y)	29.1 \pm 6.2	31.1 \pm 5.6	30.2 \pm 5.9	0.259
Gestational age (week)	29.7 \pm 7.6	29.7 \pm 5.6	29.7 \pm 6.5	0.586
Gravida	2.3 \pm 1.2	1.9 \pm 1.2	2.1 \pm 1.2	0.385
Body mass index (kg/m ²)	26.4 \pm 3.0	26.9 \pm 3.6	26.7 \pm 3.4	0.420
Ventilation status; No. (%)				0.105
O ₂ mask	2 (50.0)	10 (71.4)	12 (20.3)	
Nasal O ₂ cannula	0 (0)	3 (21.4)	3 (5.1)	
Reserve mask	2 (50.0)	1 (7.1)	3 (5.1)	
Pulmonary involvement in CT scan (%)	29.8 \pm 13.1	31.7 \pm 15.8	30.9 \pm 14.6	0.672

^a Data are presented as mean \pm SD unless otherwise indicated.^b Mean \pm SD compared by Mann-Whitney U test. No. (%) compared by the chi-square test.^c Significant level was considered P < 0.05.**Table 2.** Comparison of the Laboratory Indices of the Participants in Each Group at Admission^a

Characteristics	Methylprednisolone	Dexamethasone	Total	P-Value ^b
FBS	104.2 \pm 11.6	102.6 \pm 15.4	103.3 \pm 13.8	0.340
WBC	8.4 \pm 2.8	7.3 \pm 2.5	7.7 \pm 2.7	0.240
Neutrophil	6.6 \pm 2.1	5.7 \pm 2	6.1 \pm 2.1	0.151
Lymphocyte	1.8 \pm 0.9	1.6 \pm 0.7	1.7 \pm 0.8	0.783
Lymphocyte/Neutrophil ratio	0.27 \pm 0.1	0.29 \pm 0.1	0.28 \pm 0.1	0.436
Hemoglobin	11.6 \pm 1.2	12 \pm 2	11.9 \pm 1.7	0.448
Platelet	198.1 \pm 63.7	188.4 \pm 56.5	192.7 \pm 59.5	0.531
ESR	85.9 \pm 40.9	86.2 \pm 31.1	86.1 \pm 35.4	0.691
CRP	73.7 \pm 36.3	75.5 \pm 23.2	74.7 \pm 29.5	0.502
Creatinine	0.7 \pm 0.1	0.6 \pm 0.1	0.6 \pm 0.1	0.220
AST	33.8 \pm 14.4	32 \pm 9.8	32.8 \pm 12	0.915
ALT	34 \pm 12.9	32.2 \pm 9.8	33 \pm 11.2	0.903
LDH	38 \pm 12.5	36.7 \pm 10.3	37.3 \pm 11.3	0.888
Ferritin	153.2 \pm 181.1	109.6 \pm 76.4	128.3 \pm 131.7	0.492
D-dimer	1352 \pm 2512.3	75 \pm 362.5	591.9 \pm 1715.3	0.033

Abbreviations: FBS, fasting blood sugar (glucose); WBC, white blood cell; ESR, erythrocyte sedimentation rate; CRP, C-reactive protein; AST, aspartate aminotransferase; ALT, alanine transaminase; LDH, lactate dehydrogenase.

^a Data are presented as mean \pm SD. Mean \pm SD compared by the Mann-Whitney U test.^b Significant level was considered P < 0.05.

contribute to insulin resistance and dysfunction of beta cells (21).

In non-pregnant cases with severe COVID-19 infection, the administration of systemic corticosteroids has been associated with lower mortality (4). This is achieved by ameliorating lung inflammation and preventing acute respiratory distress syndrome (22), as well as reducing the duration of mechanical ventilation (23). However, the use of corticosteroids in treating acute COVID-19 can

exacerbate insulin resistance. Additionally, SARS-CoV-2 may worsen the existing pro-inflammatory state seen in type 2 diabetes. Glycemic abnormalities have been observed in hospitalized COVID-19 patients for up to 2 months after their initial diagnosis (24). These findings emphasize the importance of continuous monitoring and management of blood sugar levels in individuals who have recovered from COVID-19.

While corticosteroids are not entirely safe during

Table 3. Comparison of the Patients' Characteristics Between Hyperglycemic and Normal Groups at Admission^a

Characteristics	Glucose Intolerance	Normal	P-Value ^b
Age (y)	30.5 ± 6.5	30.0 ± 5.5	0.610
Gestational age (week)	30.4 ± 3.9	29.2 ± 7.8	0.770
Gravida	2.1 ± 1.4	2.1 ± 1.1	0.790
Body mass index (kg/m ²)	26.5 ± 2.9	26.8 ± 3.6	0.798
O ₂ saturation	94.4 ± 2.1	94.7 ± 2.2	0.651
Ventilation status; No. (%)			0.504
O ₂ mask	3 (50.0)	9 (75.0)	
Nasal O ₂ cannula	1 (16.7)	2 (16.7)	
Reserve mask	2 (33.3)	1 (8.3)	
Pulmonary involvement in CT scan	34.0 ± 16.7	28.7 ± 12.8	0.296

Abbreviation: CT, computed tomography.

^a Data are presented as mean ± SD. Mean ± SD compared by the Mann-Whitney U test.^b Significant level was considered P < 0.05.**Table 4.** Comparison of the Patients' Laboratory Indices Between Glucose Intolerance and Normal Groups at Admission

Characteristics	Glucose Intolerance	Normal	P-Value ^b
FBS	111.6 ± 13.5	97.3 ± 10.7	< 0.000
2-hour postprandial glucose (morning)	110.4 ± 7.6	110.3 ± 8.3	0.614
2-hour postprandial glucose (midday)	120.0 ± 10.3	109.5 ± 7.6	0.705
2-hour postprandial glucose (afternoon)	124.5 ± 27.2	116.0 ± 9.1	0.614
WBC	7.1 ± 1.9	8.2 ± 3	0.151
Neutrophil	5.6 ± 1.6	6.4 ± 2.3	0.123
Lymphocyte	1.5 ± 0.6	1.7 ± 0.9	0.758
Hemoglobin	11.5 ± 1.3	12.1 ± 1.9	0.215
Platelet	207.6 ± 72.8	182.5 ± 46.7	0.547
ESR	90.2 ± 24.8	83.2 ± 41.3	0.090
CRP	77.6 ± 24.6	72.7 ± 32.6	0.294
Creatinine	0.6 ± 0.1	0.6 ± 0.1	0.212
AST	33 ± 8.7	32.7 ± 13.9	0.371
ALT	33.7 ± 9.7	32.5 ± 12.2	0.371
LDH	38.3 ± 11.7	36.6 ± 11.1	0.613
Ferritin	176.3 ± 188.7	95.7 ± 56.4	0.124
D-dimer	159 ± 476.5	916.6 ± 2195.6	0.247

Abbreviations: FBS, fasting blood glucose; WBC, white blood cell; ESR, erythrocyte sedimentation rate; CRP, C-reactive protein; AST, aspartate aminotransferase; ALT, alanine transaminase; LDH, lactate dehydrogenase.

^a Data are presented as mean ± SD. Mean ± SD compared by the Mann-Whitney U test.^b Significant level was considered P < 0.05.

pregnancy, they are recommended for pregnant women with moderate-to-severe COVID-19. In cases where oral prednisolone or intravenous hydrocortisone is not an option, methylprednisolone can be considered (25). The World Health Organization (WHO) makes a similar recommendation, though it does not explicitly specify

which corticosteroid should be administered to women with moderate-to-severe COVID-19 (26).

Pregnancy is known to induce significant acquired insulin resistance, posing a short-term challenge to β -cells, with GDM developing in women whose β -cells are unable to cope (27). Previous research

Table 5. Duration of Insulin Requirement in and after Pregnancy in the Glucose Intolerance Group (n = 15)^a

Insulin Requirement	Methylprednisolone	Dexamethasone	P-Value
During hospitalization	1 (16.6)	4 (44.4)	0.611
1 week after discharge	2 (33.3)	2 (22.2)	0.847
10 days after discharge	0	1 (11.1)	0.409
2 weeks after discharge	0	1 (11.1)	0.409
4 weeks after discharge	3 (50.0)	1 (11.1)	0.633

^a Data are presented as No. (%). No. (%) compared by the chi-square test or Fisher's exact test.

has suggested that the use of corticosteroids, such as betamethasone and dexamethasone, during pregnancy can lead to maternal hyperglycemia. This occurs because corticosteroids antagonize insulin synthesis and increase gluconeogenesis, resulting in elevated blood glucose levels. This effect is particularly concerning for pregnant women with diabetes, as it can further disrupt glucose homeostasis (11).

Although the placenta plays a role in protecting the fetus from the side effects of corticosteroids by extensively metabolizing them into inactive products (28), repeated use of corticosteroids can lead to reduced fetal body movements and breathing, potential intrauterine growth restriction, low birth weight, fetal hypoglycemia, and an increased risk of early-onset neonatal sepsis. Impaired glucose tolerance and reduced insulin sensitivity induced by corticosteroid therapy in a mother, even if transient, might be enough to affect fetal glucose hemostasis (29, 30). Overall, these findings underscore the importance of managing and monitoring blood glucose levels in pregnant women receiving corticosteroids to minimize disruptions in glucose metabolism and potential adverse effects on both maternal and neonatal health (14).

Collectively, these findings suggest that pregnant women with COVID-19 who also receive corticosteroids are at a higher risk of developing glucose intolerance. Although there is no available data on the number of people developing new hyperglycemia after COVID-19 and corticosteroid therapy, COVID-19 can induce new-onset diabetes in high-risk and susceptible populations and can worsen blood sugar control in those with preexisting diabetes (31). This aligns with our results, where the mean fasting blood sugar levels upon hospitalization were higher than 100 mg/dL, considered hyperglycemia during pregnancy, in women with COVID-19.

In a recent study comparing methylprednisolone and dexamethasone for the treatment of COVID-19 in non-pregnant individuals, hyperglycemia was more frequent in the methylprednisolone group, but it was managed without complications (32). Although this study

did not report impaired glucose tolerance rates, our study found that approximately 40% of individuals developed glucose intolerance after receiving corticosteroids. However, all of them returned to normal glucose values within 4 weeks. Additionally, the rate of increased blood glucose was similar between the methylprednisolone and dexamethasone treatment groups.

The follow-up results for patients were also promising. The rates of preterm delivery and cesarean section, which are considered complications of COVID-19, were low and did not differ between treatment groups or patients with or without hyperglycemia.

This is the first study to evaluate the prevalence of transient hyperglycemia in pregnant women after corticosteroid therapy for COVID-19 infection. However, the current study had several limitations, including a small sample size in each group and limited follow-up data. Further studies with larger sample sizes and longer follow-up periods are required to assess the potential harmful and beneficial effects of different corticosteroid types in pregnant women with COVID-19 and their neonates.

Acknowledgments

We thank all the staff of the Perinatology Ward of Vali-e Asr Hospital for their sincere cooperation.

Footnotes

Authors' Contribution: M.S, conception and design of the study, revising the manuscript; M.G, conception and design of the study, drafting and revising the manuscript; S.M, data collection, analysis, and interpretation of data, drafting the manuscript; S.A, analysis and interpretation of data, critical revision of the manuscript; A.H.N, analysis and interpretation of data, critical revision of the manuscript; F.G.H, data collection, critical revision of the manuscript; Z.P, data collection, analysis, and interpretation of data, critical revision of the manuscript;

N.H, interpretation of data, critical revision of the manuscript; A.E, analysis and interpretation of data; S.N, data collection; N.E, data collection, analysis, and interpretation of data; A.B, data collection; F.A, statistical analysis; Z.M, critical revision of the manuscript for important intellectual content; S.H, contribution to the design of the study and critical revision of the manuscript; E.S.B, contribution to the design of the study, critical revision of the manuscript; S.J.R, contribution to the design of the study; S.H, conception and design of the study, analysis and interpretation of data, critical revision of the manuscript, and study supervision. All the authors have read and approved the final version of the manuscript.

Conflict of Interests: We declare that second author is a member of the editorial board of this journal. However, her involvement in the editorial board does not pose any conflict of interest with the content of this manuscript.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The study was approved by the Ethics Committee of the Tehran University of Medical Sciences with the IRB number of [IR.TUMS.IKHC.REC.1400.351](#).

Funding/Support: The authors have not received any financial support or research grants that are relevant to the subject matter of this manuscript.

Informed Consent: Written informed consent was obtained from all the patients or their legally authorized representatives after an explanation of the aims and methods of the study according to their knowledge.

References

- Kotlar B, Gerson EM, Petrillo S, Langer A, Tiemeier H. The impact of the COVID-19 pandemic on maternal and perinatal health: A scoping review. *Reprod Health*. 2021;18(1):10. [PubMed ID: [33461593](#)]. [PubMed Central ID: [PMC7812564](#)]. <https://doi.org/10.1186/s12978-021-01070-6>.
- Ellington S, Strid P, Tong VT, Woodworth K, Galang RR, Zambrano LD, et al. Characteristics of Women of Reproductive Age with Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status - United States, January 22-June 7, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(25):769-75. [PubMed ID: [32584795](#)]. [PubMed Central ID: [PMC7316319](#)]. <https://doi.org/10.15585/mmwr.mm6925a1>.
- Moore KM, Suthar MS. Comprehensive analysis of COVID-19 during pregnancy. *Biochem Biophys Res Commun*. 2021;538:180-6. [PubMed ID: [33384142](#)]. [PubMed Central ID: [PMC7759124](#)]. <https://doi.org/10.1016/j.bbrc.2020.12.064>.
- W. H. O. Rapid Evidence Appraisal for COVID-19 Therapies Working Group, Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, et al. Association between administration of systemic corticosteroids and mortality among critically ill Patients With COVID-19: A Meta-analysis. *JAMA*. 2020;324(13):1330-41. [PubMed ID: [32876694](#)]. [PubMed Central ID: [PMC7489434](#)]. <https://doi.org/10.1001/jama.2020.17023>.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. [PubMed ID: [31986264](#)]. [PubMed Central ID: [PMC7159299](#)]. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- Tanacan A, Yazihan N, Erol SA, Anuk AT, Yucel Yetiskin FD, Biriken D, et al. The impact of COVID-19 infection on the cytokine profile of pregnant women: A prospective case-control study. *Cytokine*. 2021;140:155431. [PubMed ID: [33503581](#)]. [PubMed Central ID: [PMC7810028](#)]. <https://doi.org/10.1016/j.cyto.2021.155431>.
- Blanke CD. In response: Diabetes is a risk factor for the progression and prognosis of COVID-19. *Diabetes Metab Res Rev*. 2020;36(7):e3331. [PubMed ID: [32452598](#)]. [PubMed Central ID: [PMC7267111](#)]. <https://doi.org/10.1002/dmrr.3331>.
- Chen M, Zhu B, Chen D, Hu X, Xu X, Shen WJ, et al. COVID-19 may increase the risk of insulin resistance in adult patients without diabetes: A 6-month prospective study. *Endocr Pract*. 2021;27(8):834-41. [PubMed ID: [33887468](#)]. [PubMed Central ID: [PMC8054613](#)]. <https://doi.org/10.1016/j.eprac.2021.04.004>.
- Khunti K, Del Prato S, Mathieu C, Kahn SE, Gabbay RA, Buse JB. COVID-19, Hyperglycemia, and New-Onset Diabetes. *Diabetes Care*. 2021;44(12):2645-55. [PubMed ID: [34625431](#)]. [PubMed Central ID: [PMC8669536](#)]. <https://doi.org/10.2337/dc21-1318>.
- Ceriello A. Hyperglycemia and COVID-19: What was known and what is really new? *Diabetes Res Clin Pract*. 2020;167:108383. [PubMed ID: [32853690](#)]. [PubMed Central ID: [PMC7445137](#)]. <https://doi.org/10.1016/j.diabres.2020.108383>.
- Jolley JA, Rajan PV, Petersen R, Fong A, Wing DA. Effect of antenatal betamethasone on blood glucose levels in women with and without diabetes. *Diabetes Res Clin Pract*. 2016;118:98-104. [PubMed ID: [27351800](#)]. <https://doi.org/10.1016/j.diabres.2016.06.005>.
- Mariotti V, Marconi AM, Pardi G. Undesired effects of steroids during pregnancy. *J Maternal-Fetal & Neonatal Medicine*. 2009;16(2):5-7. <https://doi.org/10.1080/jmfm.16.2.5.7>.
- Uquillas KR, Lee RH, Sardesai S, Chen E, Ihenacho U, Cortessis VK, et al. Neonatal hypoglycemia after initiation of late preterm antenatal corticosteroids. *J Perinatol*. 2020;40(9):1339-48. [PubMed ID: [32060360](#)]. <https://doi.org/10.1038/s41372-020-0589-1>.
- Kalra S, Kalra B, Gupta Y. Glycemic management after antenatal corticosteroid therapy. *N Am J Med Sci*. 2014;6(2):71-6. [PubMed ID: [24696828](#)]. [PubMed Central ID: [PMC3968568](#)]. <https://doi.org/10.4103/1947-2714.127744>.
- Kemp MW, Newnham JP, Challis JG, Jobe AH, Stock SJ. The clinical use of corticosteroids in pregnancy. *Hum Reprod Update*. 2016;22(2):240-59. [PubMed ID: [26590298](#)]. <https://doi.org/10.1093/humupd/dmv047>.
- Magala Ssekandi A, Sserwanja Q, Olal E, Kawuki J, Bashir Adam M. Corticosteroids Use in Pregnant Women with COVID-19: Recommendations from Available Evidence. *J Multidiscip Healthc*. 2021;14:659-63. [PubMed ID: [33758509](#)]. [PubMed Central ID: [PMC798138](#)]. <https://doi.org/10.2147/JMDHS301255>.
- McIntosh JJ. Corticosteroid Guidance for Pregnancy during COVID-19 Pandemic. *Am J Perinatol*. 2020;37(8):809-12. [PubMed ID: [32274772](#)]. [PubMed Central ID: [PMC7356057](#)]. <https://doi.org/10.1055/s-0040-1709684>.
- National Institutes of Health. Clinical Spectrum of SARS-CoV-2 Infection. *COVID-19 Treatment Guidelines* 2021. 2021.
- Karagiannis T, Bekiari EMKPK, Manolopoulos K, Paletas K, Tsapas A. Gestational diabetes mellitus: Why screen and how to diagnose. *Hippokratia*. 2010;14(3):151.
- Mendes TB, Camara-de-Souza AB, Halpern B. Hospital management of hyperglycemia in the context of COVID-19: Evidence-based clinical considerations. *Diabetol Metab Syndr*. 2022;14(1):37. [PubMed ID: [35246230](#)]. [PubMed Central ID: [PMC8895065](#)]. <https://doi.org/10.1186/s13098-022-00808-x>.
- Gojda J, Koudelkova K, Ouradova A, Lang A, Krbcova M, Gvozdeva A, et al. Severe COVID-19 associated hyperglycemia is caused by beta cell

dysfunction: A prospective cohort study. *Nutr Diabetes*. 2023;13(1):11. [PubMed ID: 37460458]. [PubMed Central ID: PMC10352285]. <https://doi.org/10.1038/s41387-023-00241-7>.

22. Kakoulidis I, Ilias I, Koukkou E. SARS-CoV-2 infection and glucose homeostasis in pregnancy. What about antenatal corticosteroids? *Diabetes Metab Syndr*. 2020;14(4):519-20. [PubMed ID: 32388332]. [PubMed Central ID: PMC7202835]. <https://doi.org/10.1016/j.dsx.2020.04.045>.

23. Villar J, Ferrando C, Martinez D, Ambros A, Munoz T, Soler JA, et al. Dexamethasone treatment for the acute respiratory distress syndrome: A multicentre, randomised controlled trial. *Lancet Respir Med*. 2020;8(3):267-76. [PubMed ID: 32043986]. [https://doi.org/10.1016/S2213-2600\(19\)30417-5](https://doi.org/10.1016/S2213-2600(19)30417-5).

24. Montefusco L, Ben Nasr M, D'Addio F, Loretelli C, Rossi A, Pastore I, et al. Acute and long-term disruption of glycometabolic control after SARS-CoV-2 infection. *Nat Metab*. 2021;3(6):774-85. [PubMed ID: 34035524]. [PubMed Central ID: PMC9931026]. <https://doi.org/10.1038/s42255-021-00407-6>.

25. Saad AF, Chappell L, Saade GR, Pacheco LD. Corticosteroids in the management of pregnant patients with coronavirus disease (COVID-19). *Obstet Gynecol*. 2020;136(4):823-6. [PubMed ID: 32769659]. <https://doi.org/10.1097/AOG.0000000000004103>.

26. World Health Organization. *Corticosteroids for COVID-19*. 2020. Available from: <https://www.who.int/publications/item/WHO-2019-nCoV-Corticosteroids-2020.1>.

27. Retnakaran R, Qi Y, Sermer M, Connelly PW, Hanley AJ, Zinman B. Glucose intolerance in pregnancy and future risk of pre-diabetes or diabetes. *Diabetes Care*. 2008;31(10):2026-31. [PubMed ID: 18628572]. [PubMed Central ID: PMC2551649]. <https://doi.org/10.2337/dc08-0972>.

28. Murphy VE, Fittcock RJ, Zarzycki PK, Delahunty MM, Smith R, Clifton VL. Metabolism of synthetic steroids by the human placenta. *Placenta*. 2007;28(1):39-46. [PubMed ID: 16549198]. <https://doi.org/10.1016/j.placenta.2005.12.010>.

29. Ogueh O, Johnson MR. The metabolic effect of antenatal corticosteroid therapy. *Hum Reprod Update*. 2000;6(2):169-76. [PubMed ID: 10782575]. <https://doi.org/10.1093/humupd/6.2.169>.

30. Felder L, Mehta V, Roman A, McCurdy R. Maternal hyperglycemia and neonatal hypoglycemia following betamethasone administration in non-diabetic women. *American J Obstetrics Gynecology*. 2022;226(1). <https://doi.org/10.1016/j.ajog.2021.11.493>.

31. Sosale A, Sosale B, Kesavadev J, Chawla M, Reddy S, Saboo B, et al. Steroid use during COVID-19 infection and hyperglycemia - What a physician should know. *Diabetes Metab Syndr*. 2021;15(4):102167. [PubMed ID: 34186344]. [PubMed Central ID: PMC8189750]. <https://doi.org/10.1016/j.dsx.2021.06.004>.

32. Ranjbar K, Moghadami M, Mirahmadizadeh A, Fallahi MJ, Khaloo V, Shahriarirad R, et al. Methylprednisolone or dexamethasone, which one is superior corticosteroid in the treatment of hospitalized COVID-19 patients: A triple-blinded randomized controlled trial. *BMC Infect Dis*. 2021;21(1):337. [PubMed ID: 33838657]. [PubMed Central ID: PMC8035859]. <https://doi.org/10.1186/s12879-021-06045-3>.