

The IFN- λ Genetic Polymorphism Association With the Viral Clearance Induced by Hepatitis C Virus Treatment in Pakistani Patients

Imran Tipu ^{1,2,*}; Fiona Marriage ^{2,3}; Zia-ur-Rahman Farooqi ^{2,4}; Hazel Platt ³; Muhammad Amin Athar ¹; Philip John Day ^{2,3}; Andrea Short ³

¹Institute of Biochemistry and Biotechnology, University of the Punjab, Lahore, Pakistan

²Manchester Institute of Biotechnology, University of Manchester, Manchester, UK

³Centre for Integrated Genomic Medical Research, University of Manchester, Manchester, UK

⁴National University of Science and Technology, Islamabad, Pakistan

*Corresponding Author: Imran Tipu, Institute of Biochemistry and Biotechnology, University of the Punjab, Lahore, Pakistan. Tel: +92-3214029804, E-mail: imran.tipu@gmail.com

Received: September 27, 2013; Revised: December 10, 2013; Accepted: January 9, 2014

Background: Polymorphisms in the interferon λ (INF λ) genes on chromosome 19 have been associated with clearance of hepatitis C virus (HCV) induced by interferon and ribavirin therapy however there is no such data available for Pakistani patients with HCV infection.

Objectives: In this study, the effects of single nucleotide polymorphisms (SNPs) have been investigated in response to treatment with interferon- α and ribavirin in a cohort of 75 HCV genotype 3a patients.

Patients and Methods: A total number of 50 SNPs from the Interferon λ region on chromosome 19 were genotyped to investigate allelic associations with the treatment response in HCV type 3a patients. Thirteen SNPs were associated with HCV clearance, with the most significant alleles being RS8109886 (Fisher's P = 0.0001), RS8113007 (Fisher's P = 0.0001) and RS12979860 (Fisher's P = 0.0002).

Results: These SNPs were found to be the most suitable SNPs for predicting treatment response in the present study. These findings support those reported previously. This could be used to improve HCV treatment strategies and suggest that Pakistani patients should be genotyped for the relevant SNPs to identify the patients who are more likely to respond to interferon and ribavirin therapy.

Conclusions: This therapy is costly and can be accompanied by several adverse side-effects, hence pre-treatment prediction of patients who are most likely to benefit would have both economic and patient benefits in the long term.

Keywords: Polymorphism, Genetic; Polymorphism, Single Nucleotide; Antiviral Agents; Interferons; Ribavirin; Hepacivirus

1. Background

Hepatitis C virus (HCV) infection is one of the leading causes of chronic liver disease and has emerged as a global concern of public health, affecting about 3% of the world's population. Pakistan is the sixth most populated country in the world and has a HCV prevalence rate of 5.9% (1). While there are different subtypes of HCV, genotype 3a is the most common form in patients from Pakistan, with frequency ranging from 28.6% (2) to 89% (3) depending on the province (4). The clinical outcome of HCV infection is determined by the interplay between viral, environmental and host related factors (5). The host's immune system is the most important factor in viral persistence and innate immunity is the first line of defense, intervening with interferons and natural killer cells (6). This immune response is influenced by genetic polymorphisms in cytokines, their receptors (7) and the polymorphic genetic makeup of human populations. Genetic variations and T-cell responses are responsible for the outcome of HCV treatment (8). The most common type of genetic varia-

tions are single nucleotide polymorphisms (SNPs) which occur approximately every 300 nucleotides in the human genome and can be used as biological markers for diseases or conditions. The majority of SNPs have no effect on health, but if SNPs are located within a gene or regulatory region, they can be functional in disease susceptibility and/or treatment response.

Studies have found that infected individuals with same HCV genotype differ in ability to spontaneously resolve infection, even if they have the same ethnic background with similar demographic features and are taking the same IFN- α /ribavirin therapy (9). The host genetics have been identified as key factors in the natural clearance of HCV and host SNPs have been already identified as associated factors in a number of studies in patients from different genetic backgrounds (Table 1). In particular, SNP RS12979860, present 3Kb upstream of the Interleukin 28B gene on chromosome 19, has been associated with a three-fold change in response to treatment against HCV infection in African-American and European cohorts (7). Another SNP from

Implication for health policy/practice/research/medical education:

The SNPs analyzed in this study showed significant association with response to the therapy which can be helpful for guiding the treatment of HCV patients in our population.

Copyright © 2014, Kowsar Corp; Published by Kowsar Corp. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

this region of chromosome 19, RS8099917, has been associated with HCV clearance in Australian (10) and Asian populations (11) (Table 1) and is located 8Kb upstream of the IFNL3 gene. In humans, four functional type III IFN λ (IFNL) genes are clustered around this region of chromosome 19 encoding cytokines IL29 (IFNL1), IL28A (IFNL2), IL28B (IFNL3) (12) and IFNL4 (13) and have a number of roles in controlling HCV infection including increasing the antiviral efficacy as a result of increased sub-saturating levels of IFN- α (14). IFN- λ binds to the heterodimeric receptors IFN- λ R1 and IL10R2 forming interferon stimulated genes (ISGs) complex and initiating a signal transduction cascade (15) leading to up-regulation of several ISGs with anti-viral effects (16). The IFN- λ receptors are present on the plasmacytoid dendritic cells, peripheral B cells, hepa-

tocytes and epithelial cells only so they can be used to target specific cell responses and can also help in avoiding adverse events of INF- α therapy (17). The role of SNPs present in the IFNL3 and IFNL4 genes in the spontaneous clearance of HCV was investigated, in addition to the associative role of SNPs present in the up- and down-stream regions of genes encoding IFN- λ . This data could be of value for predicting the response to interferon and ribavirin therapy in Pakistani patients and though would be of economic and patient benefit in the long term.

2. Objectives

In this study, the effects of SNPs have been investigated in response to treatment with interferon- α and ribavirin in a cohort of 75 patients with genotype 3a HCV.

Table 1. Previous Studies Which Have Reported SNP Allelic Associations

Author	Region	HCV genotype	RS12979860	RS8099917	RS12980275	RS4803219	RS8103142	RS8105790	RS10853728	RS7248668	RS4823221	RS1881222	RS4803217	RS28416813
Ge et al. (7)	USA	1	✓	✓					✓					
Suppiah et al. (10)	Australia			✓										
Tanaka et al. (11)	Japan	1		✓	✓									
Rauch et al. (18)	Switzerland	1, 4		✓										
Abe et al. (19)	Japan				✓									
Mangia et al. (20)	Italy	2, 3	✓											
McCarthy et al. (21)	USA			✓										
Thompson et al. (22)	USA			✓										
Bochud et al. (23)	Switzerland		✓	✓	✓									
Smith et al. (24)	Europe		✓	✓						✓	✓			
Yu.M.Lin et al. (25)	Taiwan			✓										
Chen et al. (26)	Taiwan		✓	✓	✓	✓			✓	✓	✓			✓
Scherzer et al. (27)	Austria		✓	✓										
Ridruejo et al. (28)	Argentine	1	✓	✓										
Yu et al. (25)	Taiwan	2					✓			✓				
Jun-qiang et al. (29)	China			✓										
Pedergnana et al. (30)	Egypt	4	✓					✓						
Shi et al. (31)	China		✓	✓	✓				✓					
de Castellarnau et al. (32)	Spain		✓	✓			✓	✓				✓	✓	
Grandi et al. (33)	Brazil	1	✓											
Prokunina-Olsson et al. (13)	USA			✓										
Stenkvist et al. (34)	Sweden			✓										
Gelinas et al. (35)	France			✓	✓									
Ezzikouri et al. (36)	Morocco			✓	✓									
Jung et al. (37)	Korea			✓	✓									

3. Patients and Methods

3.1. Selection and Description of Participants

Following ethical approval from the Institutional Review Board (University of Punjab, Pakistan) written informed consent for genetic testing including IFN- λ SNPs was obtained from each patient participating in the study. Patients were recruited from different areas of Punjab who visited National Genetics Laboratory, Lahore during March 2010 to May 2011. Patients displaying HCV like symptoms of infection ($n=150$) were screened for HCV RNA using an in-house PCR detection technique, of the 150 patients screened, 100 were positive for HCV RNA and 75 were classified as genotype 3a. Each patient was interviewed and a structured questionnaire was completed to figure out the demographic data.

3.2. Technical Information

3.2.1. HCV Detection

HCV viral RNA was extracted from the patient's serum using a QIAamp viral RNA extraction kit (Qiagen). The HCV RNA was detected in 100 individuals using sequence specific primers designed to target the highly conserved 5' UTR sequence in HCV (Table 2). The viral genotype was detected by nested PCR using unique antisense primers which amplify the 5' conserved sequence of HCV within the genotype and their poor homology with the sequence derived from other genotypes (Appendix 1). Only 75 patients identified with the HCV genotype 3a were selected for further study, this comprised 75% of the patients screened and thus the study avoided the effect of HCV genotypes on therapy response.

3.2.2. Treatment

All patients received three million IU of IFN- α three times a week subcutaneously and ribavirin (10 mg/day/kg body weight) for a total period of six months. Doses of IFN- α were adjusted according to platelet and white blood cell counts of patients. Ribavirin dose varied according to the haemo-

globin (Hb) levels and weight of individual patients. The therapy response was monitored by alanine aminotransferase (ALT) and HCV RNA levels at the beginning and end of treatment. The HCV RNA quantification was performed by the Artus HCV RT-PCR (Qiagen) kit using a Rotor-Gene 3000 (Corbett Robotics, Australia) instrument.

3.2.3. DNA Extraction

Human genomic DNA was extracted from peripheral blood mononuclear cells using a QIAamp blood DNA mini kit (Qiagen). DNA was quantified using a Nanodrop-ND1000 spectrophotometer (lab technologies) and concentrations were normalized to 15 ng/ μ L.

3.2.4. SNP Selection and Genotyping

In total, 50 SNPs were genotyped. Twenty five were from the coding region of the IL28B gene, 23 SNPs covered the 3' and 5' UTR's of all four IFN- λ genes and the remaining two SNPs were from the newly discovered IFNL-4 gene. The details of SNPs are given in supplementary data (Appendix 2 and 3). Genotyping was performed using the iPLEX assay on a SEQUENOM MassARRAY® platform. The primers were designed using the assay designing suite v1.0.1 (SEQUENOM) (Appendix 4). An initial PCR amplified a 50-60 bp region flanking the polymorphic site. The product was treated with 1 U/ μ L of shrimp alkaline phosphatase at 37°C for 40 minutes to dephosphorylate any unincorporated dNTPs. The iPLEX reaction product was desalted using a cationic resin, pre-treated with acidic reagents, for optimizing mass spectrophotometric analysis. The desalted iPLEX product was spotted on the SpectroCHIP using a Nano spotter (Sequenom) and loaded on to the mass spectrometer. Each spot was then subjected to a laser under vacuum by the matrix-assisted laser desorption ionization-time-of-flight (MALDI-TOF) method. Assays were designed to SNPs on chromosome 19q13.13 covering the region encoding the IFN- λ genes. After genotyping, SNPs and samples were quality checked.

Table 2. Significantly Associated SNPs ($P < 0.05$) With Sustained Virological Response to Interferon and Ribavirin Therapy^a

SNPs	MAF	Responder MAF ($n = 47$)	Non-Responder MAF ($n = 28$)	OR (95% CI)	P Value
RS8109886	0.41	0.32	0.44	3.6 (1.9-6.5)	0.0001
RS8113007	0.25	0.19	0.33	3.6 (1.9-6.5)	0.0001
RS12979860	0.3	0.23	0.41	3.1 (1.7-5.3)	0.0002
RS11665818	0.38	0.29	0.5	2.9 (1.6-5.3)	0.0003
RS955155	0.33	0.26	0.38	2.9 (1.6-5.1)	0.0004
RS688187	0.31	0.27	0.38	2.7 (1.5-4.7)	0.0011
RS4803217	0.3	0.25	0.38	2.7 (1.5-4.7)	0.0011
RS8105790	0.19	0.16	0.25	2.6 (1.4-4.6)	0.0022
RS4803221	0.22	0.16	0.27	2.6 (1.4-4.6)	0.0022
RS8099917	0.19	0.16	0.25	2.6 (1.4-4.6)	0.0022
RS7248668	0.19	0.16	0.25	2.6 (1.4-4.6)	0.0022
RS12972991	0.22	0.17	0.3	2.5 (1.4-4.5)	0.0024
RS11671087	0.41	0.32	0.5	2.2 (1.2-3.9)	0.0130

^a Abbreviations: CI, confidence interval; MAF, minor allele frequency; OR: odds ratio.

3.2.5. SNP Quality Controlled

SNPs were excluded from the analyses if the call rate < 90%, Minor Allele Frequencies < 0.05 and the cohort (responders + non-responders) was not in Hardy-Weinberg equilibrium (HWE, $P < 0.05$). Samples were excluded if the call rate was less than 90%. Call rate, Hardy-Weinberg equilibrium, minor allele frequencies, allelic and haplotypic associations and linkage disequilibrium (LD) were performed using BC|GENE version 3.5-087 software (Biocomputing Platforms, Sweden) whilst Microsoft Excel was used for the determination of means and averages.

3.3. Statistics

3.3.1. Association Analyses

Association of the genetic variants and spontaneous HCV clearance, was determined using logistic regression. The major alleles (as RS12979860 C) were compared with minor alleles (rs12979860 T) in statistical analyses to determine odds ratios (OR) and 95% confidence intervals (CI 95%).

3.3.2. Linkage Disequilibrium and Haplotype Analysis

Linkage disequilibrium between marker loci was assessed and haplotypic blocks were constructed using BC|GENE version 3.5-087 software (Biocomputing platforms, Sweden) and Haplovview 4.2 (<http://www.broadinstitute.org/haplovview/haplovview>).

3.3.3. Treatment Response

The effectiveness of IFN- λ loci SNPs was estimated for predicting the treatment response by comparing the sensitivity, specificity, positive predictive value (PPV) and

negative predictive value (NPV) for minor allele homozygotes. The most clinically useful parameter to investigate the treatment failure is PPV.

4. Results

4.1. Demographics

Out of 75 patients with genotype 3a HCV enrolled into the study, 46 were male and 29 were female. The virological response was monitored by quantification of HCV RNA at the beginning and at the end of the six months period of the therapy revealing that 63% of subjects (47) showed Sustained Virological Response (SVR) and 37% (28) patients were HCV RNA positive at the end of therapy. It also emerged that 75% of the patients were infected with HCV genotype 3a. These results were consistent with a recent review (4) showing the predominance of genotype 3a in the Pakistani population. The base line demographic, virological and clinical features of patients are shown in Table 3.

4.2. Sample and SNP Quality Control

We analyzed the region of ~ 62.4 kb (Chr 19, nucleotide positions, 39719200-39781600; build GRCh37.p10) containing 50 SNPs (Tables 2 and 4) present in the IFN- λ loci. Out of 50 SNPs, one failed the quality control (QC) criteria and was excluded from the analyses (SNP RS11881222 call rate = 80%); all other samples satisfied the inclusion criteria (> 90% call rate, HWE > 0.05). Twenty four SNPs present in the coding region of the IL28B gene were monomorphic in the studied Pakistani population and were therefore excluded from allelic association and haplotype analysis.

Table 3. Demographic and Clinical Characteristics of the Responders and Non-responders to Interferon and Ribavirin Therapy Against HCV Infection ^a

	Responders	Range	Non Responder	Range
Number of Patients, No. (%)	47 (63)		28 (37)	
Average Age, y	43	(21-60)	48	(28-63)
Gender				
Male	30		16	
Female	17		12	
Laboratory parameters				
Hb, g/dL	12.7	(8.2-16.4)	12.8	(7.1-17.1)
WBC, 10 \times 9/L	5.64	(2.8-11)	5.84	(3.3-9.4)
PLT, 10 \times 9/L	232	(93-402)	165	(67-287)
ALT, IU/L	63	(15-224)	93	(38-235)
HCV-RNA, KIU/mL, Initial	1200	(125-9900)	1034	(146-5000)
HCV-RNA, KIU/mL, End of treatment	below threshold	below threshold	2647	(120-9800)

^a Abbreviations: ALT, alanine transaminase; Hb, Haemoglobin; PLT: platelets; WBC: white blood cells.

4.3. Allelic Association

The allelic association revealed that a region of ~39 Kb (Chr 19, nucleotide positions, 39729450-39768250; build GRCh37. p10) containing 13 polymorphic SNPs in Pakistani population is strongly associated (Fisher's P value = 0.0003-0.0130) with spontaneous clearance and for all of these SNPs, spontaneous HCV clearance was more common with the major alleles. The most significant results were obtained with RS109886 (Odds ratio of presenting HCV clearance [OR] for C vs. A = 3.6 [95% CI: 1.9-6.5] Fisher's P = 0.0001), RS113007 (A vs. T OR = 3.6 [1.9-6.5]; Fisher's P = 0.0001) and RS12979860 (C vs. T OR = 3.1 [1.7-5.3]; Fisher's P = 0.0002). Among individuals, taking RS12979860 as an example, the proportion of HCV clearance was much higher in samples with major allele (80% SVR) as compared to minor T allele (34% SVR). The as-

sociation analysis of response to treatment by IFN- λ SNPs is described in Table 2.

4.4. Linkage Disequilibrium

Estimation of linkage disequilibrium was performed between 23 polymorphic IFN- λ region SNPs, which revealed three haplotypic blocks: haplotype block I, of eight Kb, included eight SNPs (RS35790907, RS12972991, RS12980275, RS12982533, RS8105790, RS688187, RS4803217 and RS12979860) in strong linkage disequilibrium ($r^2 \geq 0.85$) haplotype block II, of 4Kb included seven SNPs (RS4803221, RS1549928, RS10853727, RS109886, RS113007, RS8099917, RS7248668) in strong linkage disequilibrium ($r^2 \geq 0.95$) and block III contained just two SNPs (RS1671087 and RS11665818) lying approximately 6 kb apart from each other and in strong linkage disequilibrium ($r^2 \geq 0.85\%$)(Figure 1).


Table 4. Haplotypes With Odds Ratios ^{a, b}

Haplotype	Frequency, %	Responders, %	Non-responders, %	OR (95% CI)	P Value
AAATTGCCCATCATG	58.3	66.0	44.7	2.37 (1.34-4.20)	0.0028
TCGCCAATGATATGA	14.0	12.8	17.9	0.68 (0.31-1.48)	0.3286
AAATTGCCATAATA	9.60	7.40	14.0	0.46 (0.18-1.2)	0.106
TCGCTAATCGCATTG	8.00	4.20	12.5	0.28 (0.09-0.89)	0.022
TAGCCAATGGTATGA	5.30	3.20	7.10	0.41 (0.10-1.64)	0.194
AAATTGCTATAATA	2.00	3.20	1.90	1.52 (0.25-9.27)	0.650

^a The odds ratio has been calculated as carrying of haplotype vs. not carrying the haplotype.

^b The frequency of six haplotypes in responders and non-responders for a haplotype block covering 13 Kb IFN- λ . The SNP order is RS35790907, RS12972991, RS12980275, RS12982533, RS8105790, RS688187, RS4803217, RS10853727, RS109886, RS113007, RS8099917, RS7248668, RS6973285 RS1, RS805728 RS1, RS280602 RS1, RS4803224, RS10571087 RS1, RS165818 RS1 and RS7248931.

Figure 1. Analysis of Pairwise Linkage Disequilibrium (LD) Plot of IFN- λ Region



The linkage disequilibrium between the 17 SNPs in three LD blocks is shown. The red coloured squares represent $r^2 = 1.0$ and blue coloured squares represent $r^2 \leq 0.01$.

4.5. Haplotype Analysis

A total number of 6 haplotypes were investigated comprising of 15 SNPs using the Haplovview (MIT/Harvard/Brod Institute), among which haplotype one (AAATT-GCCCATCATG) comprising of major alleles of 14 SNPs (RS35790907, RS12972991, RS12980275, RS12982533, RS8105790, RS688187, RS4803217, RS12979860, RS4803221, RS1549928, RS10853727, RS8109886, RS8113007, RS8099917 and RS7248668) had most significant association (OR = 2.37, 95% CI = 1.34-4.20, P = 2.8x10-3) with therapy response in comparison with other detected haplotypes. The minor allele frequencies of each haplotype in responders and non-responders to the therapy with their odd ratios are shown in Table 1.

4.6. Treatment Response

The three highly associated SNPs with the treatment response; RS8109886 (PPV of 89%, 95 % CI = 81.17-94.37), RS8113007 (PPV of 74%, 95 % CI = 64.27-82.26) and RS12979860 (PPV of 74%, 95 % CI = 64.27-82.26) are also best indicators for predicting the treatment response. The sensitivity, specificity, prevalence, NPV and PPV of the IFN- λ loci SNPs has been shown in Supplementary Appendix 5.

5. Discussion

The treatment of patients with HCV is based on clinical, demographic and virological characteristics of the disease, which are helpful from a population perspective but these baseline parameters are not suitable for predicting the treatment response in HCV patients infected with the most common genotype, 3a. Two SNPs have been most frequently associated with viral clearance across all HCV genotypes in different populations of the world: RS8099917 and RS12979860 (Table 1) and efforts have been largely directed at determining which of them is most likely to be more suitable for establishing the most useful diagnostic test for predicting treatment. Genotype 3a is the most common genotype of HCV infections in Pakistan (4, 38). In a new cohort of 75 type 3a Pakistani patients SNPs in the up-and down-stream regions of IFN- λ and SNPs from IFNL3 and IFNL4 with known association to HCV clearance in other patient populations, were genotyped (Table 1). The allelic associations of four SNPs that have been reported previously in a number of populations were confirmed here (RS8105790, RS12979860, RS8099917 and RS7248668, Table 1) and a novel associations in Pakistani patients was identified (Table 2). The most significant SNPs (RS8109886 and RS8113007) detected by the present this in addition to six other SNPs have not been reported previously to have any association with HCV clearance in other populations and could be relevant to patients of Pakistani origin, although this requires follow-up studies to be fully confirmed.

Five SNPs reported in the literature were excluded from

this study (RS4803219, RS8103142, RS4823221, RS28416813 and RS11881222). SNP RS11881222 failed our QC and was excluded for a low call rate (< 80%) but the other five SNPs were not included because the Sequenom primer design software was unable to design suitable primers and probes for them. Excluding these SNPs from our study could represent missed associations in Pakistani patients and constitute additional analyses in this cohort and in future studies to determine whether they have any role in HCV clearance in Pakistani patients as well as the ones reported in patients from Taiwan, Spain, China and Europe (Table 1). None of the SNPs associated with HCV clearance in this study were in coding regions; but were located in regions up-or down-stream of genes or in the 3' or 5'UTR. This suggests that they have a regulatory function rather than directly affecting protein structure. The 13 SNPs associated with HCV clearance in this study formed 6 haplotypes, of which the major alleles of SNPs RS8109886, RS8113007, RS12979860 and RS8099917 were all present on haplotype I, the haplotype with the highest Odds Ratio for predicting the treatment response (Table 4). The role of these SNPs has been established as having effects on the binding of different transcription factors and alterations of methylation sites resulting in reduced expression of IL28B, and up-regulation of ISGs in the responder haplotypes in response to IFN- α stimulation therapy (24) while IL28B non-responders have high ISG expression in infected hepatocytes, and that high ISG levels independently predicts poor response to the therapy (39). HCV clearance is a complex process, dependent on the type of HCV infection and the host's immunity-related genetic factors. Some SNPs associated with HCV clearance in Pakistani patients are the same as those that have been detected to have associations in other cohorts too (Table 1) and suggest a common genetic background across multiple populations for HCV clearance. However, number of alleles identified in this study were unique to the present study which could suggest Pakistani-specific factors for HCV clearance, particularly for type 3a. It is important to consider, however, that this data comprised a small sample size and that repeating this study in a larger cohort could affect the findings and alter the outcome of some markers. For this reason, the data presented here should be interpreted with caution until it can be further verified. These findings, however, do support results widely reported from other populations where host genotype has been a proven factor in HCV clearance and treatment response (Table 1). Restricting this analysis to type 3a patients introduced a selection bias meaning if genotyping were to be introduced as a screening strategy, patients would require screening for HCV type prior to genotyping for treatment response. This selection strategy was chosen because type 3a is the most common form of HCV in Pakistan and so represents the largest treatment group. Confirming the association of these SNPs and HCV clearance, in other HCV types requires further investigation. Tailoring treatments to target potential responders,

as opposed to generalized, universal treatment strategies, will be of economic benefit but, more importantly, will have substantial benefits for patients, as they would recover quicker and be less likely to require multiple 'trial-and-error' treatments. Data from the present study support the associations of SNPs (Table 2) present in the

IFN-λ genes with HCV clearance after interferon and ribavirins combined therapy in Pakistani individuals infected with genotype 3a and provide preliminary evidence to suggest patients should be genotyped for the relevant SNPs in order to predict drug response before starting therapy.

Appendix 1. The Details of Single Nucleotide Polymorphisms (SNPs) Present in the up- and Down- Stream Region of IFNL-λ Genes. The Annotation of SNPs According to Their Position is Listed With Their Hardy-Weinberg Equilibrium P Values (HW p)

SNP RS No.	SNP Position	Role of SNP	Alleles	HW p
RS11083519	chr19:39719263	IFNL3 Downstream	A:T	0.820
RS955155	chr19:39729479	IFNL3 Downstream	C:T	0.304
RS35790907	chr19:39730755	IFNL3 Downstream	A:T	0.551
RS12972991	chr19:39731747	IFNL3 Downstream	A:C	0.831
RS12980275	chr19:39731783	IFNL3 Downstream	A:G	0.551
RS12982533	chr19:39731904	IFNL3 Downstream	T:C	0.551
RS8105790	chr19:39732501	IFNL3 Downstream	T:C	0.906
RS688187	chr19:39732752	IFNL3 Downstream	G:A	0.394
RS4803217	chr19:39734220	IFNL4 Exon	C:A	0.919
RS12979860	chr19:39738787	IFNL4 Intron	C:T	0.173
RS4803221	chr19:39739129	IFNL3 Promoter	C:G	1.000
RS1549928	chr19:39739709	IFNL3 Promoter	A:G	0.625
RS10853727	chr19:39740463	IFNL3 Promoter	T:C	0.118
RS8109886	chr19:39742762	IFNL3 Promoter	C:A	0.339
RS8113007	chr19:39743103	IFNL3 Promoter	A:T	0.225
RS8099917	chr19:39743165	IFNL3 Promoter	T:G	0.906
RS7248668	chr19:39743821	IFNL3 Promoter	G:A	0.906
RS16973285	chr19:39744696	IFNL3 Promoter	C:T	0.081
RS10853728	chr19:39745146	IFNL3 Promoter	G:C	0.387
RS12980602	chr19:39752820	IFNL2 Promoter	T:C	0.041
RS4803224	chr19:39753014	IFNL2 Promoter	G:C	0.976
RS11671087	chr19:39761790	IFNL2 Downstream	T:C	0.122
RS11665818	chr19:39768216	IFNL2 Downstream	G:A	0.039
RS7248931	chr19:39781583	IFNL1 Promoter	A:G	0.812

Appendix 2. The Details of SNPs Located in IL28B Gene (IFNL-3) Listed According to Amino Acid Position. The Amino Acid Present in Normal (amino acid: context) and Change of Amino Acid Due to Allele Change (amino acid: SNP) Are Listed Accordingly

SNP rs No.	Amino Acid Posi- tion.	SNP Position	Amino Acid: Context	Amino Acid: SNP	Allele Change
RS200289435	1	chr19:39735606	Methionine	Threonine	ATG →ACG
RS143935261	1	chr19:39735607	Methionine	Valine	ATG →GTG
RS202126177	2	chr19:39735603	Threonine	Serine	ACC →ATC
RS630388	2	chr19:39735602	Threonine	Threonine	ACC →ACT
RS150569967	3	chr19:39735601	Glycine	Arginine	GGG →AGG
RS199952257	57	chr19:39735438	Lysine	Arginine	AAA →AGA

RS202143862	72	chr19:39735101	Arginine	Cysteine	CGC →TGC
RS145428712	101	chr19:39734754	Threonine	Methionine	ACG →ATG
RS200889156	104	chr19:39734744	Valine	Valine	GTT →GTC
RS148543092	108	chr19:39734734	Threonine	Alanine	ACC →GCC
RS202101632	108	chr19:39734732	Threonine	Threonine	ACC →ACT
RS201376760	114	chr19:39734716	Alanine	Threonine	GCC →ACC
RS199801376	116	chr19:39734708	Glycine	Glycine	GGG →GGA
RS200058568	123	chr19:39734687	Leucine	Leucine	CTT →CTC
RS201605224	126	chr19:39734678	Leucine	Leucine	CTG →CTT
RS199655870	132	chr19:39734662	Glutamine	Stop Codon	CAG →TAG
RS149832972	133	chr19:39734659	Leucine	Phenylalanine	CTC →TTC
RS139176035	134	chr19:39734656	Arginine	Tryptophan	CGG →TGG
RS201566097	138	chr19:39734544	Glutamine	Stop Codon	CAG →TAG
RS145946971	164	chr19:39734465	Lysine	Threonine	AAG →ACG
RS143748522	179	chr19:39734328	Phenylalanine	Valine	TTC →GTC
RS150748693	180	chr19:39734325	Arginine	Cysteine	CGC →TGC
RS201746548	183	chr19:39734314	Threonine	Threonine	ACG →ACA
RS200180353	191	chr19:39734290	Serine	Serine	AGC →AGT
RS201888594	194	chr19:39734282	Leucine	Proline	CTG →CCG

Appendix 3 . The Primers Used for Detection and Genotyping of HCV. (HCF1: HCV Outer Forward Primer, HCR1: HCV Outer Reverse Primer, HCF2: HCV Internal Forward Primer, HCR2: HCV Internal Reverse Primer, HCGF1: HCV Genotype Outer Forward Primer, HCGR1: HCV Outer Reverse Primer, HCGF2: HCV Internal Forward Primer, Rest All Are Specific for Every HCV Genotype With Their Amplified Product Size Using Same Internal Primer

Primer Name	5'-3' Sequence	Product Size (bp)
HCF1	CCCTGTGAGGAACCTACTGTCTTCACGC	270
HCR1	ACTCGCAAGCACCCATCAGGCAGTAC	
HCF2	AAAGCGTCTAGCCATGGCG	210
HCR2	CACAAGGCCTTCGCGACC	
HCGF1	TTGTGGTACTGCCTGATAGGG	470
HCGR1	GGATGTACCCATGAGGATCG	
HCGF2	GTGCCCGGGAGGTCTCGTAG	
G1a	ACTCCACCAACGATCTGACC	129
G1b	AGCCTTGGGGATAGGTTGTC	233
G1c	CTTACCAAATTGCGTGACC	391
G2a	CTCCGAAGTCTCTTGTGCG	190
G2b	AGCAAGTAAACTCCGCCAAC	178
G2c	ACCGTTCGGAAGTTTCCTC	202
G3a	ACTCCACCAACGATCTGCC	258
G3b	AGCCTTGGGGATAAGGTGAC	232
G3c	GTGACCGCTCGGAAGTCTTA	197
G4a	CCGTAAGAGGCCATGGATA	288
G5a	AATCCGCACGTTAGGGTATG	417
G6a	CAGCCTCGCTTCCATAAAG	300

Appendix 4. The Primers and Probes Used During the iPLEX Assay on SEQUENOM are Given in Detail With Each SNP Corresponding to the Sequence of Forward and Reverse Primer With the Mass (Daltons) of PCR Product After First PCR. The Extended Product and Mass Represents the Change of Mass With the Different Incorporation of Base and Thus Explaining the Principal Behind the iPLEX Assay (PCR Mass: Mass of Initial PCR Product, Ext.1 and 2 Products: The Extended Base Which is Complementary to one Present in Initial PCR Product, Ext.1 and 2 Mass: The Masses of Final Products)

SNP ID	Forward Primer	Reverse Primer	PCR Mass	Probe	Ext. 1 Product	Ext. 1 Mass	Ext. 2 Product	Ext.2 Mass
RS12979860	ACGTGGATGTCGTGCC- GTCGTGACTGA	ACGTGGATGAGGCGC- GAGTCGAATCAAC	4563	AGCTCCCCGAAGGGC	C	4810.2	T	4890.1
RS143748522	ACGTGGATGTCCTCCCA- CAGGAGTCCC	ACGTGGATGCAAC- CAANTCAGGTCTCGC	4752.1	TGTCACCTCAACCTC	C	5039.3	A	5079.2
RS201566097	ACGTGGATGTCAGC- GTCCTTCCCCCTG	ACGTGGATGATGAGC- GTCCTGGCCCTGCGT	5115.3	GAECTGCCCACAGATC	G	5362.5	A	5442.4
RS148543092	ACGTGGATGTCGGC- CAAGACATCCCCAG	ACGTGGATGCTGACGT- GAAGGTCTCTG	5242.4	CTGGTAGTGTAGCGGG	C	5489.6	T	5569.5
RS11881222	ACGTGGATGACACTG- TACCCCTTCC	ACGTGGATGAGAACAGT- GAAGGTGACAG	5282.4	ACCCCTTCTCTGCTCC	G	5529.6	A	5609.5
RS3105790	ACGTGGATGCTCTC- GACATCACTCCAAAT	ACGTGGATGAGCAT- CATTAGCGGAAG	5394.5	CATCACTCCAATGCTCTG	C	5641.7	T	5721.6
RS202126177	ACGTGGATG- GCTCCCTTCTCTGTGAC	ACGTGGATGACAGGAAC- GCTCCAGTCAC	5796.8	CTCTGACACAGACATGA	G	6044	C	6084
RS150748693	ACGTGGATGAGGCCCT- GTCACCTCAAC	ACGTGGATGTCAT- GACTGGCGGAAGG	5938.9	CTGTACCTTCAACCTCTC	G	6186.1	A	6266
RS10665818	ACGTGGAT- GAAGAAAGACCTCCAC- CATGC	ACGTGGATGAGTGT- CACCCCTATTCATCTAGC	5947.9	TATCATCTGCCCAACTC	A	6219.1	G	6235.1
RS4803221	ACGTGGATGTCGTG- CACGGTGAATGCC	ACGTGGATGCCCTCAGC- GCCTTGGCAG	6319.1	CCCAAGGGCTGCCT- GCTCTC	G	6566.3	C	6606.3
RS19980376	ACGTGGATGATGGTG- CAGGGTGTGAAG	ACGTGGATGCTGAGCT- GAAGGTCTCTG	6456.2	ACGGGGCTGGTCCAGAGA- CATC	C	6703.4	T	6783.3
RS12980602	ACGTGGATGACTTTATT- AGTGGTAAAC	ACGTGGATGCTCTG- GTTTTGTCTCATCTG	6505.3	GAACAAATGAAAGG- CAGAGA	C	6752.5	T	6832.4
RS1549928	ACGTGGATGTCCTC- CAACACTGGTT	ACGTGGATGAGAAGATA- AAGACAACAGG	6643.3	GCCTAAATTGTCCTGTCCT- GT	G	6890.5	A	6970.4
RS688187	ACGTGGATGTCCTATT- GAATCCATTAC	ACGTGGATGCTGACAC- TAACAGTCACAG	6651.4	GCACATGCGCAACACAC- CACA	A	6922.6	G	6938.6
RS12980275	ACGTGGATGTCCTATT- ACCCCTCCCGC	ACGTGGATGATGAGGT- GCTGAGAGAAGTC	7016.6	ACGGGAAATATTAGA- CACGTC	G	7263.8	A	7343.7
RS11671087	ACGTGGAT- GAAGCTCCCTGGAG- TAAC	ACGTGGATGAGATGC- CACCCCAAAGTC	7056.6	CCTGTCGGAGTAACATAA- GATA	C	7303.8	T	7383.7

RS139176035	ACGTGGATGTCA- CACCTGACCATTC	ACGTGGATGTCT- CACAGCTCACAGACCT	7168.7	CTGAAC- CATATCCTCTCCAGCTC	G	7415.9	A	7495.8
RS4803217	ACGTGGATGATAATAGC- GAATGGTGAC	ACGTGGATGCCAGTCATG- CAACCTGAGAT	7449.9	GCGACTGGGTGACATA- AATTAGG	C	7697.1	A	7721.1
RS7248668	ACGTGGATGGAGTGGC- GATTGTCACCA	ACGTGGATGGCTTGTG- CAAGAGAGGGTTG	7552.9	CCAGAATTGTCACACTAC- TATGCTC	G	7800.1	A	7880
RS16973285	ACGTGGATGGCAC- GTTTCAATTGTTA	ACGTGGATGATCTTA- GCCACCCATCTTA	7593.9	CACGTTCATTGTTATT- GATTC	C	7841.1	T	7921
RS12982533	ACGTGGATGAAGAGA- GTTCTGGAGATTGC	ACGTGGATGTTACAG- GTCTGGTCCTAGT	7762	GGGTGAGATGCTTGCCT- GAACATG	C	8009.2	T	8089.1
RS3113007	ACGTGGAT- GACAAAAGGAGGAAACAGT- GAC	ACGTGGATGGAGAGTA- AAAGTAAGTCITG	7960.2	TGACAATTTGTA- AAAAATATTIACC	T	8231.4	A	8287.3
RS201605224	ACGTGGATGATGTCITG- GACCAGCCCCCTT	ACGTGGATGGCCCTGAC- GACTCACACA	8132.3	TGTGGGAC- CAGCCCCCTCACACCCCT	C	8419.5	A	8459.4
RS7248931	ACGTGGATGCTCATCTCT- CAAGAACCTAGG	ACGTGGATGGCTCATCT- TATGATGGC	8333.5	GATAATGAAACTAG- GAAAATCTCAAG	G	8580.7	A	8660.6
RS2000058568	ACGTGGATGGTGCACAT- GACCCAGCCCT	ACGTGGATGGCCCTGAC- GACTCACACA	4488.9	CTTGGGACAGCCCTT	G	4736.1	A	4816
RS202101632	ACGTGGATGCTGACGCT- GAAGGTCTG	ACGTGGATGGTGCCT- CAAGACATCCCCAG	4609	GGTTCCTGGAGGCCAC	G	4856.2	A	4936.1
RS145946971	ACGTGGATGCGCTC- CACCATGGCTG	ACGTGGATGCTCTCTC- CAGCTGCCCA	4893.2	CAGGAGCCCCAAAAAA	G	5140.4	T	5164.4
RS202143862	ACGTGGATGTCACCTG- CAGCTGCCCA	ACGTGGATGCTTGTCT- GTCTAGGAAGAG	5036.3	TGAAAGAGGGGGAGGC	A	5307.5	G	5323.5
RS201376760	ACGTGGATG- GCTCCCTTCTCTGTGAC	ACGTGGATGACAGGAACCT- GTTCAGTCAC	5100.3	CCGCTGACACTGACCA	T	5371.5	C	5387.5
RS630388	ACGTGGATGCTGACGCT- GAAGGTCTG	ACGTGGATGAGGGCTG- GTCAAGACAT	5203.4	TGTGACACAGACATGAC	G	5450.6	A	5530.5
RS955155	ACGTGGATGAC- TATGGGGCAACACTGTC	ACGTGGATGACTGGTAT- GTAGCTCCCTCG	5211.4	TGTGACTGAGGGCCCA	T	5482.6	C	5498.6
RS10853728	ACGTGGATGAGA- CAGACTCTCATCCCTCAC	ACGTGGATGTCATTCATC- CATTCCTGCTC	5676.7	CCATCCCTACCAAAAGCTTA	G	5923.9	C	5963.9
RS201746548	ACGTGGATGGTGCAT- GAATGGGGAA	ACGTGGATGCTCTGT- CACCTCTAACCTC	5756.8	CAACACAAATTCAAGGCTCG	C	6004	T	6083.9
RS199952257	ACGTGGATGATGGT- GACCTTGGAGTGC	ACGTGGATGGAGGCTG- CAGGCCCTTAAG	5787.8	TGGACTCAAGGATCT	C	6035	T	6114.9
RS35790907	ACGTGGATGACATGTC- GAGAGCGAAATC	ACGTGGATGCTCTGCG- CAGGTAGAAGC	5885.9	GCTGTACAGGTGAGAACAA	A	6157.1	T	6212.9

RS2000180353	ACGTTGGATGTCCTAGGT- GCATGACCTGGC	ACGTTGGATGCTCACGC- GAGCCIGAAATIG	6336.1	CITTCAGACACACAG- GTCCCC	A	6607.3	G	6623.3
RS8099917	ACGTTGGATGCAATTGT- CACTGTTCTCC	ACGTTGGATGACTATGA- CAGCATGGTTC	6368.1	TTTTCTTCTGTGAG- CAAT	G	6655.4	T	6695.2
RS11083519	ACGTTGGATGAAAGC- CAACTCAATTGAGG	ACGTTGGATGTTGATC- CACTTTCTGCC	6460.2	TTGAG- GAAGAATAGCCTTTC	A	6731.4	T	6787.3
RS10853727	ACGTTGGATGACGCTAC- CATTTGGTGAAC	ACGTTGGATGATGATA- AGCATGCGCAGAGAG	6825.5	GAAGACATCATATGAAAGAG- GCA	C	7072.7	T	7152.6
RS4803224	ACGTTGGATGATGCTCTA- AGCAGCTGGAG	ACGTTGGATGACACAGT- GAGACCCCATC	6994.5	GCTTGAAGCTGAG- GCACCCACCA	G	7241.7	C	7281.8
RS2000889156	ACGTTGGATGCGGTG- GCTTGGAGGTGA	ACGTTGGATGTTGATT- GAGACAGACAGAGC	4593	CTTGACGGCTGAAGGT	G	4840.2	A	4920.1
RS8109886	ACGTTGGATGTTCTCCT- GTCTCTCTCTGGC	ACGTTGGATGTTGATT- GAGACAGACAGAGC	4810.2	TCCAAACAGCATCTTG	C	5057.3	A	5081.4
RS200289435	ACGTTGGATG- GCTCCCTTCCTCTCTGAC	ACGTTGGATGACAGGAACAT- GCTCCAGTCAC	5154.4	TCTCTGTGACACACACA	G	5401.6	A	5481.5
RS201888594	ACGTTGGATGTCAG- GTTCATGACTTGG	ACGTTGGATGGGGAGACT- GAATTGTGTTG	5253.4	GGAAAGGGTAGACACACAC	A	5524.6	G	5540.6
RS199655870	ACGTTGGATGATGTTG- GACCAGCCCCCT	ACGTTGGATGGCCCTGAC- GACTCACACA	5355.5	CGGCACCATATCCCTCC	G	5602.7	T	5626.7
RS12972991	ACGTTGGATGGAAATTI- GACTCACACA	ACGTTGGATGAGACT- GAATAAGCCAGCTC	5435.5	GGCTCTAGCACCTCATG	C	5722.7	A	5762.6
RS149832972	ACGTTGGATGGCCCTGAC- GACTCACACA	ACGTTGGATGATGTCCTTG- GACCAAGCCCCCT	4547	TCACACAGCCCCGA	A	4818.2	G	4834.2
RS143935261	ACGTTGGATG- GCTCCCTTCCTCTGAC	ACGTTGGATGACAGGAACAT- GCTCCAGTCAC	5130.4	CTCTCTGTGACACACAC	T	5401.6	C	5417.6
RS145428712	ACGTTGGATGAGGGGCTG- GICCAAGACAT	ACGTTGGATGCGGTG- GCTTGGAGGCTGA	4786.1	CTCCAGAACCTTCAGC	A	5057.3	G	5073.3
RS150569967	ACGTTGGATGTTCTCT- GTGACACAGAC	ACGTTGGATGACAGGAACAT- GCTCCAGTCAC	4859.2	TGACACAGACATGACC	T	5130.4	C	5146.4

Appendix 5. Prediction of Response to Therapy With Homozygous Responder IFN- λ Region SNPs^a

SNP	Sensitivity, % (95% CI)	Specificity, % (95%CI)	Prevalence, % (95%CI)	PPV, % (95%CI)	NPV, % (95%CI)
RS955155	51.39 (42.92-59.79)	53.57 (39.75-67.01)	72.00 (65.23-78.10)	74.00 (64.27-82.26)	30.00 (21.24-39.98)
RS12972991	59.35 (50.12-68.11)	64.94 (53.21-75.46)	61.50 (54.38-68.28)	73.00 (63.20-81.39)	50.00 (39.83-60.17)
RS8105790	57.14 (48.28-65.68)	64.18 (51.53-75.53)	66.50 (59.50-73.00)	76.00 (66.43-83.97)	43.00 (33.14-53.29)
RS688187	52.48 (43.91-60.95)	55.93 (42.40-68.84)	70.50 (63.66-76.72)	74.00 (64.27-82.26)	33.00 (23.92-82.26)
RS4803217	52.48 (43.91-60.95)	55.93 (42.40-68.84)	70.50 (63.66-76.72)	74.00 (64.27-82.26)	33.00 (23.92-82.26)
RS12979860	56.92 (47.95-65.57)	62.86 (50.48-74.11)	65 (57.95-71.59)	74.00 (64.27-82.26)	44.00 (34.08-54.28)
RS4803221	51.23 (42.80-60.04)	53.23 (40.12-66.01)	69.00 (62.09-75.33)	71.00 (61.07-79.64)	33.00 (23.92-43.12)
RS8109886	60.09 (52.55-68.92)	79.63 (66.47-89.35)	73.00 (66.28-79.02)	89.00 (81.17-94.37)	43.00 (33.14-53.29)
RS8113007	52.48 (43.91-60.95)	55.93 (42.40-68.84)	70.05 (63.66-76.72)	74.00 (64.27-82.26)	33.00 (23.92-43.12)
RS8099917	42.01 (34.47-49.83)	6.45 (0.98-21.26)	84.50 (78.73-89.22)	71.00 (61.07-79.64)	2.00 (0.30-7.05)
RS7248668	42.01 (34.47-49.83)	6.45 (0.98-21.26)	84.50 (78.73-89.22)	71.00 (61.07-79.64)	2.00 (0.30-7.05)
RS11671087	59.35 (50.12-68.11)	64.94 (53.21-75.46)	61.50 (54.38-68.28)	73.00 (63.20-81.39)	50.00 (39.20-81.39)
RS11665818	62.30 (53.07-70.91)	69.23 (57.76-79.19)	61.00 (53.87-67.80)	76.00 (66.43-83.97)	54.00 (43.74-64.01)

^a Abbreviations: 95% CI, 95% confidence interval; PPV, Positive predictive value; NPV, Negative predictive value

Acknowledgements

We would like to thank the technical staff at CIGMR and National Genetics Laboratory, Lahore for making this study possible by contributing their professional skills. This study was supported by the Higher Education Commission Pakistan.

Author' Contribution

Conceived and designing the experiments: Philip Day, Fiona Marriage, Amin Athar, Imran Tipu; performed the experiments: Imran Tipu, Hazel Plat, Zia Farooqi, analysed the data: Andrea Short, Imran Tipu; wrote the paper: Imran Tipu, Philip Day, Andrea Short.

Conflict of Interest

We do not have any conflict of interest.

Financial Disclosure

Imran Tipu received financial assistance from Higher education commission of Pakistan to conduct this research.

Funding/Support

This study was the supported by international research support initiative program (IRSIP) grant code: IRSIP 21: BMS 38. By Higher education commission of Pakistan.

References

1. Lavanchy D. Evolving epidemiology of hepatitis C virus. *Clin Microbiol Infect.* 2011;17(2):107-15.
2. Rehman IU, Idrees M, Ali M, Ali I, Butt S, Hussain A, et al. Hepatitis C virus genotype 3a with phylogenetically distinct origin is circulating in Pakistan. *Genet Vaccines Ther.* 2011;9(1):2.
3. Azhar MA, Bukhari MH, Ghanni U, Khan A, Malik JI, Shah AH. Prevalence of hepatitis C virus and its serotypes in Bahawalpur Division. *Biomed.* 2003;19:18-22.
4. Attaullah S, Khan S, Ali I. Hepatitis C virus genotypes in Pakistan: a systemic review. *Virol J.* 2011;8:433.
5. Amini M, Poustchi H. Hepatitis C virus spontaneous clearance: immunology and genetic variance. *Viral Immunol.* 2012;25(4):241-8.
6. Raulet DH. Interplay of natural killer cells and their receptors with the adaptive immune response. *Nat Immunol.* 2004;5(10):996-1002.
7. Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature.* 2009;461(7262):399-401.
8. Cox AL, Netski DM, Mosbruger T, Sherman SG, Strathdee S, Ompad D, et al. Prospective evaluation of community-acquired acute-phase hepatitis C virus infection. *Clin Infect Dis.* 2005;40(7):951-8.
9. Bonis PA. Clinical outcomes after hepatitis C infection from contaminated anti-D immune globulin. *N Engl J Med.* 1999;341(10):763.
10. Suppiah V, Moldovan M, Ahlenstiell G, Berg T, Weltman M, Abate ML, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet.* 2009;41(10):1100-4.
11. Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet.* 2009;41(10):1105-9.
12. Sheppard P, Kindsvogel W, Xu W, Henderson K, Schlutsmeyer S, Whitmore TE, et al. IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol.* 2003;4(1):63-8.
13. Prokunina-Olsson L, Muchmore B, Tang W, Pfeiffer RM, Park H, Dickensheets H, et al. A variant upstream of IFNL3 (IL28B) creating a new interferon gene IFNL4 is associated with impaired clearance of hepatitis C virus. *Nat Genet.* 2013;45(2):164-71.
14. Marcello T, Grakoui A, Barba-Spaeth G, Machlin ES, Kotenko SV, MacDonald MR, et al. Interferons alpha and lambda inhibit hepatitis C virus replication with distinct signal transduction and gene regulation kinetics. *Gastroenterology.* 2006;131(6):1887-98.
15. Gad HH, Dellgren C, Hamming OJ, Vends S, Paludan SR, Hartmann R. Interferon-lambda is functionally an interferon but

structurally related to the interleukin-10 family. *J Biol Chem.* 2009;284(31):20869-75.

16. Pagliaccetti NE, Eduardo R, Kleinstein SH, Mu XJ, Bandi P, Robek MD. Interleukin-29 functions cooperatively with interferon to induce antiviral gene expression and inhibit hepatitis C virus replication. *J Biol Chem.* 2008;283(44):30079-89.
17. Sommereyns C, Paul S, Staeheli P, Michiels T. IFN-lambda (IFN-lambda) is expressed in a tissue-dependent fashion and primarily acts on epithelial cells in vivo. *PLoS Pathog.* 2008;4(3).
18. Rauch A, Katalik Z, Descombes P, Cai T, Di Julio J, Mueller T, et al. Genetic Variation in IL28B Is Associated With Chronic Hepatitis C and Treatment Failure: A Genome-Wide Association Study. *Gastroenterol.* 2010;138(4):1338-45. e7.
19. Abe H, Ochi H, Maekawa T, Hayes CN, Tsuge M, Miki D, et al. Common variation of IL28 affects gamma-GTP levels and inflammation of the liver in chronically infected hepatitis C virus patients. *J Hepatol.* 2010;53(3):439-43.
20. Mangia A, Thompson AJ, Santoro R, Piazzolla V, Tillmann HL, Patel K, et al. An IL28B polymorphism determines treatment response of hepatitis C virus genotype 2 or 3 patients who do not achieve a rapid virologic response. *Gastroenterology.* 2010;139(3):821-1.
21. McCarthy JJ, Li JH, Thompson A, Suchindran S, Lao XQ, Patel K, et al. Replicated association between an IL28B gene variant and a sustained response to pegylated interferon and ribavirin. *Gastroenterology.* 2010;138(7):2307-14.
22. Thompson AJ, Muir AJ, Sulkowski MS, Ge D, Fellay J, Shianna KV, et al. Interleukin-28B polymorphism improves viral kinetics and is the strongest pretreatment predictor of sustained virologic response in genotype 1 hepatitis C virus. *Gastroenterology.* 2010;139(1):120-18.
23. Bochud PY, Bibert S, Negro F, Haagmans B, Soulier A, Ferrari C, et al. IL28B polymorphisms predict reduction of HCV RNA from the first day of therapy in chronic hepatitis C. *J Hepatol.* 2011;55(5):980-8.
24. Smith KR, Suppiah V, O'Connor K, Berg T, Weltman M, Abate ML, et al. Identification of improved IL28B SNPs and haplotypes for prediction of drug response in treatment of hepatitis C using massively parallel sequencing in a cross-sectional European cohort. *Genome Med.* 2011;3(8):57.
25. Yu ML, Huang CF, Huang JF, Chang NC, Yang JF, Lin ZY, et al. Role of interleukin-28B polymorphisms in the treatment of hepatitis C virus genotype 2 infection in Asian patients. *Hepatology.* 2011;53(1):7-13.
26. Chen JY, Lin CY, Wang CM, Lin YT, Kuo SN, Shiu CF, et al. IL28B genetic variations are associated with high sustained virological response (SVR) of interferon-alpha plus ribavirin therapy in Taiwanese chronic HCV infection. *Genes Immun.* 2011;12(4):300-9.
27. Scherzer TM, Hofer H, Staettelmayer AF, Rutter K, Beinhardt S, Steindl-Munda P, et al. Early virologic response and IL28B polymorphisms in patients with chronic hepatitis C genotype 3 treated with peginterferon alfa-2a and ribavirin. *J Hepatol.* 2011;54(5):866-71.
28. Ridruejo E, Solano A, Marciano S, Galdame O, Adrover R, Cocozella D, et al. Genetic variation in interleukin-28B predicts SVR in hepatitis C genotype 1 Argentine patients treated with PEG IFN and ribavirin. *Ann Hepatol.* 2011;10(4):452-7.
29. Xie JQ, Guo XY, Zhang XH, Lin BL, Xie DY, Gao ZL, et al. Relationship between the genetic variation in interleukin 28B and response to antiviral therapy in patients with chronic hepatitis C. *Chin Med J (Engl).* 2012;125(13):2334-8.
30. Pedergnana V, Abdel-Hamid M, Guergnon J, Mohsen A, Le Fouler L, Theodorou I, et al. Analysis of IL28B variants in an Egyptian population defines the 20 kilobases minimal region involved in spontaneous clearance of hepatitis C virus. *PLoS One.* 2012;7(6).
31. Shi X, Pan Y, Wang M, Wang D, Li W, Jiang T, et al. IL28B genetic variation is associated with spontaneous clearance of hepatitis C virus, treatment response, serum IL-28B levels in Chinese population. *PLoS One.* 2012;7(5).
32. de Castellarnau M, Aparicio E, Parera M, Franco S, Tural C, Clotet B, et al. Deciphering the interleukin 28B variants that better predict response to pegylated interferon-alpha and ribavirin therapy in HCV/HIV-1 coinfected patients. *PLoS One.* 2012;7(2).
33. Grandi T, da Silva CM, Amaral KM, Picon PD, Costi C, da Fre NN, et al. Response to treatment in Brazilian patients with chronic hepatitis C is associated with a single-nucleotide polymorphism near the interleukin-28B gene. *Mem Inst Oswaldo Cruz.* 2013;108(1):48-53.
34. Stenkvist J, Sonnerborg A, Weiland O. HCV RNA decline in chronic HCV genotype 2 and 3 during standard of care treatment according to IL28B polymorphism. *J Viral Hepat.* 2013;20(3):193-9.
35. Gelinas JF, Fabre T, Willems P, Leung RC, George J, Willems B, et al. IL28B SNP screening and distribution in the French Canadian population using a rapid PCR-based test. *Immunogenetics.* 2013;65(6):397-403.
36. Ezzikouri S, Alaoui R, Rebbani K, Brahim I, Fakhir FZ, Nadir S, et al. Genetic variation in the interleukin-28B gene is associated with spontaneous clearance and progression of hepatitis C virus in Moroccan patients. *PLoS One.* 2013;8(1).
37. Jung YK, Kim JH, Ahn SM, Yang JW, Park SJ, Kim JW, et al. Role of interleukin 28B-related gene polymorphisms in chronic hepatitis C and the response to antiviral therapy in Koreans. *J Clin Gastroenterol.* 2013;47(7):644-50.
38. Aziz H, Raza A, Murtaza S, Waheed Y, Khalid A, Irfan J, et al. Molecular epidemiology of hepatitis C virus genotypes in different geographical regions of Punjab Province in Pakistan and a phylogenetic analysis. *Int Infect Dis.* 2013;17(4):e247-53.
39. Honda M, Sakai A, Yamashita T, Nakamoto Y, Mizukoshi E, Sakai Y, et al. Hepatic ISG expression is associated with genetic variation in interleukin 28B and the outcome of IFN therapy for chronic hepatitis C. *Gastroenterology.* 2010;139(2):499-509.