



Is Affected and Unaffected Side Lateral Spinal Mobility Different in Unilateral Cerebral Palsy? A Cross-sectional Study

Seda Ayaz Taş^{1,*}, Tamer Çankaya¹

¹ Physiotherapy and Rehabilitation Department, Faculty of Health Sciences, Bolu Abant Izzet Baysal University, Bolu, Turkey

*Corresponding Author: Physiotherapy and Rehabilitation Department, Faculty of Health Sciences, Bolu Abant Izzet Baysal University, Bolu, Turkey. Email: ptsedaayaztas@gmail.com

Received: 1 February, 2025; Revised: 9 April, 2025; Accepted: 19 May, 2025

Abstract

Background: Asymmetric posture is a characteristic feature of unilateral cerebral palsy (UCP). There is limited information on how this postural disturbance affects lateral spinal mobility on the affected and unaffected sides.

Objectives: The present study aimed to compare lateral trunk mobility on the affected and unaffected sides in children with UCP.

Methods: This cross-sectional study included 60 children with UCP, conducted at Duzce Gokkusagi and Eregli Gokkusagi special education and rehabilitation centers from October 2023 to October 2024. Participants were aged 6 - 18 years and classified as Gross motor function classification system (GMFCS) levels I and II. Two children were excluded due to scoliosis surgery, resulting in a final sample of 58 children. Demographic characteristics such as height, body weight, gender, age, and affected side were recorded. Eligible children, who voluntarily participated, were evaluated by the same examiner. Spinal posture and mobility were assessed in the frontal plane using the spinal mouse (SM) in a standing position. Spinal mobility was evaluated by comparing trunk lateral flexion angles on the affected and unaffected sides. Descriptive statistics were presented as mean and standard deviation for continuous variables and frequency and percentage for categorical variables. The Shapiro-Wilk test was used to assess normal distribution. A dependent sample *t*-test was performed to compare bilateral lateral spinal mobility, with a significance level set at $P < 0.05$.

Results: Among the children with UCP, 62.9% had right-sided involvement, and 37.1% had left-sided involvement. No significant difference was found between the affected and unaffected side spinal lateral mobility angles of the sacrum-hip ($P = 0.353$), thoracic ($P = 0.602$), lumbar ($P = 0.079$), and inclination ($P = 0.352$). However, there was a significant difference in sacrum-hip ($P = 0.011$) and lumbar ($P = 0.000$) angles in right and left lateral flexion mobility.

Conclusions: This study revealed that children with UCP and asymmetric postural patterns have similar lateral spinal mobility on the affected and unaffected sides. However, differences in right and left lateral mobility angles at certain levels were observed, potentially due to the children's postural patterns. Further research is needed to investigate spinal posture and mobility specific to different postural patterns.

Keywords: Cerebral Palsy, Posture, Spinal Mouse, Spine, Trunk

1. Background

Damage to the developing brain results in impaired postural control and motor function in cerebral palsy (CP) (1). The location of involvement in CP can be either unilateral or bilateral. Unilateral cerebral palsy (UCP), which affects one side of the body, including the extremities and trunk, represents approximately 30% of all CP cases (2). Bilateral cerebral palsy (BCP) is

characterized by the involvement of both sides of the body, including the extremities and torso (3). As the severity of motor impairment increases in CP, postural control is negatively impacted. Although children with UCP tend to have better functional outcomes, their postural control is still affected (4, 5), and they are at risk for developing asymmetric spinal posture (6). While children with CP are born with a normal musculoskeletal system, they are at significant risk for

the early development and rapid deterioration of asymmetric postural issues and abnormal spinal movement patterns (7).

In children with CP, abnormal posture may emerge as a result of deficiencies in the spinal alignment process, leading to secondary spinal deformities and movement disorders. Early determination of such abnormalities may help prevent secondary disorders in CP (8). Posture-oriented assessment and intervention programs can play a role in preventing contractures, deformities, pain, and asymmetry (9). Trunk control plays a key role in effective postural control. The stabilization of the trunk enables the effective movement of the head and distal segments, supports the visual and vestibular control of posture, facilitates the development of goal-directed activities and gross motor skills, and allows for the proper alignment of the body for communication and social skills (10, 11). When trunk control is impaired during early life, it can significantly affect overall motor development and, through delayed and limited motor functions, even impact cognitive and emotional development in children. The interaction between the trunk, upper extremities, and head changes with age, depending on maturation. Children are more susceptible to developing spinal deformities than adults due to muscle weakness, which also affects trunk movement and stability (12). Trunk movements are important for maintaining mobility and postural reactions during limb movements (8).

In children with CP, abnormal posture may emerge as a result of deficiencies in the spinal alignment process, leading to secondary spinal deformities and movement disorders (9). The disorder consisting of pathologic muscle tone, increased angle of curvature, and secondary inhibitory vertebral growth disorders in the concavity of the curve predisposes to progressive vertebral and spinal deformity as well as curve stiffness (13). Scoliosis has been reported in 40% of children with UCP (14). Postural deformity may also begin when an individual spends too much time in a particular asymmetrical position (7). Asymmetric muscle activity does not clearly explain deformity patterns; no relationship has been established between the direction of increased muscle tone and the direction of scoliosis (15). Galvis et al. found that the lateral flexion angles of healthy adolescents with scoliosis were similar on both sides. Increased mobility was observed especially in the segments below and above the apex of the scoliotic curve, indicating that the scoliotic spine is flexible and can compensate near the apex (16).

Healthcare professionals need a reliable and valid method to estimate spinal range of motion limitations

to support clinical decisions regarding individual prognosis and management (17). Evaluation of lateral mobility on the affected and unaffected side in children with UCP with asymmetric postural involvement will clinically shed light on interventions in terms of providing information about the flexibility of the spine and development of right posture. Posture-oriented assessment and intervention programs can play a role in preventing contractures, deformities, pain, and asymmetry (18).

2. Objectives

There are few studies that describe asymmetric spinal alignment in UCP (6, 14). However, no study has compared spinal alignment and spinal mobility between the affected and unaffected sides in UCP. The present study is to compare spinal posture and lateral spinal mobility between the affected and unaffected sides in children with UCP.

3. Methods

3.1. Participants

This cross-sectional study included 60 children with UCP attending Duzce Gokkusagi and Eregli Gokkusagi special education and rehabilitation centers. Children who met the inclusion criteria and whose parents and themselves voluntarily agreed to participate were included. Based on power analysis, the minimum sample size was determined to be 36, with a 90% power and a 0.05 alpha error margin. Two children were excluded due to scoliosis surgery. Consequently, evaluations were conducted on 58 children with CP, and informed consent was obtained from both the children and their families. Inclusion criteria were: Being between the ages of 6 and 18 years with UCP, and being levels I or II according to the Gross motor function classification system (GMFCS). Exclusion criteria included having undergone scoliosis surgery and having a rigid contracture in the spine. Ethical approval was obtained from the Clinical Research Ethics Committee of Bolu Abant izzet Baysal University (Decision number: 2023/211, date: 19.09.2023), in accordance with the Helsinki Declaration.

3.2. Measurements

Demographic information, including age, gender, affected side, height, and body weight, was collected. Spinal posture and mobility were assessed using a computer-assisted and non-invasive electromechanical device, the Spinal Mouse (SM) (Idiag, Voletswil,

Switzerland) (Figure 1). Assessments were conducted in a quiet, calm room with the assessor, child, and parent present. Measurements were performed with the SM when the child had no clothing on the trunk and was not wearing any orthoses or shoes. The assessment with the SM was performed in a standing position with the feet shoulder-width apart in the frontal plane. The SM device was moved along the spinal processes from C7 to S2 on the skin surface. Data obtained from the device was transmitted via bluetooth to a computer and recorded. Three measurements were taken in the frontal plane: Standing posture in an upright position, lateral flexion of the trunk on the affected side, and lateral flexion of the trunk on the unaffected side (Figure 2).

3.3. Statistical Analysis

Sample size calculation was performed using G-Power 3.1.9.7 software. Since no study in the literature compares SM and lateral spinal mobility in children with UCP, it was determined that at least 36 participants were needed at 90% power and a 5% error level for a *t*-test in dependent groups with a medium effect size of $d = 0.5$. It was also assumed that there would be a 20% loss of participants, and it was determined that at least 44 participants should be studied. Statistical analyses were conducted using IBM SPSS Statistics for Windows, v. 26.0 (Armonk, NY: IBM) and R Statistical Software (v4.3.0; R Core Team, 2023). Descriptive statistics were calculated as mean and standard deviation for continuous variables, and frequency and percentage for categorical variables. The Shapiro-Wilk test was used to assess the normality assumption. For numerical variables, a dependent samples *t*-test was performed to compare bilateral lateral spinal mobility. A significance level of $P < 0.05$ was considered statistically significant.

4. Results

The demographic characteristics of the 58 children with UCP who participated in this study, including age, height, body weight, Body Mass Index, gender, and affected side, are presented in Table 1. The spinal posture angles and mobility comparisons in the frontal plane, measured with the SM, are shown in Table 2. No significant difference was found between the affected and unaffected sides in terms of spinal lateral mobility angles at the sacrum-hip ($P = 0.353$), thoracic ($P = 0.602$), lumbar ($P = 0.079$), and inclination ($P = 0.352$) angles. However, a significant difference was observed in right and left lateral flexion mobility, with differences in sacrum-hip ($P = 0.011$) and lumbar ($P = 0.000$) angles.

5. Discussion

In this study, which aimed to compare the spinal mobilities of the affected and unaffected sides in children with CP, it was concluded that the lateral spinal mobility angles of the affected and unaffected sides were approximately similar. It was observed that the involvement of the affected half of the trunk in CP did not negatively impact trunk lateral flexion mobility compared to the unaffected side. Studies have reported high rates of lateral curvature in children with UCP (14, 19). No difference was observed between the lateral spinal mobility angles of the trunk due to compensatory mechanisms secondary to these curvatures (16).

Postural patterns vary in children with UCP. In UCP, two primary postural patterns have been described: The pro-gravitational postural pattern (PGPP) and the anti-gravitational postural pattern (AGPP). The differences between PGPP and AGPP not only involve characteristic weight-bearing on the unaffected or affected side of the body but also include significant differences in the orientation of the spine, pelvis, and shoulder girdle. In children with AGPP, the pelvis exhibits upward obliquity on the affected side, whereas in children with PGPP, the pelvis shows downward obliquity. Lateral spinal curvature has been reported in the majority of the children examined (84%). In all children with AGPP, convexity of the spine is observed toward the unaffected side, while in children with PGPP, the convexity is directed toward the affected side. More importantly, in children with AGPP, the affected side bears less weight, whereas in children with PGPP, the affected side bears more weight. The characteristic postural pathology in all children with CP should be addressed, as it affects overall functional efficiency (19).

The postural patterns of the children included in our study were not assessed. The similarity between the lateral spinal mobility angles of the affected and unaffected sides in this study may be attributed to the postural patterns specific to CP. One of the findings of our study was that the sacrum-hip angles were greater on the right side. The majority of the children in this study (62.1%) had right-sided involvement. In the right lateral flexion position, which is the affected side for most children, sacrum-hip and thoracic angles were greater, while in the left lateral flexion position, lumbar angles were greater. We hypothesize that the increased lumbar mobility on the left side may serve as a compensatory mechanism. This result is based on the compensatory mechanisms that occur in the trunk as a result of postural deformities to keep the center of gravity within the support surface and maintain balance (16).




Figure 1. A, The lateral view of the spinal mouse (SM); and B, the results of the frontal plane measurements taken with the SM



Figure 2. The measurements taken with the spinal mouse (SM) in the frontal plane in the following positions: A, Standing posture; B, right trunk lateral flexion; and C, left trunk lateral flexion

Table 1. Demographic Characteristics of Participants with Unilateral Cerebral Palsy^a

Variables	UCP Children (n = 58)
Age (y)	9.4 (6 - 18)
Height (cm)	134.1 (92.5 - 179)
Weight (kg)	34.2 (12.5 - 105.2)
BMI (kg/m ²)	17.3 (11.8 - 32.87)
Gender	
Female	30 (51.7)
Male	28 (48.3)
Affected side	
Right	36 (62.1)
Left	22 (37.9)

Abbreviation: BMI, Body Mass Index; UCP, unilateral cerebral palsy.

^aValues are expressed as median IQR (25/75) or No. (%).

Table 2. Comparison of Lateral Spinal Mobility Values in Frontal Plane

Inclination Degree Curvatures; UP	Unilateral Cerebral Palsy Children; n = 58, Median (IQR 25/75)		
Sacrum-hip	4.8 (0/12)		
Thorax	8.3 (0/17)		
Lumbar	8.4 (0/20)		
Total spine	2.6 (0/13)		
Mobility	UP-Affected side Latflex P; n = 58, Median (IQR 25/75)	UP-Nonaffected side Latflex P; n = 58, Median (IQR 25/75)	P-Value ^a
Sacrum-hip	9.2 (0/37)	8 (0/35)	0.353
Thorax	24.2 (2/44)	23.1 (4/59)	0.602
Lumbar	12.2 (0/31)	16.3 (2/36)	0.079
Total spine	22.2 (7/42)	21.1 (4/48)	0.352
UP	UP-Right Latflex P	UP-Left Latflex P	P-Value ^a
Sacrum-hip	10.2 (0/37)	7 (0/35)	0.011
Thorax	25.6 (3/41)	21.9 (2/59)	0.066
Lumbar	8.1 (0/28)	20.4 (4/36)	0.000
Total spine	21.8 (7/34)	21.5 (4/48)	0.855

Abbreviations: IQR, interquartile range; UP, upright position; Latflex P, lateral flexion position.

^a Paired sample t-test.

Scoliosis is a deformity that negatively affects spinal alignment and mobility. In scoliosis, a smaller lateral curvature angle in the frontal plane indicates higher frontal spinal mobility (20). Porsnok et al. assessed only the presence of scoliosis and scoliosis angles in the frontal plane in their study on spinal alignment in children with UCP. No evaluation was made regarding lateral spinal mobility. Based on these findings, scoliosis was reported in 40% of children and they were found to be at risk for developing spinal deformities (14). In this study, the presence of scoliosis in children with UCP was not assessed. Scoliosis is considered a factor that affects lateral spinal mobility (20). We hypothesize that any scoliosis present in the children may have influenced the lateral spinal mobility angles.

There are very few studies in the literature that have assessed spinal alignment and mobility in children with UCP (6, 14). Spinal angulations in a study in which spinal posture and mobility of children with UCP were evaluated in the frontal plane with a SM were similar to the angle values in our study. However, in these studies, the lateral spinal mobility of the affected and unaffected sides in children with UCP has not been compared (6). Therefore, we are unable to relate the results of our study to those of other studies. Suh et al. reported differences in thoracolumbar kyphosis, lumbar lordosis, pelvic tilt, and sacral slope angles in children with CP compared to typically developing children in their study in which they examined spinopelvic mobility in the sagittal plane by radiography (21). In the literature, studies examining spinal posture and mobility in the

frontal plane in individuals with CP emphasized scoliosis and included interventions for scoliosis (13, 16). In studies conducted in the sagittal plane, the relationship between spinal angulations was emphasized. Accordingly, it was emphasized that there was a relationship between sacral inclination and lumbar lordosis and between lumbar lordosis and thoracic kyphosis (22).

It has been reported that frontal curvature angles are higher in children with UCP compared to their peers (14). In this study, the lateral spinal curvatures and lateral spinal mobility angles of the affected and unaffected sides were found to be similar. We hypothesize that this may be due to the involvement of not only the affected half of the body in children with UCP but also the "unaffected" half, which we typically consider as intact. In UCP, sensorimotor integration, bimanual coordination, and motor planning impairments affect both sides of the body (23). In the side where more severe sensory impairments are observed, the kinematics of the upper extremity may be affected, which in turn can influence spinal alignment and mobility (24). It has been suggested that the dominant side in UCP should be considered as the less affected side rather than the unaffected side (25).

There are several limitations in our study. The first limitation is that spinal mobility was assessed only in the frontal plane. The rotational movements of the spine on both sides of the body could not be evaluated. The SM provides information only on spinal angles and mobility in the sagittal and frontal planes (6). The

second limitation is that lower extremity anthropometric measurements, which could influence spinal alignment, were not taken in this study. Also, postural patterns specific to the children were not assessed. In future studies, the existing postural patterns of children with UCP could be identified, and spinal alignment and mobility could be evaluated and compared according to PGPP and AGPP patterns.

5.1. Conclusions

The lateral spinal mobility angles of the affected and unaffected sides in children with UCP are similar. We attribute these results to the fact that UCP is a condition that affects the entire body. Somatosensation is a parameter that influences functionality and posture in children with UCP. Including the somatosensory system in the assessment protocols of these children may help identify the underlying causes of postural disorders (26). In the rehabilitation of children with CP, healthcare professionals should focus on developing: (A) Proper spinal curvature (kyphosis or lordosis) and a neutral pelvic position in the sagittal plane, as well as the symmetry of pelvic, trunk, and shoulder girdle orientation in the coronal plane; (B) motor control of pelvic rotation, hip abduction, knee flexion, and ankle dorsiflexion (27). Additionally, attention should be paid to preventing the development of asymmetric posture during maturation (7).

Acknowledgements

We would like to thanks to the children and their families who participated the study. There was no funding resource in this study.

Footnotes

Authors' Contribution: Study concept and design: S. A. T. and T. Ç.; Acquisition of data: S. A. T.; Analysis and interpretation of data: S. A. T. and T. Ç.; Drafting of the manuscript: S. A. T. and T. Ç.; Critical revision of the manuscript for important intellectual content: S. A. T. and T. Ç.; Statistical analysis: T. Ç.; Administrative, technical, and material support: S. A. T. and T. Ç.; Study supervision: S. A. T. and T. Ç.

Conflict of Interests Statement: The authors declare no conflict of interests.

Data Availability: Data can be shared after the article is published by asking the corresponding author but

cannot be shared publicly, because participants were not notified that the data would be shared publicly.

Ethical Approval: Ethical approval was received from the Clinical Research Ethics Committee of Bolu Abant İzzet Baysal University (Decision number: 2023/211, date: 19.09.2023).

Funding/Support: The present study received no funding/support.

Informed Consent: Written informed consent was taken from the children and their parents.

References

- Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, et al. A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl.* 2007;109:8-14. [PubMed ID: 17370477].
- Johnson A. Cerebral palsy: epidemiology and causal pathways. *Arch Dis Child.* 2000;83(3):279A. [PubMed ID: 10952658]. [PubMed Central ID: PMC1718461]. <https://doi.org/10.1136/adc.83.3.279a>.
- Cans C. Surveillance of cerebral palsy in Europe: a collaboration of cerebral palsy surveys and registers. *Developmental Med Child Neurol.* 2007;42(12):816-24. <https://doi.org/10.1111/j.1469-8749.2000.tb00695.x>.
- Kallem Seyyar G, Aras B, Aras O. Trunk control and functionality in children with spastic cerebral palsy. *Dev Neurorehabil.* 2019;22(2):120-5. [PubMed ID: 29652201]. <https://doi.org/10.1080/17518423.2018.1460879>.
- Heyrman I, Desloovere K, Molenaers G, Verheyden G, Klingels K, Monbaliu E, et al. Clinical characteristics of impaired trunk control in children with spastic cerebral palsy. *Res Dev Disabil.* 2013;34(1):327-34. [PubMed ID: 23000634]. <https://doi.org/10.1016/j.ridd.2012.08.015>.
- Tas SA, Cankaya T. Effects of structured training on spinal posture and selective motor control in children with unilateral spastic cerebral palsy. *Gait Posture.* 2024;109:22-7. [PubMed ID: 38244393]. <https://doi.org/10.1016/j.gaitpost.2024.01.007>.
- Porter D, Michael S, Kirkwood C. Is there a relationship between preferred posture and positioning in early life and the direction of subsequent asymmetrical postural deformity in non ambulant people with cerebral palsy? *Child Care Health Dev.* 2008;34(5):635-41. [PubMed ID: 18796054]. <https://doi.org/10.1111/j.1365-2214.2008.00852.x>.
- Dewar R, Love S, Johnston LM. Exercise interventions improve postural control in children with cerebral palsy: a systematic review. *Dev Med Child Neurol.* 2015;57(6):504-20. [PubMed ID: 25523410]. <https://doi.org/10.1111/dmcn.12660>.
- Ozer Kaya D, Kocak UZ, Emuk Y, Olgac Dundar N, Bozkaya Yilmaz S, Gençpinar P. The comparison of regional spinal curvatures and movements in sitting posture in ambulatory children with cerebral palsy having minimal-to-moderate functional limitations. *Gait Posture.* 2021;90:408-14. [PubMed ID: 34571351]. <https://doi.org/10.1016/j.gaitpost.2021.09.192>.
- Pierret J, Beyaert C, Vasa R, Rumilly E, Paysant J, Caudron S. Rehabilitation of Postural Control and Gait in Children with Cerebral Palsy: the Beneficial Effects of Trunk-Focused Postural Activities. *Dev Neurorehabil.* 2023;26(3):180-92. [PubMed ID: 36959769]. <https://doi.org/10.1080/17518423.2023.2193269>.
- Peeters LHC, de Groot IJM, Geurts ACH. Trunk involvement in performing upper extremity activities while seated in neurological

patients with a flaccid trunk - A review. *Gait Posture*. 2018;62:46-55. [PubMed ID: 29524797]. <https://doi.org/10.1016/j.gaitpost.2018.02.028>.

12. Sveistrup H, Schneiberg S, McKinley PA, McFadyen BJ, Levin MF. Head, arm and trunk coordination during reaching in children. *Exp Brain Res.* 2008;188(2):237-47. [PubMed ID: 18392615]. <https://doi.org/10.1007/s00221-008-1357-1>.
13. Hasler C, Brunner R, Grundshtain A, Ovadia D. Spine deformities in patients with cerebral palsy; the role of the pelvis. *J Child Orthop.* 2020;14(1):9-16. [PubMed ID: 32165976]. [PubMed Central ID: PMC7043121]. <https://doi.org/10.1302/1863-2548.14.190141>.
14. Porsnok D, Mutlu A, Livanelioglu A. Assessment of spinal alignment in children with unilateral cerebral palsy. *Clin Biomech (Bristol)*. 2022;100:105800. [PubMed ID: 36279632]. <https://doi.org/10.1016/j.clinbiomech.2022.105800>.
15. Young NL, Wright JG, Lam TP, Rajaratnam K, Stephens D, Wedge JH. Windswept Hip Deformity in Spastic Quadriplegic Cerebral Palsy. *Pediatric Physical Therapy.* 1998;10(3). <https://doi.org/10.1097/00001577-199801030-00002>.
16. Galvis S, Burton D, Barnds B, Anderson J, Schwend R, Price N, et al. The effect of scoliotic deformity on spine kinematics in adolescents. *Scoliosis Spinal Disord.* 2016;11:42. [PubMed ID: 27800560]. [PubMed Central ID: PMC5080732]. <https://doi.org/10.1186/s13013-016-0103-x>.
17. Bartlett D, Purdie B. Testing of the spinal alignment and range of motion measure: a discriminative measure of posture and flexibility for children with cerebral palsy. *Dev Med Child Neurol.* 2005;47(11):739-43. [PubMed ID: 16225736]. <https://doi.org/10.1017/S0012162205001556>.
18. Paleg G, Livingstone R. Evidence-informed clinical perspectives on postural management for hip health in children and adults with non-ambulant cerebral palsy. *J Pediatr Rehabil Med.* 2022;15(1):39-48. [PubMed ID: 35275575]. <https://doi.org/10.3233/PRM-220002>.
19. Domagalska-Szopa M, Szopa A. Postural pattern recognition in children with unilateral cerebral palsy. *Ther Clin Risk Manag.* 2014;10:113-20. [PubMed ID: 24600228]. [PubMed Central ID: PMC3933466]. <https://doi.org/10.2147/TCRM.S58186>.
20. Ucurum SG, Uzunlar H, Kirmizi M, Polat K, Ozdemir E, Sahin A, et al. Relationship of sagittal and frontal spinal curvatures and mobility with balance and respiratory functions in adolescent idiopathic scoliosis: Preliminary report. *Gait & Posture.* 2024;113:179-80. <https://doi.org/10.1016/j.gaitpost.2024.07.193>.
21. Suh SW, Suh DH, Kim JW, Park JH, Hong JY. Analysis of sagittal spinopelvic parameters in cerebral palsy. *Spine J.* 2013;13(8):882-8. [PubMed ID: 23541886]. <https://doi.org/10.1016/j.spinee.2013.02.011>.
22. Novikov VA, Umnov VV, Umnov DV, Zvozil AV, Zharkov DS, Mustafaeva AR, et al. Correlation between frontal X-ray parameters of the hip joint and sagittal vertebral-pelvic profile in patients with cerebral palsy. *Pediatric Traumatol, Orthopaed Reconstructive Surgery.* 2023;11(2):149-58. <https://doi.org/10.17816/ptors321909>.
23. Gordon AM, Bleyenheuft Y, Steenbergen B. Pathophysiology of impaired hand function in children with unilateral cerebral palsy. *Dev Med Child Neurol.* 2013;55 Suppl 4:32-7. [PubMed ID: 24237277]. <https://doi.org/10.1111/dmcn.12304>.
24. Perruchoud D, Murray MM, Lefebvre J, Ionta S. Focal dystonia and the Sensory-Motor Integrative Loop for Enacting (SMILE). *Front Hum Neurosci.* 2014;8:458. [PubMed ID: 24999327]. [PubMed Central ID: PMC4064702]. <https://doi.org/10.3389/fnhum.2014.00458>.
25. Burn MB, Gogola GR. Dexterity of the Less Affected Hand in Children With Hemiplegic Cerebral Palsy. *Hand (N Y).* 2022;17(6):1114-21. [PubMed ID: 33605176]. [PubMed Central ID: PMC9608304]. <https://doi.org/10.1177/1558944721990803>.
26. Jovellar-Isiegas P, Cuesta Garcia C, Jaen-Carrillo D, Palomo-Carrion R, Pena Alonso C, Roche-Seruendo LE. Somatosensation and motor performance in the less-affected and more-affected hand of unilateral cerebral palsy children: a cross-sectional study. *Disabil Rehabil.* 2023;45(21):3500-10. [PubMed ID: 36172643]. <https://doi.org/10.1080/09638288.2022.2127938>.
27. Szopa A, Domagalska-Szopa M, Siwiec A, Kwiecien-Czerwieniec I. Canonical correlation between body-posture deviations and gait disorders in children with cerebral palsy. *PLoS One.* 2020;15(6):e0234654. [PubMed ID: 32544177]. [PubMed Central ID: PMC7297316]. <https://doi.org/10.1371/journal.pone.0234654>.