



Increasing the Anaerobic Capacity Resulting from the Improvement of Buffering and Hypoxia Indicators in Healthy Men

Samira Nasiri  ¹, Ebrahim Banitalebi  ^{1,*}, Mohammad Faramarzi  ²

¹Department of Sports Sciences, Shahrekord University, Shahrekord, Iran

²Department of Exercise Physiology, Faculty of Sports Sciences, University of Isfahan, Isfahan, Iran

*Corresponding Author: Department of Sports Sciences, Shahrekord University, Shahrekord, Iran. Email: banitalebi@sku.ac.ir

Received: 29 September, 2024; Revised: 30 October, 2024; Accepted: 5 November, 2024

Abstract

Background: Increasing anaerobic capacity is a crucial factor in enhancing performance in strength sports.

Objectives: This study investigated the effect of 10 weeks of weightlifting training with increased respiratory dead space on buffering capacity and anaerobic capacity in healthy men.

Methods: Eighteen male weightlifters (mean age: 28.2 ± 3.02 years, BMI: 24.27 ± 1.34 kg/m²) were selected through convenience sampling and randomly assigned to two training groups: With and without masks. Both groups performed selected weightlifting exercises three times per week for 10 weeks at 80% of their one-repetition maximum. The mask group trained using a mask and tube to increase respiratory dead space throughout the training period. The Borg scale was used to measure perceived effort, and the Wingate test assessed anaerobic power and Fatigue Index (FI). Blood samples were taken to measure lactate, CO₂, and hypoxia-inducible factor 1-alpha (HIF-1α) levels. Data analysis was conducted using ANOVA with repeated measures ($P \leq 0.05$).

Results: Ten weeks of weightlifting training with a mask led to a significant increase in bicarbonate levels ($P = 0.029$), lactate ($P \leq 0.001$), FI ($P \leq 0.001$), and rating of perceived exertion (RPE) ($P = 0.041$) compared to the group without a mask. There was no significant difference between the groups in HIF-1α levels ($P = 0.079$) or anaerobic power ($P = 0.534$). However, the percentage changes were greater in the mask group (anaerobic power = 5.9%, HIF-1α = 1.97%) compared to the group without a mask (anaerobic power = -0.9%, HIF-1α = 1.07%).

Conclusions: Using the strategy of increasing respiratory dead space during weightlifting training has proven beneficial for enhancing buffering capacity and anaerobic power.

Keywords: Bicarbonates, Carbon Dioxide, HIF-1alpha Protein, Respiratory Dead Space, Resistance Training

1. Background

Improving sports performance is of paramount importance. Coaches and athletes focus on physiological factors such as strength, VO₂max, and anaerobic capacity to achieve success (1). Weightlifters, in particular, require these physiological adaptations to lift heavy weights effectively. These athletes demonstrate exceptional explosive power compared to others (2, 3), which is a critical component for optimal performance (3). The positive impact of weight training on anaerobic capacity and the cardiovascular system has been well established (4). However, the search for innovative

methods to enhance power and reduce fatigue in weightlifters continues.

Research indicates that weightlifting exercises improve jumping ability and speed performance (5). However, there remains a gap in understanding how to further enhance performance and power in weightlifting. Blood flow restriction (BFR) training has been shown to improve anaerobic power, but it is often limited by movement restrictions, pain, and athletes' intolerance at higher intensities (6).

An alternative training method used in endurance exercises involves the use of an added respiratory dead space (ARDS) device. This device, which includes an additional corrugated tube of a specified length without

a valve to increase respiratory resistance, has shown potential for improving buffering capacity (7, 8). Studies on the ARDS method have reported increases in partial pressure of carbon dioxide (PCO_2) and decreases in blood pH (9). These physiological changes, along with increased bicarbonate (HCO_3^-) concentrations, may lead to delayed acidosis, improved buffering capacity, and enhanced energy generation through anaerobic metabolism (10).

In mammalian cells, oxygen homeostasis is regulated by the transcription factor hypoxia-inducible factor 1-alpha (HIF-1 α). Hypoxia-inducible factor 1-alpha regulates the expression of genes that enable cell adaptation to low oxygen tension, including genes involved in angiogenesis, glucose metabolism, and glucose transport. Research highlights a significant relationship between anaerobic power and HIF-1 α in athletes (11-13). Notably, scientists have observed that athletes focused on power and explosive performance possess a unique form of HIF-1 α , which enhances their performance (11, 12). Additionally, lactate (La^-), a byproduct of glycolysis and a key signaling molecule, has been shown to prevent the degradation of HIF-1 α (13).

However, to the best of our knowledge, the ARDS training method has only been studied in the context of aerobic exercises. Studies by Zaton et al. and Smolka et al. reported improved exercise performance following endurance training interventions using the ARDS method (9, 14). Similarly, in the study by Danek et al., increased performance was observed in sprint interval exercises (SIE) with varying volumes of ARDS, without significant changes in the rating of perceived exertion (RPE) or fatigue (15).

Overall, since the ARDS training method facilitates the development of anaerobic metabolism, the present study aimed to investigate whether applying this method to weightlifting exercises could capitalize on the increase in lactate (La^-) to enhance HIF-1 α and ultimately improve anaerobic power.

2. Objectives

Therefore, this study examined the effects of increased respiratory CO_2 during exercise on buffering capacity, HIF-1 α levels, anaerobic power, Fatigue Index (FI), and RPE over the course of 10 weeks of weightlifting training with added dead space.

3. Methods

Among 30 male weightlifters with a maximum of six months of weightlifting training experience (mean age:

28.2 ± 3.02 years), 20 men were selected through purposive convenience sampling. The sample size was determined using G*Power software and data from previous studies, identifying a range of 18 - 20 participants. The health status of the participants was assessed through a self-report form and the Physical Activity Readiness Questionnaire (PAR-Q). They were then randomly assigned, based on their body mass index (BMI between 20 and 25), into two groups: weightlifting with a mask (WARDS = 10 participants) and weightlifting without a mask (WT = 10 participants).

Inclusion criteria were no history of cardiovascular disease, diabetes, hypertension, respiratory diseases, non-smoking, and not following any diet or weight gain/loss programs. Exclusion criteria included taking any medication without prior notice, skeletal and respiratory diseases, and missing more than one-third of the training sessions. Written consent forms were obtained from all participants. Ultimately, data from 18 participants were evaluated (Table 1).

3.1. Technical Information

On the first day of the participants' arrival at the laboratory, measurements of height (using a Seca stadiometer, Germany), weight, and body composition (using an InBody 770 analyzer, South Korea) were conducted. These measurements were taken 48 hours after the protocol and under similar environmental conditions.

3.1.1. Anaerobic Power and Fatigue Index

The 30-second Wingate test was performed on a cycle ergometer (Monarch, model 894) with a load equivalent to 7.5% of the participant's body mass. All tests were conducted in the early morning following a light breakfast. Participants were instructed to start pedaling 10 seconds before the workload was applied and to continue with maximum effort for the duration of the 30-second test. Variables including peak anaerobic power and FI obtained from the test were recorded.

3.1.2. Rate of Perceived Exertion

The Borg scale (10-point) was used to measure RPE. Participants were verbally asked to report their perceived exertion after completing each exercise set, and the average RPE was calculated for that training day.

3.1.3. One Repetition Maximum

Participants selected the weight based on their initial estimate and performed the movement until failure. The amount of weight lifted and the number of

Table 1. General Characteristics of the Subjects ^a

Variables and Groups	Pre-test	Post-test	Percentage of Change	P		
				Intergroup	Between-Group	F
Age (y)						
WARDs	28.3 ± 2.17	-	-	-	-	-
WT	28.2 ± 3.83	-	-	-	-	-
Body height (cm)						
WARDs	174.5 ± 9.47	-	-	-	-	-
WT	179.7 ± 7.07	-	-	-	-	-
Body mass (kg)					0.714	0.457
WARDs	74.9 ± 8.86	76.4 ± 9.63	2.01	0.983		
WT	77.8 ± 8.47	79.5 ± 8.39	2.18	0.977		
BMI (kg/m²)					0.510	0.787
WARDs	24.5 ± 1.59	25.0 ± 1.62	1.95	0.883		
WT	24.0 ± 1.07	24.5 ± 1.08	2.20	0.844		
BFP (%)					0.942	0.130
WARDs	13.7 ± 1.11	13.4 ± 0.98	-2.46	0.929		
WT	13.5 ± 1.44	13.5 ± 1.22	-0.29	1.000		

Abbreviations: WARDs, group with mask; WT, group without mask.

^a Values are expressed as mean ± SD.

repetitions completed were used in the relevant formula to calculate the 1RM:

$$1RM = \frac{weight \ (kg)}{((1.0278) - (0.0278 \times repetition))}$$

3.2. Laboratory Method

Blood samples were collected to measure lactate (La^-), partial pressure of carbon dioxide (PCO_2), and bicarbonate (HCO_3^-) levels at six stages: Twenty-four hours and one hour before the start of the first training session, one hour before the start of the last training session, immediately after the first and last training sessions, and 48 hours after the last training session.

Additionally, a diagnostic kit was used to measure La^- and HIF-1 α levels at two stages: Before and after the training protocol.

3.3. Training Protocol

Participants underwent weightlifting training under the supervision of an experienced international coach for two weeks, attending three sessions per week to address and correct technical flaws. Following this preparatory phase, all participants engaged in the main weightlifting training program for eight weeks. Each session included two out of four primary weightlifting movements: Power snatch, squat snatch, power clean

and push jerk, and squat clean and split jerk. The training sessions progressively increased in frequency and intensity (Table 2). The primary distinction between the two groups was that the WARDs group used masks and tubes throughout the training sessions, except during the warm-up, cool-down, and a 10-minute rest period between exercises.

3.3.1. Dead Space Increasing Device

The device used consisted of an oxygen mask connected to a ventilator tube measuring 402 cm in length and 2.5 cm in diameter, creating 1000 mL of dead space (15).

3.4. Statistical Method

The test results were analyzed statistically by calculating the mean and standard deviation values. Normality and homogeneity of variance were assessed using the Shapiro-Wilk and Levene's tests. To examine the main effects and interactions between the group factor (WT vs. WARDs) and time factor (pre-training vs. post-training), a repeated measures ANOVA with a between-group factor was applied. If significant differences between groups were identified, Bonferroni post hoc tests were used for further analysis. All statistical calculations were conducted using SPSS software version 22, and charts were created in Excel. A significance level of $P \leq 0.05$ was considered for all tests.

Table 2. Weight Lifting Training Method ^a

Weeks	Activity	Frequency	Rep × Set (for Each Movements)	Intensity (RPE)	Intensity (%) iRM
1-2	Coordination-practice	3	5 × 6-8	14-16	80
3-4	Main exercise	3	5 × 6-8	14-16	80
5-6	Main exercise	4	5 × 6-8	14-16	80
7-10	Main exercise	5	5 × 6-8	14-16	80

Abbreviations: RPE, Rating of Perceived Exertion Based on the Borg Scale; iRM, one repetition maximum.

^aThe iRM of the subjects is measured every two weeks; and the training intensity percentage is calculated based on the new iRM.

Table 3. Indexes and Blood Gases ^a

Variables and Groups	24 Hours Before Training	Before the First Session	Immediately After the First Session	Before the Last Session	Immediately After the Last Session	48 Hours After the Last Session	P-Value ^b
PCO ₂ (mmHg)							0.031
WARDs	51.1 ± 3.6	52.6 ± 5.78	71 ± 11.5 ^c	56.6 ± 5.8	92.3 ± 13.8 ^c	51.3 ± 4.1	
WT	51.3 ± 4.9	52.6 ± 7.11	56.3 ± 7.18 ^c	52.3 ± 6.9	72.9 ± 17.5	51.2 ± 4.9	
HCO ₃ ⁻ (mg/L)							0.029
WARDs	30.0 ± 3.54	31.3 ± 2.07	35.8 ± 2.70 ^c	34.2 ± 2.47	38.6 ± 2.68 ^c	30.6 ± 2.56	
WT	30.2 ± 2.53	30.8 ± 1.96	31.6 ± 0.45	30.7 ± 2.94	31.7 ± 0.83	30.6 ± 2.96	
La ⁻ (mmol/L)							0.000
WARDs	1.01 ± 0.19	1.06 ± 0.19	4.23 ± 0.75 ^c	1.44 ± 0.29	5.87 ± 0.46 ^c	1.37 ± 0.19	
WT	0.97 ± 0.30	0.98 ± 0.33	2.24 ± 0.30	1.29 ± 0.28	2.23 ± 0.35	1.28 ± 0.38	

Abbreviations: WARDs, group with mask; WT, group without mask.

^aValues are expressed as mean ± SD.

^bP < 0.05 was considered statistically significant.

^cSignificant within-group changes.

4. Results

The results regarding the general characteristics of the subjects are presented in [Table 1](#). The findings indicate that after 10 weeks of training, the indices for weight (P = 0.714), BMI (P = 0.514), and body fat percentage (BFP, P = 0.942) did not exhibit significant between-group changes.

[Table 3](#) presents the results related to blood gases and La⁻ across six stages. The findings reveal that after 10 weeks of training, there was a significant inter-group change in the variables PCO₂ (P = 0.031), HCO₃⁻ (P = 0.029), and La⁻ (P ≤ 0.001). The intra-group analysis showed a significant difference in the variable PCO₂ in the WARDs group (after the first and last session) and in the WT group (after the first session). Additionally, significant differences were observed in the variables HCO₃⁻ and La⁻ in the WARDs group (after the first and last session).

The results for HIF-1α, anaerobic power, FI, and RPE are displayed in [Figures 1](#) and [2](#). The findings indicate that changes in HIF-1α concentration (P = 0.079) and anaerobic power (P = 0.534) did not exhibit significant intergroup differences. However, significant intergroup differences were observed in FI (P ≤ 0.001) and RPE (P = 0.041).

5. Discussion

The aim of this study was to investigate the effect of weightlifting exercises with ARDS on buffering capacity, HIF-1α, anaerobic power, FI, and RPE in weightlifters. The results indicated that using added dead space in weightlifting exercises enhanced buffering capacity, resulting in a 1.97% increase in HIF-1α levels, a 5.9% increase in anaerobic power, and a 23.2% reduction in FI in the WARDs group. Additionally, the RPE Index during training sessions decreased in the WARDs group, showing no significant difference between the two training groups on the final training day. Athletes and




Figure 1. Changes in hypoxia-inducible factor 1-alpha (HIF-1 α), anaerobic power and power drop after 10 weeks of weight lifting exercises with a mask (WARDS) and without a mask (WT). *: Significant intra-group difference ($P \leq 0.05$).

coaches can adopt this low-cost and efficient training method to improve buffering capacity, enhance anaerobic power, and reduce fatigue.

The adaptations resulting from increased mean partial pressure of CO_2 during ARDS training and the elevated HCO_3^- concentration from the carbonic anhydrase reaction (15-17) contribute to delayed acidosis and improved buffering capacity (18), which can further promote anaerobic metabolism (10). La^- , a potent signaling molecule, induces an increase in the HIF-1 α factor (13). In line with our study's findings, implementing the ARDS training method caused a significant rise in La^- levels in the WARDS group, consistent with the 1.97% increase in HIF-1 α levels in this group. Factors such as acute exercise, acidosis, oxidative stress, and heat have been shown to activate HIF-1 α expression independently of hypoxia (19). Additionally, La^- can increase HIF-1 α levels even in the presence of oxygen (13).

However, the results of our study contrast with those of Selfridge *et al.*, who reported that increased CO_2

responses reduce HIF transcription activity and that low pH conditions facilitate the lysosomal degradation of HIF- α protein (20).

The reason for this difference may lie in the fact that Selfridge *et al.* examined acute CO_2 exposure conditions, whereas in our study, participants trained under high CO_2 conditions for 10 weeks, and we measured the adaptations resulting from these conditions. Selfridge also suggested that the acidic pH conditions associated with high CO_2 exposure might deprive cells of nutrients, prompting a response in the form of lysosomal degradation of HIF- α protein (20). However, with the observed increase in La^- and HCO_3^- levels in the WARDS group of our study, it seems that enhanced buffering capacity facilitated better H^+ elimination, thereby preventing the reduction of HIF-1 α .

Research has demonstrated that RPE is a valuable tool for prescribing exercise intensity (21). Prescribing RPE-based training programs allows individuals to maintain exercise intensity within a predetermined RPE range, which is closely associated with objective




Figure 2. Changes in rating of perceived exertion (RPE) following 10 weeks of weight lifting exercises with a mask (WARDS) and without a mask (WT). *: Significant intra-group difference ($P \leq 0.05$). #: Significant between groups difference ($P \leq 0.05$).

physiological markers of intensity, such as heart rate, oxygen consumption, or blood La^- levels (22). Numerous studies have reported a strong correlation between blood La^- and RPE during exercise (23, 24). In our study, RPE levels showed significant differences between the two groups only in the initial training sessions, with no differences observed by the seventh and tenth weeks. This suggests that using a mask and tube during training was not perceived as more difficult by participants in the WARDS group. Furthermore, the WARDS group trained under higher CO_2 and La^- conditions with similar RPE levels and demonstrated superior results in buffering capacity, anaerobic power, and FI.

Our findings align with the results of studies by Danek et al. and López-Cabral et al. (15, 25). In Danek et al.'s study, 11 active individuals performed six 10-second repetitions with four minutes of active recovery over four laboratory sessions. The work done in the mask group (4.4%) and the average HCO_3^- concentration (6.7%) were higher, with no difference in RPE between groups (15). Similarly, López-Cabral's study stated that reductions in RPE were associated with increases in La^- due to metabolic adaptations during resistance training (25).

However, our results were inconsistent with findings by Miller et al. and Green et al. (22, 26). Miller et al.

reported that exogenous La^- intake during exercise did not affect RPE levels (26). Additionally, Green et al.'s study found a negative correlation between La^- concentration and RPE during cycling exercise, where RPE levels were lower when La^- levels were highest (22). In these studies, the exercise duration was much shorter than in our research, and the researchers focused on the immediate response of La^- to RPE. In contrast, our study examined the adaptations created over a prolonged training period, which are likely to influence the measured factors differently (27). Players with higher anaerobic power and lower FI are capable of superior performance in high-intensity activities (28). Researchers emphasize that elevated metabolic stress is a key factor in enhancing anaerobic power after training (6). Improved buffering capacity, through increased bicarbonate and its role in compensating for the energy demands of the anaerobic system (29), contributes to elevated La^- levels, as observed in this study. Alongside the rise in La^- levels, an increase in HIF-1 α levels was also noted. Several studies have reported a specific relationship between HIF-1 α levels and anaerobic power. It has been demonstrated that the distribution of HIF genotypes in strength and power athletes, such as weightlifters, sprinters, and short-distance swimmers, differs from the general population. Athletes with certain HIF alleles exhibit resistance to hypoxic conditions and possess enhanced glycolysis and

angiogenesis capabilities, making them particularly adept at power sports (12).

In studies conducted by Ahmetov et al. (11) and Cieszczyk et al. (12), a positive correlation was observed between the frequency of the HIF-1 α allele in weightlifters and their level of success. The frequency of this allele was notably higher in athletes compared to the control group. In the present study, findings from the WARDS group compared to the WT group revealed that an increase in HIF-1 α levels (1.97% vs. 1.07%) was accompanied by an increase in anaerobic power (5.9% vs. -0.9%) and a significant decrease in the FI (-23.2% vs. -1.6%).

Both the aforementioned studies involved a large number of athletes and non-athletes to address potential issues of population stratification and assessed HIF-1 α distribution through DNA testing and biopsies. While the present study had a smaller sample size, it supports the findings of these larger studies, demonstrating similar trends and validating the relationship between HIF-1 α levels, anaerobic power, and reduced fatigue.

5.1. Conclusions

Our study results demonstrated that employing the ARDS training strategy in weightlifting exercises enhanced buffering capacity and, through increased HIF-1 α levels, improved anaerobic power while reducing FI, without elevating RPE. Trainers and athletes involved in anaerobic sports activities who aim to enhance performance efficiently and reduce fatigue can benefit from incorporating this training method.

However, our study had limitations. The small sample size introduces potential ambiguities, preventing a robust confirmation of the hypothesized relationship between a specific type of HIF-1 α and anaerobic power. It is important to note that athletic performance is a multi-gene trait, and further exploration of other performance-related factors is warranted. Additionally, the findings of this study should be corroborated by future research involving longer durations, elite athletes, and female participants to provide a more comprehensive understanding of the impact of ARDS training on performance.

Footnotes

Authors' Contribution: Study concept and design: S. N.; Acquisition of data: S. N.; Analysis and interpretation of data: S. N. and E. B.; Drafting of the manuscript: S. N.; Critical revision of the manuscript for important intellectual content: S. N., E. B., and M. F.; Statistical

analysis: S. N. and E. B.; Administrative, technical, and material support: E. B. and M. F.; Study supervision: E. B. and M. F.

Clinical Trial Registration Code:
IRCT20200626055276N1.

Conflict of Interests Statement: The authors declared that they have no competing interests.

Data Availability: No new data were created or analyzed in this study. Data sharing does not apply to this article.

Ethical Approval: This study is approved under the ethical approval code of IR.SKU.REC.1401.016.

Funding/Support: This work is based upon research funded by Iran National Science Foundation (INSF) under project No. 4013749.

Informed Consent: Written consent forms were obtained from all participants. Ultimately, data from 18 participants were evaluated.

References

1. Ravasi A, Aminian T, Haghghi A. [Investigation and comparison of body composition, VO₂max strength and anaerobic power of elite weightlifters and non-athletes]. *Institute Humanities Cultural Stud.* 2004;21(21):5-17. FA.
2. Stone MH, Pierce KC, Sands WA, Stone ME. Weightlifting. *Strength Conditioning J.* 2006;28(1):50-66. <https://doi.org/10.1519/00126548-200602000-00010>.
3. Rogozkin V. Weightlifting and Power Events. In: Maughan RJ, editor. *Nutrition in Sport*. Blackwell, Oklahoma: Blackwell; 2008. p. 622-31. <https://doi.org/10.1002/9780470693766.ch47>.
4. Murugavel K, Balaji E. Impact of resistance training plyometric training and maximal power training on strength endurance and anaerobic power of team handball players. *Solid State Technol.* 2020;63(3):4259-71.
5. García-Valverde A, Manresa-Rocamora A, Hernández-Davó JL, Sabido R. Effect of weightlifting training on jumping ability, sprinting performance and squat strength: A systematic review and meta-analysis. *Int J Sports Sci Coaching.* 2021;17(4):917-39. <https://doi.org/10.1177/17479541211061695>.
6. Mostafa Farkhani B, Saqib Ju M, Hosseini Kakhk A, Hedayati M. [The effect of endurance training in productive speed along with blood flow limitation in rest periods on serum levels of VEGF and HIF-1 α and aerobic and anaerobic performance of male soccer players]. *J Practical Stud Biosciences Sport.* 2022;10(2):68-84. FA.
7. Szczepan S, Michalik K, Borkowski J, Zaton K. Effects of Swimming with Added Respiratory Dead Space on Cardiorespiratory Fitness and Lipid Metabolism. *J Sports Sci Med.* 2020;19(1):95-101. [PubMed ID: 32132832]. [PubMed Central ID: PMC7039034].
8. Szczepan S, Danek N, Michalik K, Wroblewska Z, Zaton K. Influence of a Six-Week Swimming Training with Added Respiratory Dead Space on Respiratory Muscle Strength and Pulmonary Function in Recreational Swimmers. *Int J Environ Res Public Health.* 2020;17(16). [PubMed ID: 32784446]. [PubMed Central ID: PMC7459907]. <https://doi.org/10.3390/ijerph17165743>.

9. Smolka L, Borkowski J, Zaton M. The effect of additional dead space on respiratory exchange ratio and carbon dioxide production due to training. *J Sports Sci Med*. 2014;13(1):36.
10. Woorons X, Mollard P, Pichon A, Duvallet A, Richalet JP, Lamberto C. Effects of a 4-week training with voluntary hypoventilation carried out at low pulmonary volumes. *Respir Physiol Neurobiol*. 2008;160(2):123-30. [PubMed ID: 18160351]. <https://doi.org/10.1016/j.resp.2007.09.010>.
11. Ahmetov I, Hakimullina AM, Lyubaeva EV, Vinogradova OL, Rogozkin VA. Effect of HIF1A gene polymorphism on human muscle performance. *Bull Exp Biol Med*. 2008;146(3):351-3. [PubMed ID: 19240858]. <https://doi.org/10.1007/s10517-008-0291-3>.
12. Cięszczyk P, Eider J, Arczewska A, Ostaniek M, Leońska-Duniec A, Sawczyn S, et al. The Hif1a Gene Pro582ser Polymorphism in Polish Power-Oriented Athletes. *Biology of Sport*. 2011;28(2):111-4. <https://doi.org/10.5604/945117>.
13. Kozlov AM, Lone A, Betts DH, Cumming RC. Lactate preconditioning promotes a HIF-1alpha-mediated metabolic shift from OXPHOS to glycolysis in normal human diploid fibroblasts. *Sci Rep*. 2020;10(1):8388. [PubMed ID: 32433492]. [PubMed Central ID: PMC7239882]. <https://doi.org/10.1038/s41598-020-65193-9>.
14. Zatoń MW, Hebisz RG, Hebisz P. The effect of training with additional respiratory dead space on haematological elements of blood. *Isokinetics Exercise Sci*. 2010;18(3):137-43. <https://doi.org/10.3233/ies-2010-0372>.
15. Danek N, Michalik K, Smolarek M, Zatoń M. Acute Effects of Using Added Respiratory Dead Space Volume in a Cycling Sprint Interval Exercise Protocol: A Cross-Over Study. *Int J Environm Res Public Health*. 2020;17(24):9485.
16. Pazoki AH, Choobineh S, Akbarnejad A. [The effect of six weeks combined training on plasma levels of chemerin, serum amyloid a and c-reactive protein and plasma lipid in obese male]. *J Arak Univ Med Sci*. 2016;19(1):1-11. FA.
17. Samuel R. A Graphical Tool for Arterial Blood Gas Interpretation using Standard Bicarbonate and Base Excess. *Indian J Med Biochem*. 2018;22(1):85-9. <https://doi.org/10.5005/jp-journals-10054-0061>.
18. Saunders B, Sale C, Harris RC, Sunderland C. Effect of sodium bicarbonate and Beta-alanine on repeated sprints during intermittent exercise performed in hypoxia. *Int J Sport Nutr Exerc Metab*. 2014;24(2):196-205. [PubMed ID: 24225816]. <https://doi.org/10.1123/ijsnem.2013-0102>.
19. Metzen E, Berchner-Pfannschmidt U, Stengel P, Marxsen JH, Stolze I, Klinger M, et al. Intracellular localisation of human HIF-1 alpha hydroxylases: implications for oxygen sensing. *J Cell Sci*. 2003;116(Pt 7):1319-26. [PubMed ID: 12615973]. <https://doi.org/10.1242/jcs.00318>.
20. Selfridge AC, Cavadas MA, Scholz CC, Campbell EL, Welch LC, Lecuona E, et al. Hypercapnia Suppresses the HIF-dependent Adaptive Response to Hypoxia. *J Biol Chem*. 2016;291(22):11800-8. [PubMed ID: 27044749]. [PubMed Central ID: PMC4882447]. <https://doi.org/10.1074/jbc.M116.713941>.
21. Stoudemire NM, Wideman L, Pass KA, McGinnes CL, Gaesser GA, Weltman A. The validity of regulating blood lactate concentration during running by ratings of perceived exertion. *Med Sci Sports Exerc*. 1996;28(4):490-5. [PubMed ID: 8778555]. <https://doi.org/10.1097/00005768-199604000-00014>.
22. Green JM, McLester JR, Crews TR, Wickwire PJ, Pritchett RC, Redden A. RPE-lactate dissociation during extended cycling. *Eur J Appl Physiol*. 2005;94(1-2):145-50. [PubMed ID: 15702340]. <https://doi.org/10.1007/s00421-004-1311-2>.
23. Hetzler RK, Seip RL, Boutilier SH, Pierce E, Snead D, Weltman A. Effect of exercise modality on ratings of perceived exertion at various lactate concentrations. *Med Sci Sports Exercise*. 1991;23(1). <https://doi.org/10.1249/00005768-199101000-00014>.
24. Noble BJ, Borg GA, Jacobs IRA, Ceci R, Kaiser P. A category-ratio perceived exertion scale. *Med Sci Sports Exercise*. 1983;15(6). <https://doi.org/10.1249/00005768-198315060-00015>.
25. López-Cabral JA, Rivera-Cisneros A, Rodríguez-Camacho H, Sánchez-González JM, Serna-Sánchez I, Trejo-Trejo M. Modification of fatigue indicators using citrulline malate for high performance endurance athletes. *Revista Mexicana de Patología Clínica y Medicina de Laboratorio*. 2012;59(4):194-201.
26. Miller BF, Fattor JA, Jacobs KA, Horning MA, Navazio F, Lindinger MI, et al. Lactate and glucose interactions during rest and exercise in men: effect of exogenous lactate infusion. *J Physiol*. 2002;544(3):963-75. [PubMed ID: 12411539]. [PubMed Central ID: PMC2290635]. <https://doi.org/10.1113/jphysiol.2002.027128>.
27. Ghoochani S, Riyahi Malayeri S, Daneshjo A. [Short-term effect of Citrulline Malate supplement on LDH and Lactate levels and Resistance Exercise Performance]. *J Military Med*. 2022;22(4):154-62. FA.
28. Haugen TA, Tonnessen E, Seiler S. Anaerobic performance testing of professional soccer players 1995-2010. *Int J Sports Physiol Perform*. 2013;8(2):148-56. [PubMed ID: 22868347]. <https://doi.org/10.1123/ijsspp.8.2.148>.
29. Minaseyan V, Eslami M, Sabaghe Langroudi M. The Effect of Sodium Bicarbonate Supplement on Anaerobic Power and Blood Lactate Level of Futsal Players. *J Sport Biosciences*. 2013;5(1):19-5.