



Plant-Origin Coumarin-Hemiterpene Ethers and Their Chemotaxonomic Significance

Saba Rahimi Bahoosh¹, Mahdi Mojarab ^{2,3,*}

¹ Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran

² Pharmaceutical Sciences Research Center, Research Institute for Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

³ Department of Pharmacognosy and Pharmaceutical Biotechnology, School of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding Author: Pharmaceutical Sciences Research Center, Research Institute for Health, Kermanshah University of Medical Sciences, Kermanshah, Iran. Email: mmojarab@kums.ac.ir

Received: 6 May, 2025; Revised: 6 June, 2025; Accepted: 23 June, 2025

Abstract

Context: A particular class of coumarins, known as coumarin-hemiterpene ethers (CHEs), consists of a coumarin nucleus and a prenyloxy portion. No detailed report on natural coumarin-hemiterpene ethers (NCHes) and their plant sources is available.

Evidence Acquisition: Electronic databases, including PubMed, EMBASE, Scopus, Google Scholar, and ScienceDirect, were searched for isolation reports of NCHes. All relevant papers published up to February 2025 were collected.

Results: The NCHes have been found in 15 plant families. The Rutaceae, Asteraceae, and Apiaceae families are the richest sources of NCHes. 7-isopentenylcoumarin is the most common NCHes. The non-glycosylated forms are the predominant types of plant-derived CHEs. The NCHes possess chemotaxonomic significance, as many have been identified exclusively in a single family. The highest number of exclusive NCHes was found in the Rutaceae family, while the Asteraceae family shows the highest diversity of unique substitution patterns.

Conclusions: The NCHes are specific coumarins that exhibit considerable chemotaxonomic potential.

Keywords: Coumarin, Prenyloxy, Isopentenyl, Rutaceae, Asteraceae, Apiaceae, Chemotaxonomy.

1. Context

The prenyloxy coumarins are a class of secondary metabolites with promising pharmacological activities. These compounds comprise structures in which a prenyl side chain is linked to the coumarin scaffold through an ethereal bond. In a type of prenyloxy coumarins, the prenyl side chain is an isopentenyl moiety (1) or, in other words, a hemiterpene moiety (C₅) (2). Although a hemiterpene is an isoprene group, several five-carbon compounds containing an isopentane skeleton, which can be saturated, unsaturated, repeatedly oxygenated, or methoxylated, are known as hemiterpenes (3). Szabo et al. first used the term “Coumarin-Hemiterpene Ethers” in 1985 for coumarins bearing a hemiterpene moiety by a C-O

linkage (4). To date, numerous natural coumarin-hemiterpene ethers (NCHes) have been reported with various biological activities such as antioxidant (5, 6), anti-inflammatory (7-9), antiviral (10, 11), antifungal (12, 13), cytotoxic, apoptotic (14), and antimutagenic activities (15). There are also reports on the synthesis and biological evaluation of coumarin-hemiterpene ethers (CHEs) that have not been found in nature (16). Although most NCHes are aglycones, a limited number of glycosylated NCHes have also been documented (17, 18). However, a comprehensive report on NCHes and their plant sources has not been available. Additionally, the significance of NCHes as chemotaxonomic markers has not been thoroughly investigated or discussed.

For the first time, in this review, we have categorized NCHes based on their structural features and plant

sources and have introduced their chemotaxonomic potential.

2. Evidence Acquisition

Electronic databases, including PubMed, EMBASE, Scopus, Google Scholar, and ScienceDirect, were searched for isolation reports of NCHes. All relevant papers published up to February 2025 were collected. Relevant articles written in English or those that had at least an English abstract were included in this study. Studies involving synthetic CHes and CHes isolated from non-plant sources were excluded from the current study. The search terms were as follows: Hemiterpene, isopentenyl, prenyl, oxy prenyl, dimethyl allyl, methyl butyl, methyl butyl, isoprene, and coumarin.

3. Results

3.1. Coumarin-Hemiterpene Ethers and Their Plant Sources

As previously mentioned, only a few glycosylated NCHes, such as 4'-O-(β -D-glucopyranosyl) desoxylacaryl and 5-O-(β -D-glucopyranosyl) lacaryl from the aerial parts of *Artemisia armeniaca* (Asteraceae) (17), and 6-O-[β -D-apiofuranosyl-(1-6)- β -D-glucopyranosyl]-prenyletin from the roots of *Prangos uloptera* (Apiaceae) (18), have been reported to date. The structures of these glycosylated NCHes are illustrated in Figure 1.

The non-glycosylated NCHes are found in 15 plant families, including Rutaceae, Asteraceae, Apiaceae, Cucurbitaceae, Amaranthaceae, Myrtaceae, Campanulaceae, Solanaceae, Polygalaceae, Convolvulaceae, Simaroubaceae, Thymelaeaceae, Theaceae, Scrophulariaceae, and Lamiaceae. Their structures, arranged based on the attachment site of the hemiterpene ether portion to the coumarin scaffold, and plant sources containing CHes, are depicted and presented in Figure 2.

Regarding to this table, there are no reports of 3-prenyl coumarins of plant origin. 4-prenyl coumarins and 6-prenyl coumarins have a very limited distributions. The findings suggest that 5-prenyl coumarins are more common, but they are restricted to the Rutaceae, Asteraceae, and Apiaceae families. 7-prenyl coumarins are the most prevalent NCHes, found in 13 families. Among all known NCHes, 7-

isopentenylcoumarin and its 6-methoxy derivative are highly prevalent.

A remarkable point about 8-prenyl coumarins is that two-thirds of these compounds have been isolated from the Asteraceae (genus *Artemisia*). Prenyl lacaryl and marianin A are NCHes that contain two hemiterpene ether moieties. These two NCHes are categorized in a distinct section named diprenyl coumarins. Figure 3 presents the occurrence of NCHes in different plant families.

The Rutaceae (53 compounds), Asteraceae (37 compounds), and Apiaceae (12 compounds) families are the richest sources of NCHes. However, a limited number of NCHes (10 compounds) have been reported from the other plant families. The various types of NCHes isolated from the three families – Rutaceae, Asteraceae, and Apiaceae – are displayed in Table 1.

The following sections provide detailed descriptions of NCHes in different plant families.

3.2. Natural Coumarin-Hemiterpene Ethers from Rutaceae

Almost half of the reports on the isolation of NCHes have resulted from phytochemical investigations on Rutaceae members. There are no reports of 4-prenyl coumarins, 6-prenyl coumarins, and diprenyl coumarins in Rutaceae. The genera *Citrus* and *Agathosma* had the highest occurrence of 5-prenyl coumarins and 7-prenyl coumarins, respectively.

3.3. Natural Coumarin-Hemiterpene Ethers from Asteraceae

Asteraceae is the sole plant family in which NCHes with all different substitution patterns (based on the attachment site of the hemiterpene ether portion) have been identified. The genera *Pterocaulon* and *Artemisia* had the highest number of 7-prenyl coumarins and 8-prenyl coumarins, respectively.

3.4. Natural Coumarin-Hemiterpene Ethers from Apiaceae

There are no reports of 4-prenyl, 6-prenyl, 8-prenyl, and diprenyl coumarins in Apiaceae.

3.5. Natural Coumarin-Hemiterpene Ethers from Other Plant Families

Several phytochemical findings have confirmed that other families, including Cucurbitaceae (109), Amaranthaceae (45, 46), Myrtaceae (47), Campanulaceae




Figure 1. The plant-derived glycosylated coumarin-hemiterpene ethers (CHEs)




Figure 3. The occurrence of coumarin-hemiterpene ethers (CHEs) in plant families

(48), Solanaceae (46), Polygalaceae (64), Convolvulaceae (6), Simaroubaceae (77) Thymelaeaceae (78), Theaceae (79), Scrophulariaceae (104), and Lamiaceae (110), also

contain NCHes. The NCHes from the other plant families are summarized in Table 2. As is evident, most of these families have provided 7-prenyloxy coumarins,

Table 1. The Coumarin-Hemiterpene Ether-Rich Plant Families

Variables	Families
Rutaceae (53 compounds)	
4-prenyloxy coumarins	-
5-prenyloxy coumarins (8 compounds)	<i>Citrus reticulata</i> (20), <i>C. aurantiifolia</i> (21), <i>C. medica</i> (22), <i>C. meyeri</i> (23), <i>C. limon</i> (15, 24), <i>Zanthoxylum nitidum</i> (27), <i>Toddalia asiatica</i> (28)
6-prenyloxy coumarins	-
7-prenyloxy coumarins (40 compounds)	<i>Agathosma collina</i> , <i>A. imbricata</i> , <i>A. martiana</i> , <i>A. mucronulata</i> , <i>A. mundii</i> , <i>A. recurvifolia</i> , <i>A. serpyllacea</i> , <i>A. spinosa</i> , <i>A. thymifolia</i> , <i>A. uncarpellata</i> , <i>A. puberula</i> (40), <i>Melicope lunu-ankenda</i> (59), <i>M. hayesi</i> (41), <i>M. semecarpifolia</i> (42), <i>M. vitiflora</i> (43), <i>M. borbonica</i> (73), <i>Diosma ramosissima</i> , <i>D. prama</i> , <i>D. recurva</i> (40), <i>D. acmaeophylla</i> (39), <i>Coleonema album</i> (37), <i>C. calycinum</i> (38), <i>C. aspalathoides</i> (38), <i>Clausena anisata</i> (60, 92), <i>C. indica</i> (61), <i>C. excavata</i> (98), <i>Citrus trifoliata</i> (93), <i>C. medica</i> (72), <i>C. limon</i> (24), <i>Choisya arizonica</i> , <i>Ch. mollis</i> (90), <i>Ch. ternata</i> (105), <i>Myrtopsis selligii</i> (94), <i>M. corymbosa</i> (11), <i>Haplophyllum obtusifolium</i> (71, 82), <i>H. ramosissimum</i> (83), <i>Galipea panamensis</i> (95), <i>G. trifoliata</i> (91), <i>Zanthoxylum schinifolium</i> (87), <i>Z. nitidum</i> (111), <i>Poncirus trifoliata</i> (62, 96, 97), <i>Phyllosma capensis</i> (102), <i>Triphasia trifolia</i> (10), <i>Eriostemon spicatus</i> (89), <i>Flindersia brayleyana</i> (107), <i>Euodia vitiflora</i> (35), <i>Asterolasia phebalioides</i> (36), <i>Ptaeroxylon obliquum</i> (65)
8-prenyloxy coumarins (5 compounds)	<i>C. calycinum</i> (38), <i>C. album</i> (37), <i>T. asiatica</i> (28), <i>Z. nitidum</i> (111)
Diprenyloxy coumarins	-
Asteraceae (37 compounds)	
4-prenyloxy coumarins (1 compound)	<i>Gerbera crocea</i> (19)
5-prenyloxy coumarins (3 compound)	<i>Artemisia laciniata</i> (29)
6-prenyloxy coumarins (2 compound)	<i>Carduus tenuiflorus</i> (33), <i>Pterocaulon polystachyum</i> (34)
7-prenyloxy coumarins (23 compound)	<i>P. polystachyum</i> (13, 34), <i>P. virgatum</i> (74), <i>P. alopecuroides</i> (13, 84), <i>P. balansae</i> (80), <i>Pt. redolens</i> (106) <i>Artemisia glauca</i> (51), <i>A. dracunculoides</i> (75), <i>Artemisia armeniaca</i> (86), <i>A. apiacea</i> (88), <i>A. laciniata</i> (29), <i>Haplopappus multifolius</i> (5, 81), <i>H. deserticola</i> (44), <i>Isocoma tenuisecta</i> , <i>L. acradenia</i> , <i>L. rushyi</i> (50), <i>Ozothamnus rosmarinifolius</i> (68), <i>O. lycopodioides</i> (99), <i>Tagetes lucida</i> (8, 9), <i>Helianthus tuberosus</i> (76), <i>Heterotheca inuloides</i> (49), <i>Viguiera gardneri</i> (66), <i>Flourensia thurifera</i> (67), <i>Melampodium divaricatum</i> (85), <i>Psidium dentata</i> (100), <i>Gochnatia argentina</i> (103), <i>Helichrysum plicatum</i> (101)
8-prenyloxy coumarins (7 compound)	<i>Artemisia tanacetifolia</i> (4), <i>A. caruifolia</i> (108), <i>A. armeniaca</i> (4, 86), <i>A. laciniata</i> (4, 29), <i>M. divaricatum</i> (85)
Diprenyloxy coumarins (1 compound)	<i>A. armeniaca</i> (4)
Apiaceae (12 compounds)	
4-prenyloxy coumarins	-
5-prenyloxy coumarins (5 compounds)	<i>Seseli sibiricum</i> (30, 31), <i>S. rigidum</i> (32), <i>Angelica gigas</i> (25), <i>Notopterygium incisum</i> (26)
6-prenyloxy coumarins	-
7-prenyloxy coumarins (7 compounds)	<i>Heracleum mantegazzianum</i> (63), <i>H. dissectum</i> (53), <i>H. lanatum</i> (54), <i>Angelica gigas</i> (70), <i>A. archangelica</i> (52), <i>Notopterygium franchetii</i> (58), <i>N. incisum</i> (26, 58), <i>Ligusticum lucidum</i> (7), <i>Seseli libanotis</i> (35), <i>Bupleurum fruticosum</i> (69), <i>Pimpinella anisum</i> , <i>Anethum graveolens</i> (52), <i>Lomatium nevadense</i> (55), <i>Tordylium apulum</i> (56), <i>Zosima absinthifolia</i> (57), <i>Libanotis intermedia</i> (35)
8-prenyloxy coumarins	-
Diprenyloxy coumarins	-

particularly 7-isopentenylcoumarin (6, 45-48, 64, 77-79, 104), while two families, Cucurbitaceae and Lamiaceae, possess a 4,8-diprenyloxy coumarin and an 8-prenyloxy coumarin, respectively (109, 110).

3.6. Chemotaxonomic Significance of Natural Coumarin-Hemiterpene Ethers

The NCHes possess considerable chemotaxonomic significance, as a large number of them (71 compounds) have been identified exclusively in single plant families

to date. The Rutaceae family has the highest number of unique NCHes (40 compounds). Furthermore, more than three-fourths of the exclusive NCHes in the Rutaceae family (31 compounds) are genus-specific or species-specific. Two genera, *Citrus* and *Coleonema*, provide six exclusive NCHes, while the number of genus-specific NCHes in *Melicope*, *Galipea*, and *Myrtopsis* is three, two, and one, respectively. It is noteworthy that *Poncirus trifoliata*, *Flindersia brayleyana*, and *Eriostemon spicatus* possess six, three, and two species-specific

Table 2. The Other Plant Families Containing Coumarin-Hemiterpene Ethers

Families	4-prenyloxy Coumarins	5-prenyloxy Coumarins	6-prenyloxy Coumarins	7-prenyloxy Coumarins	8-prenyloxy Coumarins	Diprenyloxy Coumarins
Cucurbitaceae (<i>Cucumis bisexualis</i>)	-	-	-	-	-	Marianin A (109)
Amaranthaceae (<i>Amaranthus retroflexus</i> , <i>Spinacia oleracea</i>)	-	-	-	7-isopentenylcoumarin (45, 46)	-	-
Myrtaceae (<i>Melaleuca alternifolia</i>)	-	-	-	7-isopentenylcoumarin (47)	-	-
Campanulaceae (<i>Codonopsis pilosula</i>)	-	-	-	7-isopentenylcoumarin (48)	-	-
Solanaceae (<i>Lycium barbarum</i>)	-	-	-	7-isopentenylcoumarin (46)	-	-
Polygalaceae (<i>Polygala sabulosa</i>)	-	-	-	6-methoxy-7-isopentenylcoumarin (64)	-	-
Convolvulaceae (<i>Convolvulus trabutianus</i>)	-	-	-	6-methoxy-7-isopentenylcoumarin (6)	-	-
Simaroubaceae (<i>Ailanthus altissima</i>)	-	-	-	6-methoxy-7-isopentenylcoumarin (77), Altissimacoumarin N (77), Puberulin (77)	-	-
Thymelaeaceae (<i>Daphne oleoides</i>)	-	-	-	Virgatenol (78)	-	-
Theaceae (<i>Eurya chinensis</i>)	-	-	-	Haplopinol methyl ether, 6-demethyl obtusin (79)	-	-
Scrophulariaceae (<i>Verbascum thapsus</i>)	-	-	-	2',3'-dihydroxy puberulin (104)	-	-
Lamiaceae (<i>Gmelina arborea</i>)	-	-	-	-	Artanin (110)	-

NCHes, respectively. Additionally, each of the two species, *Clausena anisata* and *Haplophyllum obtusifolium*, contains one species-specific NCHE.

Nearly half of the exclusive NCHes in the Rutaceae family (18 compounds) are 8-C-prenylated 7-prenyloxy coumarins that were observed in eight genera, including Myrtopsis, Poncirus, Clausena, Triphasia, Citrus, Galipea, Eriostemon, and Choisya. Besides, three species-specific NCHes in *F. brayleyana* are the 8-C-prenylated 6-methoxy 7-prenyloxy coumarin derivatives. The structural pattern of 5-hydroxy 7-prenyloxy coumarin is restricted to the Poncirus and Clausena genera, while the pattern of 5-prenyloxy 7-hydroxy coumarin is characteristic of the genus Citrus and is specifically found in *Citrus reticulata*.

The occurrence of 23 exclusive NCHes has been reported in the Asteraceae family. The genus Artemisia contains 10 unique NCHes, among which four are species-specific to *A. laciniata*, and three are species-specific to *A. armeniaca*. Moreover, six exclusive NCHes belong to the genus Pterocaulon, of which two are species-specific to *Pterocaulon balansae*, and the other two are species-specific to *P. polystachyum*, while *P. virgatum* provides one species-specific NCHE. Additionally, *Melampodium divaricatum* possesses two species-specific NCHes, and one unique NCHE has been reported for each of the following species: *Gerbera crocea*, *Carduus tenuiflorus*, and *Haplopappus multifolius*.

Two other exclusive NCHes in the Asteraceae family have been found in more than one genus (Ozothamnus, Psiadia, Pterocaulon, and Helichrysum). The substitution pattern of both of them is 5-hydroxy 6-

methoxy 7-prenyloxy, which has not been observed in other plant families. Other exclusive NCHE substitution patterns observed in Asteraceae include 4-prenyloxy 5-methyl (found in the genus Gerbera), 6-prenyloxy 7-methoxy (found in the genera Carduus and Pterocaulon), 5,6-dimethoxy 7-prenyloxy (found in the genus Pterocaulon), 7-prenyloxy 8-hydroxy (found in the genera Artemisia and Melampodium), 5-hydroxy 7-methoxy 8-prenyloxy (found in the genus Artemisia), 7-hydroxy 8-prenyloxy (found in the genera Artemisia and Melampodium), and 5,8-diprenyloxy 7-methoxy (found in the genus Artemisia).

There are five NCHes restricted to the Apiaceae family. These compounds exhibit two exclusive substitution patterns, including 5-methoxy 7-prenyloxy (found in Heracleum) and 5-prenyloxy 7-methoxy 8-prenyl (found in Seseli). Besides, three NCHes are species-specific to *Seseli sibiricum*. One exclusive NCHE has been reported for each of the families Cucurbitaceae (*Cucumis bisexualis*), Theaceae (*Eurya chinensis*), and Simaroubaceae (*Ailanthus altissima*). The Cucurbitaceae family (*C. bisexualis*) provides a notable exclusive substitution pattern (4, 7-diprenyloxy 5-methyl).

In summary, the Rutaceae family possesses the highest number of exclusive NCHes, following eight substitution patterns. Half of these patterns are not observed in other plant families. The 7-prenyloxy 8-prenyl pattern is found with the greatest abundance across eight genera. The Asteraceae family exhibits the highest diversity of unique substitution patterns, mainly with a distribution limited to a single genus (4 patterns) or two genera (3 patterns). Among all plant

genera studied, *Artemisia* has the highest number of exclusive NCHes (10 compounds). The *Apiaceae* and *Cucurbitaceae* families exhibit five and one exclusive NCHes, respectively. All of these compounds also display specific substitution patterns that are observed exclusively in these two families. Moreover, one exclusive NCH has been reported from each family, *Theaceae* and *Simaroubaceae*. However, similar substitution patterns have been observed in NCHes from other families.

4. Conclusions

The CHEs are a class of coumarins in which a hemiterpene moiety (C₅) is attached to the coumarin scaffold by O-prenylation. The plant-derived CHEs were abundantly isolated from the *Rutaceae*, *Asteraceae*, and *Apiaceae* families. However, these compounds have also been reported in 12 other plant families. 7-isopentenylloxycoumarin is the most common NCH. Almost 80% of NCHes have been exclusively isolated from a single plant family. The highest number of exclusive NCHes was identified in the *Rutaceae* family, while the *Asteraceae* family exhibits the highest diversity of unique substitution patterns. Accordingly, NCHes can be considered valuable chemotaxonomic markers.

4.1. Study Limitations

The number of phytochemical studies in some plant families and genera was not sufficient to draw definitive conclusions about the chemotaxonomic significance of isolated NCHes.

4.2. Future Directions

The findings of this study may help researchers conduct phytochemical studies on the plant genera that are likely to be rich in this type of secondary metabolite. This, in turn, will increase our understanding of their chemotaxonomic significance.

Footnotes

Authors' Contribution: Study concept and design: M. M.; Acquisition of data: S. R.; Analysis and interpretation of data: M. M.; Drafting of the manuscript: S. R.; Critical

revision of the manuscript for important intellectual content: M. M.; Study supervision: M. M.

Conflict of Interests Statement: The authors declare no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Funding/Support: The present study received no funding/support.

References

- Curini M, Cravotto G, Epifano F, Giannone G. Chemistry and biological activity of natural and synthetic prenyloxycoumarins. *Curr Med Chem*. 2006;**13**(2):199-222. [PubMed ID: 16472213]. <https://doi.org/10.2174/092986706775197890>.
- Tetali SD. Terpenes and isoprenoids: a wealth of compounds for global use. *Planta*. 2019;**249**(1):1-8. [PubMed ID: 30467631]. <https://doi.org/10.1007/s00425-018-3056-x>.
- Farozi A. *Study on Chemistry of Some Medicinally Important Plants of Jammu and Kashmir [Dissertation]*. Srinagar, India: National Institute of Technology Srinagar; 2017.
- Szabó G, Greger H, Hofer O. Coumarin-hemiterpene ethers from *Artemisia* species. *Phytochem*. 1985;**24**(3):537-41. [https://doi.org/10.1016/s0031-9422\(00\)80763-3](https://doi.org/10.1016/s0031-9422(00)80763-3).
- Torres R, Faini F, Modak B, Urbina F, Labbe C, Guerrero J. Antioxidant activity of coumarins and flavonols from the resinous exudate of *Haplopappus multifolius*. *Phytochem*. 2006;**67**(10):984-7. [PubMed ID: 16684545]. <https://doi.org/10.1016/j.phytochem.2006.03.016>.
- Belaouira R, Marchioni E, Benayache F, Benayache S. On-line screening and identification of polyphenolic antioxidant compounds of *Convolvulus trabutianus*. *Nat Prod Res*. 2020;**34**(10):1490-3. [PubMed ID: 30518259]. <https://doi.org/10.1080/14786419.2018.1510398>.
- Menghini I, Epifano F, Genovese S, Marcotullio MC, Sosa S, Tubaro A. Antiinflammatory activity of coumarins from *Ligusticum lucidum* Mill. subsp. *cuneifolium* (Guss.) Tammara (*Apiaceae*). *Phytother Res*. 2010;**24**(11):1697-9. [PubMed ID: 21031631]. <https://doi.org/10.1002/ptr.3170>.
- Sandra Liliana PD, Manases GC, Enrique JF, Ruben RR, Cinthya BP, Belen MG, et al. Isolation, chemical characterization, and anti-inflammatory activity of coumarins, flavonoids, and terpenes from *Tagetes lucida*. *Nat Prod Res*. 2022;**36**(18):4751-6. [PubMed ID: 34789033]. <https://doi.org/10.1080/14786419.2021.2005592>.
- Nayeli MB, Maribel HR, Enrique JF, Rafael BP, Margarita AF, Macrina FM, et al. Anti-inflammatory activity of coumarins isolated from *Tagetes lucida* Cav. *Nat Prod Res*. 2020;**34**(22):3244-8. [PubMed ID: 30618303]. <https://doi.org/10.1080/14786419.2018.1553172>.
- Likhitwitayawuid K, Supudompol B, Sritularak B, Lipipun V, Rapp K, Schinazi RF. Phenolics with Anti-HSV and Anti-HIV Activities from *Artocarpus gomezianus*, *Mallotus pallidus*, and *Triphasia trifolia*. *Pharm Biol*. 2008;**43**(8):651-7. <https://doi.org/10.1080/13880200500383058>.

11. Coulerie P, Maciuk A, Lebouvier N, Hnawia E, Guillemot J, Canard B, et al. Phytochemical study of myrtopsis corymbosa, perspectives for anti-dengue natural compound research. *Rec Nat Prod*. 2013;7(3):250.
12. Rahalison L, Benathan M, Monod M, Frenk E, Gupta MP, Solis PN, et al. Antifungal principles of *Baccharis pedunculata*. *Planta Med*. 1995;61(4):360-2. [PubMed ID: 7480184]. <https://doi.org/10.1055/s-2006-958101>.
13. Stein AC, Alvarez S, Avancini C, Zacchino S, von Poser G. Antifungal activity of some coumarins obtained from species of *Pterocaulon* (Asteraceae). *J Ethnopharmacol*. 2006;107(1):95-8. [PubMed ID: 16574360]. <https://doi.org/10.1016/j.jep.2006.02.009>.
14. Mojarrab M, Emami SA, Delazar A, Tayarani-Najaran Z. Cytotoxic Properties of Three Isolated Coumarin-hemiterpene Ether Derivatives from *Artemisia armeniaca* Lam. *Iran J Pharm Res*. 2017;16(1):221-9. [PubMed ID: 28496477]. [PubMed Central ID: PMC5423249].
15. Matsumoto T, Takahashi K, Kanayama S, Nakano Y, Imai H, Kibi M, et al. Structures of antimutagenic constituents in the peels of Citrus limon. *J Nat Med*. 2017;71(4):735-44. [PubMed ID: 28699128]. <https://doi.org/10.1007/s11418-017-1108-3>.
16. Iranshahi M, Jabbari A, Orafaie A, Mehri R, Zeraatkar S, Ahmadi T, et al. Synthesis and SAR studies of mono O-prenylated coumarins as potent 15-lipoxygenase inhibitors. *Eur J Med Chem*. 2012;57:134-42. [PubMed ID: 23047230]. <https://doi.org/10.1016/j.ejmech.2012.09.006>.
17. Mojarrab M, Delazar A, Hamburger M, Potterat O. New coumarin-hemiterpene ether glucosides and a structurally related phenylpropanoic acid derivative from *Artemisia armeniaca*. *Nat Prod Commun*. 2010;5(10):1619-22. [PubMed ID: 21121260].
18. Razavi SM, Nazemiyeh H, Delazar A, Hajiboland R, Rahman M, Gibbons S, et al. Coumarins from the roots of *Prangos uloptera*. *Phytochem Lett*. 2008;1(3):159-62. <https://doi.org/10.1016/j.phytol.2008.07.009>.
19. Kirsch G, Abdelwahab AB, Chaimbault P. Natural and Synthetic Coumarins with Effects on Inflammation. *Molecules*. 2016;21(10). [PubMed ID: 27706093]. [PubMed Central ID: PMC6273422]. <https://doi.org/10.3390/molecules21101322>.
20. Phetkul U, Phongpaichit S, Watanapokasin R, Mahabusarakam W. New depside from Citrus reticulata Blanco. *Nat Prod Res*. 2014;28(13):945-51. [PubMed ID: 24635118]. <https://doi.org/10.1080/14786419.2014.896010>.
21. Costa R, Russo M, De Grazia S, Grasso E, Dugo P, Mondello L. Thorough investigation of the oxygen heterocyclic fraction of lime (*Citrus aurantifolia* (Christm.) Swingle) juice. *J Sep Sci*. 2014;37(7):792-7. [PubMed ID: 24478239]. <https://doi.org/10.1002/jssc.201300986>.
22. Wang C, Huang J, Zhou Z, Xu P, Shi J, Yang Y, et al. Coumarins from Jinhua Finger Citron: Separation by Liquid-Liquid Chromatography and Potential Antitumor Activity. *Molecules*. 2023;28(19). [PubMed ID: 37836760]. [PubMed Central ID: PMC10574065]. <https://doi.org/10.3390/molecules28196917>.
23. Miyake Y, Ito C, Tokuda H, Suzuki N, Itoigawa M. Evaluation for Antitumor-promoting Activity of Meyerin and 7-Methoxy-5-prenyloxycoumarin in Meyer Lemon. *Food Sci Technol Res*. 2015;21(6):879-82. <https://doi.org/10.3136/fstr.21.879>.
24. Ziegler H, Spiteller G. Coumarins and psoralens from sicilian lemon oil (*Citrus limon* (L.) Burm. f.). *Flavour Fragr J*. 2006;7(3):129-39. <https://doi.org/10.1002/ffj.2730070307>.
25. Kang SY, Lee KY, Sung SH, Park MJ, Kim YC. Coumarins isolated from *Angelica gigas* inhibit acetylcholinesterase: structure-activity relationships. *J Nat Prod*. 2001;64(5):683-5. [PubMed ID: 11374978]. <https://doi.org/10.1021/np000441w>.
26. Wu SB, Zhao Y, Fan H, Hu YH, Hamann MT, Peng JN, et al. New guaiane sesquiterpenes and furanocoumarins from *Notopterygium incisum*. *Planta Med*. 2008;74(15):1812-7. [PubMed ID: 18991202]. [PubMed Central ID: PMC4883698]. <https://doi.org/10.1055/s-0028-1088326>.
27. Deng Y, Ding T, Deng L, Hao X, Mu S. Active constituents of *Zanthoxylum nitidum* from Yunnan Province against leukaemia cells in vitro. *BMC Chem*. 2021;15(1):44. [PubMed ID: 34301301]. [PubMed Central ID: PMC8305521]. <https://doi.org/10.1186/s13065-021-00771-0>.
28. Song SC, Ren BD, Wu XW, Xie YF, Cheng B, Wei Q, et al. Asiaticasides A-O, structurally intriguing coumarins from *Toddalia asiatica* with potential inflammatory inhibitory activity. *Phytochem*. 2024;221:114042. [PubMed ID: 38417721]. <https://doi.org/10.1016/j.phytochem.2024.114042>.
29. Hofer O, Szabó G, Greger H, Nikiforov A. Leaf coumarins from the *Artemisia laciniata* group. *Liebigs Ann Chem*. 2006;1986(12):2142-9. <https://doi.org/10.1002/ljac.198619861208>.
30. Banerjee SK, Gupta BD, Kumar R, Atal CK. New coumarins from the umbels of *Seseli sibiricum*. *Phytochem*. 1980;19(2):281-4. [https://doi.org/10.1016/s0031-9422\(00\)81974-3](https://doi.org/10.1016/s0031-9422(00)81974-3).
31. Kumar R, Gupta BD, Banerjee SK, Atal CK. New coumarins from *Seseli sibiricum*. *Phytochem*. 1978;17(12):2111-4. [https://doi.org/10.1016/s0031-9422\(00\)89291-2](https://doi.org/10.1016/s0031-9422(00)89291-2).
32. Stankov-Jovanovic VP, Ilic MD, Mitic VD, Mihajilov-Krstev TM, Simonovic SR, Nikolic Mandic SD, et al. Secondary metabolites of *Seseli rigidum*: Chemical composition plus antioxidant, antimicrobial and cholinesterase inhibition activity. *J Pharm Biomed Anal*. 2015;111:78-90. [PubMed ID: 25863020]. <https://doi.org/10.1016/j.jpba.2015.03.015>.
33. Cardona L, García B, José R P, Pérez J. 6-Prenyloxy-7-methoxycoumarin, a coumarin-hemiterpene ether from *Carduus tenuiflorus*. *Phytochem*. 1992;31(11):3989-91. [https://doi.org/10.1016/s0031-9422\(00\)97569-1](https://doi.org/10.1016/s0031-9422(00)97569-1).
34. Vera N, Bardon A, Catalan CA, Gedris TE, Herz W. New coumarins from *Pterocaulon polystachyum*. *Planta Med*. 2001;67(7):674-7. [PubMed ID: 11582550]. <https://doi.org/10.1055/s-2001-17365>.
35. Lassak E, Southwell I. The Coumarins From the Resin of *Euodia vitiflora*. *Aust J Chem*. 1972;25(11). <https://doi.org/10.1071/ch9722491>.
36. Sarker SD, Armstrong JA, Gray AI, Waterman PG. Coumarins from *Asterolasia phebalioides* (Rutaceae). *Biochem Syst Ecol*. 1994;22(4). [https://doi.org/10.1016/0305-1978\(94\)90036-1](https://doi.org/10.1016/0305-1978(94)90036-1).
37. Gray AI. New coumarins from *Coleonema album*. *Phytochem*. 1981;20(7):1711-3. [https://doi.org/10.1016/s0031-9422\(00\)98560-1](https://doi.org/10.1016/s0031-9422(00)98560-1).
38. Gray AI, Meegan CJ, O'Callaghan NB. Coumarins from two *Coleonema* species. *Phytochem*. 1986;26(1):257-60. [https://doi.org/10.1016/s0031-9422\(00\)81523-x](https://doi.org/10.1016/s0031-9422(00)81523-x).
39. Campbell WE, Provan GJ, Waterman PG. Simple coumarins from two populations of *Diosma acmaeophylla*. *Phytochem*. 1982;21(6):1457-8. [https://doi.org/10.1016/0031-9422\(82\)80171-4](https://doi.org/10.1016/0031-9422(82)80171-4).
40. Cambell WE, Majal T, Bean A. Coumarins of the rutoideae: Tribe Diosmeae. *Phytochem*. 1986;25(3):655-7. [https://doi.org/10.1016/0031-9422\(86\)88018-9](https://doi.org/10.1016/0031-9422(86)88018-9).

41. Latip J, Hartley TG, Waterman PG. Lignans and coumarins metabolites from *Melicopehayesii*. *Phytochem*. 1999;**51**(1):107-10. [https://doi.org/10.1016/s0031-9422\(98\)00720-1](https://doi.org/10.1016/s0031-9422(98)00720-1).
42. Chou HC, Chen JJ, Duh CY, Huang TF, Chen IS. Cytotoxic and anti-platelet aggregation constituents from the root wood of *Melicope semecarpifolia*. *Planta Med*. 2005;**71**(11):1078-81. [PubMed ID: 16320216]. <https://doi.org/10.1055/s-2005-871295>.
43. O'Donnell F, Ramachandran VN, Smyth TJ, Smyth WF, Brooks P. An investigation of bioactive phytochemicals in the leaves of *Melicope vitiflora* by electrospray ionisation ion trap mass spectrometry. *Anal Chim Acta*. 2009;**634**(1):115-20. [PubMed ID: 19154819]. <https://doi.org/10.1016/j.aca.2008.11.066>.
44. Zdero C, Bohlmann F, Niemeyer HM. Diterpenes and umbelliferone derivatives from *Haplopappus deserticola*. *Phytochem*. 1990;**29**(1):326-9. [https://doi.org/10.1016/0031-9422\(90\)89064-g](https://doi.org/10.1016/0031-9422(90)89064-g).
45. Fiorito S, Epifano F, Palmisano R, Genovese S, Taddeo VA. A re-investigation of the phytochemical composition of the edible herb *Amaranthus retroflexus* L. *J Pharm Biomed Anal*. 2017;**143**:183-7. [PubMed ID: 28605679]. <https://doi.org/10.1016/j.jpba.2017.05.051>.
46. Fiorito S, Preziuso F, Epifano F, Scotti L, Bucciarelli T, Taddeo VA, et al. Novel biologically active principles from spinach, goji and quinoa. *Food Chem*. 2019;**276**:262-5. [PubMed ID: 30409593]. <https://doi.org/10.1016/j.foodchem.2018.10.018>.
47. Scotti L, Genovese S, Bucciarelli T, Martini F, Epifano F, Fiorito S, et al. Analysis of biologically active oxyprenylated phenylpropanoids in Tea tree oil using selective solid-phase extraction with UHPLC-PDA detection. *J Pharm Biomed Anal*. 2018;**154**:174-9. [PubMed ID: 29549856]. <https://doi.org/10.1016/j.jpba.2018.03.004>.
48. Wang R, Su P, Zhang Z, Li B, Hu F, Gao K, et al. Triterpenoids, Steroids, and Other Constituents of the Roots of *Codonopsis pilosula*. *Chem Nat Compd*. 2021;**57**(6):1160-2. <https://doi.org/10.1007/s10600-021-03577-7>.
49. Delgado G, del Socorro Olivares M, Chavez MI, Ramirez-Apan T, Linares E, Bye R, et al. Antiinflammatory constituents from *Heterotheca inuloides*. *J Nat Prod*. 2001;**64**(7):861-4. [PubMed ID: 11473412]. <https://doi.org/10.1021/np0005107>.
50. Lee ST, Cook D, Davis TZ, Gardner DR, Johnson RL, Stonecipher CA. A survey of tremetone, dehydrotremetone, and structurally related compounds in *Isonoma* spp. (goldenbush) in the southwestern United States. *J Agric Food Chem*. 2015;**63**(3):872-9. [PubMed ID: 25554830]. <https://doi.org/10.1021/jf505423h>.
51. Suleimen YM, Jose RA, Suleimen RN, Ishmuratova MY, Toppet S, Dehaen W, et al. Isolation and In Silico SARS-CoV-2 Main Protease Inhibition Potential of Jusan Coumarin, a New Dicoumarin from *Artemisia glauca*. *Molecules*. 2022;**27**(7). [PubMed ID: 35408682]. [PubMed Central ID: PMC9000794]. <https://doi.org/10.3390/molecules27072281>.
52. Taddeo VA, Genovese S, Medina P, Palmisano R, Epifano F, Fiorito S. Quantification of biologically active O-prenylated and unprenylated phenylpropanoids in dill (*Anethum graveolens*), anise (*Pimpinella anisum*), and wild celery (*Angelica archangelica*). *J Pharm Biomed Anal*. 2017;**134**:319-24. [PubMed ID: 27916506]. <https://doi.org/10.1016/j.jpba.2016.11.048>.
53. Belenovskaya IM, Sinitskii VS, Tumbaa K. 7-Isopentenylloxycoumarin from *Heracleum dissectum*. *Chem Nat Compd*. 1977;**13**(4):478. <https://doi.org/10.1007/bf00565845>.
54. Nakata H, Sashida Y, Shimomura H. A new phenolic compound from *heracleum lanatum* Michx. var. *nippinicum* Hara. II. *Chem Pharm Bull*. 1982;**30**(12):4554-6. <https://doi.org/10.1248/cpb.30.4554>.
55. Beauchamp PS, Dev BC, Dev V, Midland SL, Sims JJ. California Lomatiums, Part VII. Analysis of the Essential Oils of *Lomatium nevadense* (Watson) J. Coulter et Rose var. *Parishii* (J. Coulter et Rose) Jepson. Isolation of trans-Dauc-8-en-11-ol, a New Sesquiterpene Alcohol and Naturally Occurring 2',3',3'-Trimethyl-2',3'-Dihydroangelicin. *J Essent Oil Res*. 2007;**19**(2):117-24. <https://doi.org/10.1080/10412905.2007.9699243>.
56. Kofinas C, Chinou L, Loukis A, Harvala C, Roussakis C, Maillard M, et al. Cytotoxic Coumarins from the Aerial Parts of *Tordylium apulum* and their Effects on a Non-Small-Cell Bronchial Carcinoma Cell Line. *Planta Med*. 2007;**64**(2):174-6. <https://doi.org/10.1055/s-2006-957398>.
57. Razavi SM, Imanzadeh G, Davari M. Coumarins from *Zosima absinthifolia* seeds, with allelopathic effects. *EurAsian J BioSci*. 2010;**4**(1):17-22. <https://doi.org/10.5053/ejobios.2010.4.0.3>.
58. Xu K, Jiang S, Zhou Y, Zhang Y, Xia B, Xu X, et al. Discrimination of the seeds of *Notopterygium incisum* and *Notopterygium franchetii* by validated HPLC-DAD-ESI-MS method and principal component analysis. *J Pharm Biomed Anal*. 2011;**56**(5):1089-93. [PubMed ID: 21856104]. <https://doi.org/10.1016/j.jpba.2011.07.034>.
59. Ito C, Matsui T, Tokuda H, Tan HT, Itoigawa M. Cancer chemopreventive constituents from *Melicope lunu-ankenda*. *Phytochem Lett*. 2017;**20**:172-6.
60. Ngadjui BT, Ayafor JF, Sondengam BL, Connolly JD. Coumarins from *Clausena anisata*. *Phytochem*. 1989;**28**(2):585-9.
61. Quan NV, Xuan TD, Anh LH, Tran H. Bio-guided isolation of prospective bioactive constituents from roots of *Clausena indica* (Dalzell) Oliv. *Molecules*. 2019;**24**(24):4442.
62. Feng T, Wang R, Cai X, Zheng Y, Luo X. Anti-human immunodeficiency virus-1 constituents of the bark of *Poncirus trifoliata*. *Chem Pharm Bull*. 2010;**58**(7):971-5.
63. Glowinski K, Mroczek T, Zabza A, Cierpicki T. Isolation And Structure Elucidation Of 5,7-disubstituted Simple Coumarins In The Fruits Of *Heracleum Mantegazzianum*. *Pharm Biol*. 2000;**38**(4):308-12. [PubMed ID: 21214482]. [https://doi.org/10.1076/1388-0209\(200009\)3841-AFT308](https://doi.org/10.1076/1388-0209(200009)3841-AFT308).
64. Pizzolatti MG, Luciano C, Monache FD. Styryl- and dihydrostyryl-2-pyrone derivatives from *Polygala sabulosa*. *Phytochem*. 2000;**55**(7):819-22. [PubMed ID: 11190403]. [https://doi.org/10.1016/s0031-9422\(00\)00301-0](https://doi.org/10.1016/s0031-9422(00)00301-0).
65. Agostinho D, Boudesocque L, Thery-Kone I, Debierre-Grockiego F, Gueiffier A, Enguehard-Gueiffier C, et al. A new meroterpenoid isolated from roots of *Ptaeroxylon obliquum* Radlk. *Phytochem Lett*. 2013;**6**(4):560-6.
66. Schorr K, Garcia-Pineros AJ, Siedle B, Merfort I, Da Costa FB. Guaianolides from *Viguiera gardneri* inhibit the transcription factor NF-kappaB. *Phytochem*. 2002;**60**(7):733-40. [PubMed ID: 12127591]. [https://doi.org/10.1016/s0031-9422\(02\)00197-8](https://doi.org/10.1016/s0031-9422(02)00197-8).
67. Faini F, Labbe C, Salgado I, Coll J. Chemistry, toxicity and antifeedant activity of the resin of *Flourensia thurifera*. *Biochem System Ecol*. 1997;**25**(3):189-93.
68. Wollenweber E, Fischer R, Dörr M, Irvine K, Pereira C, Stevens JF. Chemodiversity of exudate flavonoids in *Cassia* and *Ozothamnus* (Asteraceae, Gnaphalieae). *Z Naturforsch C J Biosci*. 2008;**63**(9-10):731-9.

69. Pistelli L, Bertoli A, Bilia AR, Morelli I. Minor constituents from *Bupleurum fruticosum* roots. *Phytochem*. 1996;**41**(6):1579-82.
70. Dschang Jong Jung PA, Huneck S. Gigasol and other coumarins from *Angelica gigas*. *Phytochem*. 1991;**30**(2):710-2.
71. Batirov ÉK, Matkarimov AD, Malikov VM, Yagudaev MR, Seitmuratov E. New coumarins of *Haplophyllum obtusifolium*. *Chem Nat Compd*. 1981;**16**(6):558-61.
72. Zheng Z, Hu H, Zeng L, Yang H, Yang T, Wang D, et al. Analysis of the characteristic compounds of *Citri Sarcodactylis Fructus* from different geographical origins. *Phytochem Anal*. 2022;**33**(1):72-82. [PubMed ID: 34114292]. <https://doi.org/10.1002/pca.3069>.
73. Simonsen HT, Adersen A, Bremner P, Heinrich M, Wagner Smitt U, Jaroszewski JW. Antifungal constituents of *Melicope borbonica*. *Phytother Res*. 2004;**18**(7):542-5. [PubMed ID: 15305313]. <https://doi.org/10.1002/ptr.1482>.
74. Debenedetti SL, Nadinic EL, Coussio JD, De Kimpe N, Boeykens M. Two 6, 7-dioxygenated coumarins from *Pterocaulon virgatum*. *Phytochem*. 1998;**48**(4):707-10.
75. Herz W, Bhat SV, Santhanam PS. Coumarins of *Artemisia dracunculoides* and 3', 6-dimethoxy-4', 5, 7-trihydroxyflavone in *A. arctica*. *Phytochem*. 1970;**9**(4):891-4.
76. Jantaharn P, Mongkolthananuruk W, Senawong T, Jogloy S, McCloskey S. Bioactive compounds from organic extracts of *Helianthus tuberosus* L. flowers. *Ind Crop Prod*. 2018;**119**:57-63.
77. Yan ZY, Lv TM, Wang YX, Shi SC, Chen JJ, Bin L, et al. Terpenylated coumarins from the root bark of *Ailanthus altissima* (Mill.) Swingle. *Phytochem*. 2020;**175**:112361. [PubMed ID: 32289598]. <https://doi.org/10.1016/j.phytochem.2020.112361>.
78. Riaz M, Malik A. Novel Coumarin Glycosides from *Daphne oleoides*. *Helv Chim Acta*. 2001;**84**(3):656-61. [https://doi.org/10.1002/1522-2675\(20010321\)84:3<656::Aid-hlca656>3.0.Co;2-d](https://doi.org/10.1002/1522-2675(20010321)84:3<656::Aid-hlca656>3.0.Co;2-d).
79. Song JL, Yuan Y, Tan HB, Huang RM, Liu HX, Xu ZF, et al. Anti-inflammatory and antimicrobial coumarins from the stems of *Eurya chinensis*. *J Asian Nat Prod Res*. 2017;**19**(3):222-8. [PubMed ID: 27299182]. <https://doi.org/10.1080/10286020.2016.1191474>.
80. Medeiros-Neves B, De Barros FMC, Von Poser GL, Ferreira Teixeira H. Quantification of coumarins in aqueous extract of *Pterocaulon balansae* (Asteraceae) and characterization of a new compound. *Molecules*. 2015;**20**(10):18083-94.
81. Chiang MT, Bittner M, Silva M, Mondaca A, Zemelman R, Sammes PG. A prenylated coumarin with antimicrobial activity from *Haplopappus multifolius*. *Phytochem*. 1982;**21**(11):2753-5.
82. Matkarimov AD, Batirov ÉK, Malikov VM, Seitmuratov E. Obtusin— a new coumarin from *Haplophyllum obtusifolium*. *Chem Nat Compd*. 1980;**16**:240-2.
83. Bessonova IA, Kurbanov D, Yunusov SY. Components of the leaves of *Haplophyllum ramosissimum*. *Chem Nat Compd*. 1990;**26**(2):234.
84. Vilegas W, Boralle N, Cabrera A, Bernardi AC, Pozetti GL, Arantes SF. Coumarins and a flavonoid from *Pterocaulon alopecuroides*. *Phytochem*. 1995;**38**(4):1017-9.
85. Borges-Del-Castillo J, Martinez-Martir AI, Rodriguez-Luis F, Rodriguez-Ubis JC, Vazquez-Bueno P. Isolation and synthesis of two coumarins from *Melampodium divaricatum*. *Phytochem*. 1984;**23**(4):859-61.
86. Mojarrab M, Delazar A, Moghadam SB, Nazemiyeh H, Nahar L, Kumarasamy Y, et al. Armenin and isoarmenin—two prenylated coumarins from the aerial parts of *Artemisia armeniaca*. *Chem Biodiver*. 2011;**8**(11):2097-103.
87. Li W, Sun YN, Yan XT, Yang SY, Kim E, Kang HK, et al. Coumarins and lignans from *Zanthoxylum schinifolium* and their anticancer activities. *J Agric Food Chem*. 2013;**61**(45):10730-40.
88. Shimomura H, SASHIDA YUTAKA, OHSHIMA YUKIO. The chemical components of *Artemisia apiacea* Hance. II. More coumarins from the flower heads. *Chem Pharm Bull*. 1980;**28**(1):347-8.
89. Rashid MA, Gray AI, Waterman PG, Armstrong JA. Coumarins from *Eriostemon spicatus*. *J Nat Prod*. 1992;**55**(5):685-7.
90. Dreyer DL, Pickering MV, Cohan P. Distribution of limonoids in the Rutaceae. *Phytochem*. 1972;**11**(2):705-13.
91. Wirasutisna KR, Gleye J, Moulis C, Stanislas E, Moretti C, Galipein, a coumarin from *Galipea trifoliata*. *Phytochem*. 1987;**26**(12):3372.
92. Ngadjui BT, Ayafor JF, Sondengam BL, Connolly JD. Prenylated Coumarins from the Leaves of *Clausena anisata*. *J Nat Prod*. 1989;**52**(2):243-7. <https://doi.org/10.1021/np50062a003>.
93. Kerekes D, Horvath A, Kusz N, Borcsa BL, Szemerédi N, Spengler G, et al. Coumarins, furocoumarins and limonoids of *Citrus trifoliata* and their effects on human colon adenocarcinoma cell lines. *Heliyon*. 2022;**8**(9). e10453. [PubMed ID: 36097483]. [PubMed Central ID: PMC9463373]. <https://doi.org/10.1016/j.heliyon.2022.e10453>.
94. Hifnawy MS, Vaquette J, Pousset JL, Cave A. [Neutral products of Myrtopsis sellingii: two new coumarins, myrsellin and myrsellinol]. *Planta Med*. 1977;**31**(2):156-62. FR. [PubMed ID: 854544]. <https://doi.org/10.1055/s-0028-1097509>.
95. Arango V, Robledo S, Séon-Méniel B, Figadere B, Cardona W, Sáez J, et al. Coumarins from *Galipea panamensis* and their activity against *Leishmania panamensis*. *J Nat Prod*. 2010;**73**(5):1012-4.
96. Guiotto A, Rodighiero P, Pastorini G, Celon E. Coumarins from unripe fruits of *Poncirus trifoliata*. *Phytochem*. 1977;**16**(8):1257-60.
97. Park JS, Chung B, Lee WH, Lee J, Suh Y, Oh DC, et al. Sortase A-Inhibitory Coumarins from the Folk Medicinal Plant *Poncirus trifoliata*. *J Nat Prod*. 2020;**83**(10):3004-11. [PubMed ID: 32996318]. <https://doi.org/10.1021/acs.jnatprod.0c00551>.
98. Seo C, Ahn E-K, Kang J-S, Lee J-H, Oh JS, Hong SS. Excavosides A and B, two new flavonoid glycosides from *Clausena excavata* Burm. f. (Rutaceae). *Phytochem Lett*. 2017;**20**:93-7.
99. Rumero A, Arriaga-Giner FJ, Wollenweber E. A new oxyprenyl coumarin and highly methylated flavones from the exudate of *Ozothamnus lycopodioides* (Asteraceae). *Z Naturforsch C J Biosci*. 2000;**55**(1-2):1-4. [PubMed ID: 10739091]. <https://doi.org/10.1515/znc-2000-1-202>.
100. Fortin H, Tomasi S, Jaccard P, Robin V, Boustie J. A prenyloxycoumarin from *Psiadia dentata*. *Chem Pharm Bull*. 2001;**49**(5):619-21. [PubMed ID: 11383617]. <https://doi.org/10.1248/cpb.49.619>.
101. Vujic B, Vidakovic V, Jadranin M, Novakovic I, Trifunovic S, Tesevic V, et al. Composition, Antioxidant Potential, and Antimicrobial Activity of *Helichrysum plicatum* DC. Various Extracts. *Plants*. 2020;**9**(3). [PubMed ID: 32155955]. [PubMed Central ID: PMC7154845]. <https://doi.org/10.3390/plants9030337>.
102. Campbell WE, Cragg GM. A New Coumarin from *Phyllosma capensis*. São Paulo, Brazil: CABI Digital Library; 1979.
103. Garcia EE, Guerreiro E. Sesquiterpene lactones from *Gochnatia palosanto* and coumarins from *G. argentina*. *Phytochem*. 1988;**27**(1):288-90.

104. Tamura S, Yoshihira K, Kawano T, Murakami N. Inhibitor for FcepsilonRI expression on mast cell from *Verbascum thapsus* L. *Bioorg Med Chem Lett*. 2018;**28**(20):3342-5. [PubMed ID: 30217416]. <https://doi.org/10.1016/j.bmcl.2018.09.007>.
105. Donald GR, de Carvalho PR, Fernandes PD, Boylan F. Antinociceptive activity of puberulin and choisyine from ethanol extract of *Choisya ternata* Kunth var. *Sundance*. *Biomed Pharmacother*. 2021;**141**:111926. [PubMed ID: 34323696]. <https://doi.org/10.1016/j.biopha.2021.111926>.
106. Ishikawa T, Kanlayavattanakul M, Ruangrunsi N, Watanabe T. Chemical Constituents of *Pterocaulon redolens*. *Heterocycles*. 2003;**61**:10.3987. <https://doi.org/10.3987/COM-03-S16>.
107. Moreira AS, Mathias L, Braz-Filho R, Schripsema J, Vieira IJ. Two new diprenylated coumarins from *Flindersia brayleyana*. *Nat Prod Lett*. 2002;**16**(4):291-5. [PubMed ID: 12168767]. <https://doi.org/10.1080/10575630290020668>.
108. Barua NC, Sharma RP, Madhusudanan K, Thyagarajan G, Herzt W. Coumarins in *Artemisia caruifolia*. *Phytochem*. 1980;**19**(10):2217-8.
109. Ma QG, Wei RR, Yang M, Huang XY, Wang F, Sang ZP, et al. Molecular Characterization and Bioactivity of Coumarin Derivatives from the Fruits of *Cucumis bisexualis*. *J Agric Food Chem*. 2018;**66**(22):5540-8. [PubMed ID: 29775541]. <https://doi.org/10.1021/acs.jafc.8b00976>.
110. Kubo M, Irimajiri R, Kawata M, Takahashi Y, Hayashi K, Matsuno M, et al. Prenylated-coumarins from *Gmelina arborea* and evaluation for neurotrophic activity. *Phytochem*. 2023;**213**:113721. [PubMed ID: 37279871]. <https://doi.org/10.1016/j.phytochem.2023.113721>.
111. Zuo GY, Wang CJ, Han J, Li YQ, Wang GC. Synergism of coumarins from the Chinese drug *Zanthoxylum nitidum* with antibacterial agents against methicillin-resistant *Staphylococcus aureus* (MRSA). *Phytomed*. 2016;**23**(14):1814-20. [PubMed ID: 27912884]. <https://doi.org/10.1016/j.phymed.2016.11.001>.

The plant-derived coumarin-hemiterpene ethers and their sources







4-prenyloxy coumarins			
	Name	Structure	plants
1	4-isopentenyl-5-methyl coumarin		<i>Gerbera crocea</i> (R) (19) (Asteraceae)
5-prenyloxy coumarins			
2	5-isopentenyl-7-hydroxy coumarin		<i>Citrus reticulata</i> (L) (20) (Rutaceae)
3	5-isopentenyl-7-methoxy coumarin		<i>Citrus aurantifolia</i> (F) (21), <i>Citrus medica</i> (F) (22), <i>Citrus meyeri</i> (F) (23), <i>Citrus limon</i> (O) (24) (Rutaceae) <i>Angelica gigas</i> (Un) (25) <i>Notopterygium incisum</i> (Rh) (26) (Apiaceae)
4	Wakayamalimonol A		<i>Citrus limon</i> (P) (15) (Rutaceae)
5	Wakayamalimonol B		<i>Citrus limon</i> (P) (15) (Rutaceae)
6	Wakayamalimonol C		<i>Citrus limon</i> (P) (15) (Rutaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (15, 19-26).

Continue of Figure 1








7	Wakayamalimonol D		<i>Citrus limon</i> (P) (15) (Rutaceae)
8	5-(2',3'-epoxy-3'-methylbutoxy)-7-methoxycoumarin		<i>Citrus limon</i> (O) (24) (Rutaceae)
9	Neoartanin		<i>Zanthoxylum nitidum</i> (L&R) (27), <i>Toddalia asiatica</i> (A) (28) (Rutaceae) <i>Artemisia laciniata</i> (L) (29) (Asteraceae)
10	Neoartanin epoxide		<i>Artemisia laciniata</i> (L) (29) (Asteraceae)
11	Neoartanindiol		<i>Artemisia laciniata</i> (L) (29) (Asteraceae)
12	Sesibiricin		<i>Seseli sibiricum</i> (R & U) (30, 31) (Apiaceae)
13	Sesibiricol		<i>Seseli sibiricum</i> (U) (30) (Apiaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (15, 24, 27-31).

Continue of Figure 1






14	Sesebrinol		<i>Seseli sibiricum</i> (R & U) (30, 31), <i>Seseli rigidum</i> (R) (32) (Apiaceae)
15	Sesebrin		<i>Seseli sibiricum</i> (R & U) (30, 31) (Apiaceae)
6-prenyloxy coumarins			
16	6-isopentenylxy-7-methoxy coumarin		<i>Carduus tenuiflorus</i> (A) (33) (Asteraceae)
17	Isovirgatenol		<i>Pterocaulon polystachyum</i> (A) (34) (Asteraceae)
7-prenyloxy coumarins			
18	7-isopentenylxycoumarin		<i>Euodia vitiflora</i> (Re) (35), <i>Asterolasia phebaloides</i> (A) (36), <i>Citrus limon</i> (O) (24), <i>Coleonema album</i> (A) (36), <i>Coleonema aspalathoides</i> (A) (37), <i>Coleonema calycinum</i> (A) (38), <i>Diosma acmaeophylla</i> (A) (39), <i>Diosma ramosissima</i> (A), <i>Diosma prama</i> (A) (40), <i>Melicope hayesii</i> (W) (41), <i>Melicope semecarpifolia</i> (R) (42), <i>Melicope vitiflora</i> (L) (43) (Rutaceae) <i>Haplopappus multifolius</i> (L & S) (5), <i>Haplopappus deserticola</i> (A) (44),

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (5, 24, 30-44).

Continue of Figure 1



			<i>Heterotheca imuloides</i> (A) (49), <i>Isocoma tenuisecta</i> (A), <i>Isocoma acradenia</i> (A), <i>Isocoma rusbyi</i> (A) (50), <i>Artemisia glauca</i> (A) (51), <i>Tagetes lucida</i> (A) (8, 9) (Asteraceae) <i>Pimpinella anisum</i> (Se), <i>Angelica archangelica</i> (R) (52), <i>Heracleum dissectum</i> (R) (53), <i>Heracleum lanatum</i> (R) (54), <i>Lomatium nevadense</i> (EO) (55), <i>Tordylium apulum</i> (A) (56) <i>Zosima absinthifolia</i> (Se) (57) <i>Notopterygium incisum</i> (Se), <i>Notopterygium franchetii</i> (Se) (58), <i>Libanotis intermedia</i> (F), <i>Seseli libanotis</i>, (R) (35) (Apiaceae) <i>Amaranthus retroflexus</i> (F) (45), <i>Spinacia oleracea</i> (L) (46) (Amaranthaceae) <i>Melaleuca alternifolia</i> (EO) (47) (Myrtaceae) <i>Codonopsis pilosula</i> (R) (48) (Campanulaceae) <i>Lycium barbarum</i> (F) (46) (Solanaceae)
19	7-(2',3'-dihydroxy-3'-methyl butoxy) coumarin		<i>Coleonema album</i> (A) (37) <i>Coleonema calycinum</i> (A) (38) (Rutaceae)
20	7-(2',3'-epoxy-3'-methyl butoxy) coumarin		<i>Coleonema album</i> (A) (37) <i>Coleonema calycinum</i> (A) (38) (Rutaceae) <i>Ligusticum lucidum</i> (A) (7) (Apiaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbls; Un, underground parts; W, wood; Wh, whole herb) (8, 9, 35, 45-58).

Continue of Figure 1










21	7-(3'-carboxy but-2'-enoxy) coumarin		<i>Melicope vitiflora</i> (L) (43), <i>Euodia vitiflora</i> (Re) (35) (Rutaceae)
22	7-(3'-methoxy carbonyl but-2'-enoxy) coumarin		<i>Coleonema calycinum</i> (A) (38) (Rutaceae)
23	7-(3'-carboxy butoxy) coumarin		<i>Melicope vitiflora</i> (L) (43), <i>Euodia vitiflora</i> (Re) (35) (Rutaceae)
24	7-(3'-methyl-4'-carboxybutanoxy) umbelliferone methyl ester		<i>Melicope hayesii</i> (W) (41) (Rutaceae)
25	7-(2'-hydroxy-3'-methylbut-3'-enoxy) coumarin		<i>Coleonema calycinum</i> (A) (38) (Rutaceae)
26	6-deoxyhaplopinol		<i>Melicope lumu-ankenda</i> (T) (59) (Rutaceae) <i>Haplopappus multifolius</i> (L & S) (5) (Asteraceae)
27	Melilunumarin A		<i>Melicope lumu-ankenda</i> (T) (59) (Rutaceae)
28	Melilunumarin B		<i>Melicope lumu-ankenda</i> (T) (59) (Rutaceae)
29	Anisocoumarin B		<i>Clausena anisata</i> (Sb & R) (60), <i>Clausena indica</i> (R) (61) <i>Poncirus trifoliata</i> (Sb) (62) (Rutaceae)
30	5-methoxy-7-isopentenylloxycoumarin		<i>Heracleum mantegazzianum</i> (F) (63) (Apiaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their source (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (5, 35, 38, 41, 43, 59-63).

Continue of Figure 1



31	Prenyletin		<i>Ptaeroxylon obliquum</i> (R) (65) (Rutaceae) <i>Viguiera gardneri</i> (A) (66) <i>Haplopappus multifolius</i> (L & S) (5) <i>Pterocaulon polystachyum</i> (A) (34) <i>Flourensia thurifera</i> (L & S) (67) <i>Ozothamnus rosmarinifolius</i> (A) (68) (Asteraceae) <i>Bupleurum fruticosum</i> (R) (69) <i>Angelica gigas</i> (A) (70) (Apiaceae)
32	6-methoxy-7-isopentenylcoumarin		<i>Agathosma collina</i> (A), <i>Agathosma imbricate</i> (A), <i>Agathosma martiana</i> (A), <i>Agathosma mucronulata</i> (A), <i>Agathosma mundii</i> (A), <i>Agathosma recurvifolia</i> (A), <i>Agathosma serpyllacea</i> (A), <i>Agathosma spinosa</i> (A), <i>Agathosma thymifolia</i> (A), <i>Agathosma uncarpellata</i> (A), <i>Diosma recurva</i> (A) (40), <i>Diosma acmaeophylla</i> (A) (39), <i>Haplophyllum obtusifolium</i> (E) (71), <i>Citrus medica</i> (F) (72), <i>Melicope borbonica</i> (L) (73) (Rutaceae) <i>Tagetes lucida</i> (A) (9), <i>Flourensia thurifera</i> (L & S) (67), <i>Pterocaulon polystachyum</i> (A) (13, 34), <i>Pterocaulon virgatum</i> (A) (74), <i>Artemisia dracunculoides</i> (A) (75), <i>Helianthus tuberosus</i> (F) (76), <i>Ozothamnus rosmarinifolius</i> (A) (68) <i>Haplopappus deserticola</i> (A) (44) (Asteraceae) <i>Bupleurum fruticosum</i> (R) (69) <i>Notopterygium incisum</i> (Rh) (26) (Apiaceae) <i>Polygala sabulosa</i> (Wh) (24) (Polygalaceae) <i>Convolvulus trabutianus</i> (A) (6) (Convolvulaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (5, 6, 9, 13, 26, 34, 39, 40, 44, 64-76).

Continue of Figure 1

			<i>Ailanthus altissima</i> (Rb) (77) (Simaroubaceae)
33	6-methoxy-7-(2',3'-epoxy-3'-methyl butoxy) coumarin		<i>Pterocaulon alopecuroides</i> (A) (13), <i>Pterocaulon balansae</i> (A) (80) (Asteraceae)
34	Virgatenol		<i>Pterocaulon polystachyum</i> (A) (34), <i>Pterocaulon virgatum</i> (A) (74), <i>Tagetes lucida</i> (A) (8) (Asteraceae) <i>Bupleurum fruticosum</i> (R) (69) (Apiaceae) <i>Daphne oleoides</i> (R) (78) (Thymelaeaceae)
35	Virgatol		<i>Pterocaulon virgatum</i> (A) (74) (Asteraceae)
36	Haplopinol		<i>Haplopappus multifolius</i> (L & S) (5, 81) (Asteraceae)
37	Haplopinol methyl ether		<i>Pterocaulon polystachyum</i> (A) (34) (Asteraceae) <i>Eurya chinensis</i> (S) (79) (Theaceae)
38	Obtusinin		<i>Haplophyllum obtusifolium</i> (E) (82), <i>Haplophyllum ramosissimum</i> (L) (83) (Rutaceae) <i>Pterocaulon polystachyum</i> (A) (34), <i>Pterocaulon balansae</i> (A) (80), <i>Pterocaulon virgatum</i> (A) (74), <i>Pterocaulon alopecuroides</i> (A) (84), (Asteraceae)
39	6-demethyl obtusinin		<i>Eurya chinensis</i> (S) (79) (Theaceae)
40	3'-deoxy obtusinin		<i>Pterocaulon polystachyum</i> (A) (34) (Asteraceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigae parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (5, 8, 13, 34, 69, 74, 77-84).

Continue of Figure 1









41	7-isopentenyl-8-hydroxy coumarin		<i>Melampodium divaricatum</i> (A) (85) (Asteraceae)
42	Isoarmenin		<i>Artemisia armeniaca</i> (A) (86) (Asteraceae)
43	Lacinartin		<i>Diosma acmaeophylla</i> (A) (39), <i>Zanthoxylum schinifolium</i> (S) (87), <i>Agathosma mucronulata</i> (A), <i>Agathosma mundii</i> (A) (40) (Rutaceae) <i>Artemisia tanacetifolia</i> (L) (4), <i>Artemisia laciniata</i> (L) (29), <i>Artemisia apiacea</i> (Fl) (88) (Asteraceae)
44	Lacinartin epoxide		<i>Artemisia laciniata</i> (L) (29) (Asteraceae)
45	Lacinartindiol		<i>Artemisia laciniata</i> (L) (29) (Asteraceae)
46	Ramosin		<i>Eriostemon spicatus</i> (A) (89), <i>Choisya arizonica</i> (A), <i>Choisya mollis</i> (A) (90), <i>Myrtopsis corymbosa</i> (B) (11), <i>Galipea trifoliata</i> (Sb & Rb) (91) (Rutaceae)
47	7-isopentenyl-8-(4'-hydroxy-3'-methylbut-2'-enyl) coumarin		<i>Eriostemon spicatus</i> (A) (89) (Rutaceae)
48	7-isopentenyl-8-(2'-hydroxy-3'-methylbut-3'-enyl) coumarin		<i>Eriostemon spicatus</i> (A) (89) (Rutaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (4, 11, 29, 39, 40, 85-91).

Continue of Figure 1








49	Anisocoumarin E		<i>Clausena anisata</i> (L) (92), <i>Triphasia trifolia</i> (L) (10) (Rutaceae)
50	Anisocoumarin F		<i>Clausena anisata</i> (L) (92), <i>Eriostemon spicatus</i> (A) (89) (Rutaceae)
51	Anisocoumarin G		<i>Clausena anisata</i> (L) (92) (Rutaceae)
52	Galipein		<i>Galipea trifoliata</i> (Sb & Rb) (91) (Rutaceae)
53	Myrsellin		<i>Myrtopsis corymbosa</i> (B) (11), <i>Citrus trifoliata</i> (P) (93), <i>Myrtopsis selligii</i> (A) (94) (Rutaceae)
54	7-((3,3-dimethyloxiran-2-yl)methoxy)-8-(3-(prop-1-en-2-yl)oxiran-2-yl) coumarin		<i>Galipea panamensis</i> (L) (95) (Rutaceae)
55	Poncimarin		<i>Poncirus trifoliata</i> (F) (96) (Rutaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (10, 11, 89, 91-96).

Continue of Figure 1

56	Isoponcimarín		<i>Triphasia trifolia</i> (L) (10), <i>Poncirus trifoliata</i> (F) (96) (Rutaceae)
57	Triphasiol		<i>Clausena anisata</i> (L) (92), <i>Triphasia trifolia</i> (L) (10), <i>Poncirus trifoliata</i> (F) (97), <i>Clausena excavata</i> (L & T) (98), <i>Citrus trifoliata</i> (P) (93) (Rutaceae)
58	O-methyl triphasiol A		<i>Poncirus trifoliata</i> (F) (97) (Rutaceae)
59	Triphasiolene A		<i>Poncirus trifoliata</i> (F) (97) (Rutaceae)
60	Ponciol		<i>Poncirus trifoliata</i> (F) (97) (Rutaceae)
61	O-methyl ponciol A		<i>Poncirus trifoliata</i> (F) (97) (Rutaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Fo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (10, 92, 93, 96-98).

Continue of Figure 1

62	O-methyl ponciol B		<i>Poncirus trifoliata</i> (F) (97) (Rutaceae)
63	Myrsellinol		<i>Myrtopsis selligii</i> (A) (94), <i>Myrtopsis corymbosa</i> (B) (11) (Rutaceae)
64	Isoobtusitin		<i>Ozothamnus lycopodioides</i> (Br) (99), <i>Psiadia dentata</i> (L) (100) (Asteraceae)
65	5-hydroxy-6-methoxy-7-(2',3'-dihydroxy-3'-methylbutoxy) coumarin		<i>Pterocaulon alopecuroides</i> (A) (84), <i>Helichrysum plicatum</i> (A) (101) (Asteraceae)
66	5,6-dimethoxy-7-(2',3'-dihydroxy-3'-methylbutoxy) coumarin		<i>Pterocaulon balansae</i> (A) (80) (Asteraceae)
67	5,6 dimethoxy-7-(2',3'-epoxy-3'-methylbutoxy) coumarin		<i>Pterocaulon balansae</i> (A) (80) (Asteraceae)
68	Capensin		<i>Phyllosma capensis</i> (A) (102), <i>Haplophyllum obtusifolium</i> (E) (71), <i>Agathosma puberula</i> (A), <i>Agathosma mucronulata</i> (A) (40) (Rutaceae) <i>Gochnatia argentina</i> (A) (103), <i>Flourensia thurifera</i> (L & S) (67) (Asteraceae) <i>Bupleurum fruticosum</i> (R) (69) (Apiaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (11, 40, 67, 69, 71, 80, 84, 94, 97, 99-103).

Continue of Figure 1








69	Obtusicin		<i>Haplophyllum obtusifolium</i> (E) (71) (Rutaceae)
70	Puberulin		<i>Choisya ternata</i> (L) (105), <i>Zanthoxylum schinifolium</i> (S) (87), <i>Agathosma martiana</i> (A), <i>Agathosma mucronulata</i> (A), <i>Agathosma puberula</i> (A), <i>Agathosma recurvifolia</i> (A), <i>Agathosma spinosa</i> (A) (40) (Rutaceae) <i>Pterocaulon redolens</i> (A) (106) (Asteraceae) <i>Ailanthus altissima</i> (Rb) (77) (Simaroubaceae)
71	2',3'-dihydroxy puberulin		<i>Helichrysum plicatum</i> (A) (101), <i>Pterocaulon redolens</i> (A) (106) (Asteraceae) <i>Verbascum thapsus</i> (104) (Scrophulariaceae)
72	Altissimacoumarin N		<i>Ailanthus altissima</i> (Rb) (77) (Simaroubaceae)
73	Brayleyanin		<i>Flindersia brayleyana</i> (Sb) (107) (Rutaceae)
74	6-methoxy-8-(3-methyl-2-oxobut-3-en-1-yl)-7-((3-methylbut-2-en-1-yl)oxy)-2H-chromen-2-one		<i>Flindersia brayleyana</i> (Sb) (107) (Rutaceae)
75	1-(6-methoxy-7-((3-methylbut-2-en-1-yl)oxy)-2-oxo-2H-chromen-8-yl)-3-methylbut-3-en-2-yl acetate		<i>Flindersia brayleyana</i> (Sb) (107) (Rutaceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (40, 71, 77, 87, 101, 104-107).

Continue of Figure 1








8-prenyloxy		coumarins	
76	7-hydroxy-8-isopentenylxy coumarin		<i>Melampodium divaricatum</i> (A) (85) (Asteraceae)
77	Armenin		<i>Artemisia armeniaca</i> (A) (86) (Asteraceae)
78	7-methoxy-8-isopentenylxy coumarin		<i>Coleonema calycinum</i> (A) (38) (Rutaceae) <i>Artemisia tanacetifolia</i> (L) (4), <i>Artemisia caruifolia</i> (A) (108) (Asteraceae)
79	7-methoxy-8-(2'-hydroxy-3'-methylbut-3'-enoxy) coumarin		<i>Coleonema calycinum</i> (A) (38) (Rutaceae)
80	7-methoxy-8-(2',3'-dihydroxy-3'-methyl butoxy) coumarin		<i>Coleonema calycinum</i> (A) (38) (Rutaceae)
81	7-methoxy-8-(2',3'-epoxy-3'-methyl butoxy) coumarin		<i>Coleonema calycinum</i> (A) (38), <i>Coleonema album</i> (A) (37) (Rutaceae)
82	Deoxylacarol		<i>Artemisia armeniaca</i> (A) (4, 86), <i>Artemisia laciniata</i> (L) (4, 29), <i>Artemisia tanacetifolia</i> (L) (4) (Asteraceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (4, 37, 38, 85, 86, 108).

Continue of Figure 1






83	Lacarol		<i>Artemisia armeniaca</i> (A) (4, 86), <i>Artemisia laciniata</i> (L) (4, 29) (Asteraceae)
84	Methylacarol		<i>Artemisia armeniaca</i> (L), <i>Artemisia tanacetifolia</i> (L) (4), <i>Artemisia laciniata</i> (L) (4, 29) (Asteraceae)
85	Artanin		<i>Toddalia asiatica</i> (A) (28) <i>Zanthoxylum nitidum</i> (R) (111) (Rutaceae) <i>Artemisia tanacetifolia</i> (L) (4) (Asteraceae) <i>Gmelina arborea</i> (S) (110) (Lamiaceae)
Diprenyloxy coumarin			
86	Marianin A		<i>Cucumis bisexualis</i> (F) (109) (Cucurbitaceae)
87	Prenyl lacarol		<i>Artemisia armeniaca</i> (L) (4) (Asteraceae)

Figure 2. The plant-derived coumarin-hemiterpene ethers (CHEs) and their sources (Abbreviations: A, aerial parts; B, bark; Br, branches; E, epigeal parts; Eo, essential oil; F, fruits; Fl, flowers; L, leaves; O, oil; P, peels; R, root; Rb, root barks; Re, resin; Rh, rhizome; S, stems; Sb, stem bark; Se, seed; T, twigs; U, umbels; Un, underground parts; W, wood; Wh, whole herb) (4, 28, 29, 86, 109-111).