

Diurnal Variation of Essential of the Oil Components of *Pycnocycla spinosa* Decne. ex Boiss

Gholamreza Asghari^{1*}; Gholamali Houshfar²; Zahra Mahmoudi¹; Matin Asghari³

¹Isfahan Pharmaceutical Sciences Research Center, Isfahan University of Medical Sciences, Isfahan, IR Iran

²Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, IR Iran

³Department of Agricultural Biotechnology, Tehran University, Tehran, IR Iran

*Corresponding author: Gholamreza Asghari, Isfahan Pharmaceutical Sciences Research Center, Isfahan University of Medical Sciences, Isfahan, IR Iran. Tel: +98-3117922644, E-mail: asghari@pharm.mui.ac.ir.

Received: May 15, 2013; Revised: August 28, 2013; Accepted: October 1, 2013

Background: *Pycnocycla spinosa* Decne. ex Boiss is an aromatic plant which showed relaxant effects on isolated ileum contractions and antidiarrheal activity. Thirty four components have been extracted from *P. spinosa* essential oil, of which several major constituents were found to show seasonal variation.

Objectives: The aim of this work is to evaluate the diurnal variation of its oil constituents during specific hours of the day.

Materials and Methods: The *Pycnocycla spinosa* samples were collected at different times of the day. The hydro-distilled aerial parts oils of collected *P. spinosa* were analyzed by GC and GC/MS.

Results: Fourteen monoterpenoid and nine sesquiterpenoid components were identified, of which the fluctuating constituents were *p*-cymene, *trans*- β -ocimene, β -citronellol, citronellyl pentanoate, geranyl isovalerate, α -humulene, caryophyllene oxide, α -cadinol, and α -eudesmol. The content of *p*-cymene in the essential oil in different daily times varied from 0.16 to 4.19%, and the geranyl isovalerate 7.75-23.99%.

Conclusions: Essential oils with different qualities can be obtained according to the harvest time of the plant in a day.

Keywords: Oils, Volatile; Plants; Gas Chromatography-Mass Spectrometry

1. Background

Pycnocycla spinosa Decne. ex Boiss is a variety of *Pycnocycla* which belongs to Umbelliferae. It is an aromatic plant distributed in central parts of Iran (1). It was reported that the essential oil and extract of *P. spinosa* have relaxant effects on isolated ileum contractions in relatively low concentration. The essential oil has more potent ileum relaxant effects than its extract (2, 3). In addition, comparative studies have been shown that the antidiarrheal activity of *P. spinosa* extract and its effect on small intestinal transition of charcoal meal are relatively similar to loperamide (4). The hydrodistilled aerial parts oil of *Pycnocycla spinosa* was analyzed and thirty-four components have been identified. The most abundant constituents identified in the essential oil are geranyl isopantanoate, caryophyllene oxide, β -eudesmol, citronellol, elemicin, *p*-cymene, citronellyl acetate, α -cadinol, nonadecane, sabinene, octanal, δ -cadinene, methyl eugenol, decanal, *trans*- β -ocimene, limonene, *trans*-caryophyllene and octadecane respectively, of which several major constituents were found to show seasonal variation (5, 6). There are no data concerning the

diurnal variation of these constituents during specific hours of the day. It is well known that the concentration of phytochemicals varies during the day. Diurnal variation in plant secondary chemical components has been observed, either related to light conditions (7, 8), water deficit (9) or more complex interactions (10). Diurnal cycles characterize the accumulation of many compounds in plants. This is an expression of changes in enzymatic activities in the daily courses; such variations are suggested to be probably caused by variations in temperature or light conditions. This paper reports the diurnal variation of the essential oil from aerial parts of the plant. There has been no research regarding this issue.

The amount of a compound is usually not constant throughout the life of a plant. The stage at which a plant is collected is, therefore, very important for maximizing the yield of the desired constituents. It is well known that the concentration of many metabolites in plants varies during the day and season (11-13). Diurnal fluctuations of the alkaloid concentration in the *Lupinus* spp have previ-

Implication for health policy makers/practice/research/medical education:

The quality of essential oil influences the success of aromatherapy. This article has been tried to evaluate the diurnal variation of oil constituents during specific hours of the day.

Copyright © 2014, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences; Published by DOCS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ously been explained (14). The yield and composition of the essential oil from the leaves of Eucalypt (*Eucalyptus nicholii*), Rosemary (*Rosmarinus officinalis* L.), White Cedar (*Thuja occidentalis* L.) and Lawson Cypress (*Chamaecyparis lawsoniana*) was reported to have diurnal variation (15). However, a limited published data is available for the role of the time harvesting on chemical composition of essential oil (16-24).

2. Objectives

The essential oils composition of *Pycnocycla spinosa* was found to show seasonal variation (6). There are no data concerning the diurnal variation of its oil constituents during specific hours of the day. In this study, the diurnal variation was investigated.

3. Materials and Methods

3.1. Plant Materials

Pycnocycla spinosa var. *spinosa* was collected from Isfahan University campus and was identified by the botanist Mr. Mehregan in department of biology at Isfahan University. A voucher specimen (A24) was authenticated and then deposited in the herbarium of Faculty of Pharmacy and Pharmaceutical Sciences, Isfahan, Iran. Samples of wild *P. spinosa* were collected in the middle of the flowering period. The samples were harvested every three-hour over a 12-hour period from 7:00 AM to 7:00 PM.

3.2. GC/MS Analysis

The hydrodistilled aerial parts oils of *P. spinosa* were analyzed by GC and GC/MS. Gas chromatography analysis was carried out using Perkin-Elmer 8500 gas chromatograph with flame ionization detector (FID) detector and a BP-1 capillary column (39 m × 0.25 mm; film thickness 0.25 μ m). The carrier gas was helium with a flow rate of 2 mL/min, the oven temperature for the first 4 minutes kept

at 60 °C and then increased at 4 °C/min until reached to the temperature of 280 °C, injector and detector temperatures were set at 280 °C. Confirmation of peak identity was effected by co-chromatography with standards and GC-MS. The mass spectra were recorded on a Hewlett Packard 6890 MS detector coupled with Hewlett Packard 6890 gas chromatograph equipped with HP-5MS capillary column (30 m × 0.25 mm; film thickness 0.25 μ m). The gas chromatography condition was as mentioned previously. Mass spectrometer condition was as follow: ionized potential 70 eV, source temperature at 200 °C. Identification was based on retention data and computer matching with the Wiley 275.L library as well as by comparison of electron-impact-mass spectra (EI-MS) with those relevant reference samples and the literature (25, 26).

4. Results

The hydrodistilled aerial parts oil of *Pycnocycla spinosa*, collected on different time of day, was analyzed and thirty-four components identified, of which several constituents were found with diurnal variation. The GC chromatogram of *P. spinosa* essential oil is presented in Figure 1 and diurnal variations of fluctuating monoterpenoid and sesquiterpenoid components are shown in Table 1.

As presented in Table 1, the fluctuating monoterpenoid constituents were *p*-cymene, *trans*- β -ocimene, β -citronellol, citronellyl pentanoate, and geranyl isovalerate. Also four fluctuating sesquiterpenoid constituents were α -humulene, caryophyllen oxide, α -cadinol, and α -eudesmol (see Table 1). Several studies on aromatic plants have shown that the essential oil composition may vary considerably throughout a year (27, 28). The results showed that the content of *p*-cymene in the essential oil in different daily times 0.16 - 4.19% varied, and of geranyl isovalerate did 7.75 - 23.99% (see Table 1). Seasonal variation has also been noticed for *p*-cymene in the essential oil of *Origanum onites* (29). Results of diurnal variation of β -citronellol and citronellyl pentanoate were correlated.




Figure 1. GC Chromatogram of *P. spinosa* Essential Oil

Table 1. The Diurnal Variation of Fluctuating Constituents in the Essential oil of *P. Spinosa*^a

Compounds	7:00 AM	10:00 AM	1:00 PM	4:00 PM	7:00 PM
P-cymene	2.94	0.16	1.40	1.56	4.19
t-β-ocimene	1.21	1.49	2.19	0.11	0.00
β-citronellol	7.26	5.01	5.67	8.11	1.55
Citronellyl Pentanoate	10.32	14.31	7.31	18.17	10.53
Geranyl isovalerate	17.91	19.56	7.75	11.61	23.99
α-humulene	1.32	1.11	0.00	0.46	0.57
Caryophyllen oxide	3.99	0.00	0.00	0.0	4.60
α-cadinol	2.86	0.59	0.00	2.12	3.01
α-eudesmol	5.72	6.46	3.46	4.11	5.12

^a % in Oil, average of triplicate

β -citronellol and citronellyl pentanoate reached to their lowest level at 7:00 and 1:00 PM. respectively, while both components reached to their highest levels at 4:00 PM. Similarly, it was reported that the percentage of citronellol, main components of rose oil, increased with delay in harvesting. Geraniol content of rose oil was maximum when the flowers were harvested at 10:00 AM, but after that there was significant reduction in its concentration up to 06:00 PM (24). As illustrated in Table 1, there is severing drop in citronellyl pentanoate content at 1:00 PM during day period. It may indicate some correlation between β -citronellol content and the level of its pentanoate ester. Results of diurnal variation of sesquiterpenes, α -humulene, caryophyllen oxide, α -cadinol, and α -eudesmol are presented in Table 1. Seasonal and diurnal variation has also been noticed for *Hymenaea courbaril*, *Copaifera officinalis*, and *Copaifera pubiflora*, which content of sesquiterpenes varies greatly during the year (30). α -cadinol showed the highest concentration at 7:00 PM. Also plant harvested early in the morning at 7:00 a.m. and late evening at 7:00 PM. provided essential oil with a high caryophyllen oxide content. Seasonal variation also was observed for this compounds (5). These variations were probably are related to the temperature or light variations. Plant cells are dependent on light for their growth and development. Light is also important for the production of metabolites by plant cells, and so also are irradiance, wavelength, and exposure time. As illustrated in Table 1, α -humulene reached to its highest level at 7:00 AM, while the lowest level was observed at 1:00 PM. In contrast, this compound was reported to be at highest level at 2:0 PM in *Rosa damascene* essential oil (31). It seems its fluctuation may not be related to the temperature or light but probably to genus and genetics of the plant. Caryophyllen oxide and α -cadinol reached their highest level at 7:00 PM. The highest content of eudesmol

was observed when the plant was harvested at 10:00 AM. Variation in the other mono- and sesquiterpenes due to time were not significant.

5. Discussion

The interesting point to note is that diurnal cycles characterize the accumulation of 5 monoterpenes and 4 sesquiterpenes of thirty-four components identified in the essential oils of *P. spinosa*. Such variation has been reported in many compounds of the plants (32, 33). It seems that this is an expression of changes in enzymatic activities; such variations are suggested to be probably caused by variations in temperature or light conditions in different courses of the day (34). Light effects on the accumulation of secondary metabolites may occur on three different levels; through direct control of product concentrations, influences on membrane permeability, and influence on enzymatic reactions. The latter may occur either by varying enzyme activities or altering the concentrations of the involved enzymes. In conclusion, variations in essential oil components of *P. spinosa* in response to diurnal changes may indicate that essential oil with different qualities can be obtained according to the harvest season and harvest time of the plant.

Acknowledgements

This research was supported by a grant from the research council of the Isfahan University of Medical Sciences.

Authors' Contribution

Gholamreza Asghari and Gholamali Houshfar carried out the design and coordinated the study and participated in most of the experiments. Matin Asghari contributed in data analysis and writing and finalizing the

manuscript. Zahra Mamoudi participated in most of the experiments and in manuscript preparation.

Financial Disclosure

The authors have no financial interests related to the materials of the manuscript.

Funding/Support

The work was financially supported by the research council of the Isfahan University of Medical Sciences, Isfahan, IR Iran.

References

1. Mozaffarian V. *A dictionary of Iranian plant name*. Tehran: Farhang Moaser; 1996.
2. Sadraei H, Asghari G, Naddafi A. Relaxant effect of essential oil and hydro-alcoholic extract of *Pycnocycla spinosa* Decne. ex Boiss. on ileum contractions. *Phytother Res*. 2003;17(6):645-9.
3. Sadraei H, Asghari G, Behzad S. Bioactivity-guided isolation of spasmolytic components of *Pycnocycla spinosa* Decne ex Boiss. *Res Pharm Sci*. 2011;6(2):81-6.
4. Sadraei H, Asghari G, Shams M. Antidiarrheal action of hydroalcoholic extract of *Pycnocycla spinosa* in comparison with loperamide and dicyclomine. *Iranian J Pharm Res*. 2011;10(4):835-41.
5. Mahmoudi Z. Composition of the essential oil of *Pycnocycla Spinosa* Decne. ex. Boiss. from Isfahan. *DARU J Pharm Sci*. 2001;9(3-4):28-9.
6. Asghari G, Houshfar G, Mahmoudi Z. Seasonal Variation of Mono- and Sesquiterpenes in the Essential Oil of *Pycnocycla Spinosa* Decne. ex Boiss. *Iranian J Pharm Res*. 2010;1:61-3.
7. Dickson RE. Diurnal changes in leaf chemical constituents and (14)C partitioning in cottonwood. *Tree Physiol*. 1987;3(2):157-71.
8. Badri DV, Loyola-Vargas VM, Broeckling CD, Vivanco JM. Root secretion of phytochemicals in *Arabidopsis* is predominantly not influenced by diurnal rhythms. *Mol Plant*. 2010;3(3):491-8.
9. Rosa EAS, Heaney RK, Rego FC, Fenwick GR. The variation of glucosinolate concentration during a single day in young plants of *Brassica oleracea* var *acephala* and *capitata*. *J Sci Food Agric*. 1994;66(4):457-63.
10. Itenov K, Molgaard P, Nyman U. Diurnal fluctuations of the alkaloid concentration in latex of poppy *Papaver somniferum* is due to day-night fluctuations of the latex water content. *Phytochem*. 1999;52(7):1229-34.
11. Brown PD, Tokuhisa JG, Reichelt M, Gershenson J. Variation of glucosinolate accumulation among different organs and developmental stages of *Arabidopsis thaliana*. *Phytochem*. 2003;62(3):471-81.
12. Zeeman SC, Smith SM, Smith AM. The diurnal metabolism of leaf starch. *Biochem J*. 2007;401(1):13-28.
13. Hemm MR, Rider SD, Ogas J, Murry DJ, Chapple C. Light induces phenylpropanoid metabolism in *Arabidopsis* roots. *Plant J*. 2004;38(5):765-78.
14. Birecka H, Zebrowska J. Diurnal changes in the alkaloid content in *Lupinus albus*, *Lupinus luteus*. *Bull Acad Polon Sci Ser Sci Biol*. 1960;8:339-41.
15. Hopfinger JA, Kumamoto J, Scora RW. Diurnal variation in the essential oils of Valencia orange leaves. *Am J Botany*. 1979;111:5.
16. Ramezani S, Ramezani F, Rasouli F, Ghasemi M, Fotokian MH. Diurnal variation of the essential oil of four medicinal plants spe-
17. cies in central region of Iran. *Res J Biol Sci*. 2009;4(1):103-6.
18. Chang X, Alderson PG, Wright Ch J. Variation in the Essential Oils in Different Leaves of Basil (*Ocimum basilicum* L.) at Day Time. *Open Horticulture J*. 2009;2:13-6.
19. de Vasconcelos Silva MG, Craveiro AA, Abreu Matos FJ, Machado MIL, Alencar JW. Chemical variation during daytime of constituents of the essential oil of *Ocimum gratissimum* leaves. *Fitoterapia*. 1999;70(1):32-4.
20. Angelopoulou D, Demetzos C, Perdetzoglou D. Diurnal and seasonal variation of the essential oil labdanes and clerodanes from *Cistus monspeliensis* L. leaves. *Biochem Sys Ecol*. 2002;30(3):189-203.
21. Yaldiriz G, Sekeroglu N, Ozguven M, Kirpik M. Seasonal and diurnal variability of essential oil and its components in *Origanum onites* L. grown in the ecological conditions of Cukurova. *Grasas Aceites*. 2005;56(4):254-8.
22. Gurbuz B, Ipek A, Basalma D, Sarihan EO, Sancak C, Ozcan S. Effects of diurnal variability on essential oil composition of sweet basil (*Ocimum basilicum* L.). *Asian Jo Chem*. 2006;18(1):285-8.
23. Ayan AK, Cirak C, Yanar O. Variations in total phenolics during ontogenetic, morphogenetic, and diurnal cycles in *Hypericum* species from Turkey. *J Plant Biol*. 2006;49(6):432-9.
24. Esmaelian Y, Galavi M, Ramroodi M, Mashhadi M, Boojar A. Diurnal variability of stigma compounds of saffron (*Crocus sativus* L.) at different harvest times. *Ann Biol Res*. 2012;3(3):1562-8.
25. Baj T, Ludwiczuk A, Sieniawska E, Skalicka-Wozniak K, Widelski J, Zieba K, et al. GC-MS analysis of essential oils from *Salvia officinalis* L.: comparison of extraction methods of the volatile components. *Acta Pol Pharm*. 2013;70(1):35-40.
26. Zheljazkov VD, Astatkine T, Jeliazkova E. Year-round Variations in Essential Oil Content and Composition of Male and Female Juniper. *Hort Sci*. 2013;48(7):883-6.
27. Kumar R, Sharma S, Sood S, Agnihotri VK, Singh B. Effect of diurnal variability and storage conditions on essential oil content and quality of damask rose (*Rosa damascena* Mill.) flowers in north western Himalayas. *Sci Hortic*. 2013;154:102-8.
28. Ozkan G, Baydar H, Erbas S. The influence of harvest time on essential oil composition, phenolic constituents and antioxidant properties of Turkish oregano (*Origanum onites* L.). *J Sci Food Agric*. 2010;90(2):205-9.
29. ElSohly HN, Croom EM, Kopycki WJ, Joshi AS, McChesney JD. Diurnal and seasonal effects on the taxane content of the clippings of certain *Taxus* cultivars. *Phytochem Anal*. 1997;8(3):124-9.
30. Leach GJ, Whiffen T. Ontogenetic, seasonal and diurnal variation in leaf volatile oils and leaf phenolics of *Angophora costata*. *Aust Sys Botany*. 1989;2(1):99-111.
31. Staudt M, Bertin N, Hansen U, Seufert G, Cicciolij P, Foster P, et al. Seasonal and diurnal patterns of monoterpene emissions from *Pinus pinea* (L.) under field conditions. *Atmos Environ*. 1997;31:145-56.
32. Guemuescu A, Ipek A, Sarihan EO, Guerbuez B, Kaya MD, Arslan N. Effects of diurnal and ontogenetic variability on essential oil composition of oregano (*Origanum vulgare* var. *hirtum*). *Asian J Chem*. 2008;20(2):1290-4.
33. Bertin N, Staudt M, Hansen U, Seufert G, Cicciolij P, Foster P, et al. Diurnal and seasonal course of monoterpene emissions from *Quercus ilex* (L.) under natural conditions application of light and temperature algorithms. *Atmos Environ*. 1997;31:135-44.