




**Appendix 1. P6 point (1)**




**Appendix 2. Point K-K9 (2)**



**Appendix 3. Point Li5 (3)**



**Appendix 4:** Comparison of the mean number of additional antinausea medications over six measurements across the three groups.



**Appendix 5. Summary of Repeated Measures ANOVA for CINV scores**

Tests	p-value	
Mauchly's Test of Sphericity		< 0.001
Repeated Measures ANOVA		p-value
Greenhouse-Geisser Correction	Within-Subjects	< 0.001
	Between-Subjects	0.006
Pairwise Comparisons	P6 & K-K9	0.92
	P6 & Li5	0.006
	K-K9 & Li5	0.004

1. Genç F, Tan M. The effect of acupressure application on chemotherapy-induced nausea, vomiting, and anxiety in patients with breast cancer. *Palliative and Supportive Care*. 2014;13(2):275-84.
2. Bertalanffy P, Hoerauf K, Fleischhackl R, Strasser H, Wicke F, Greher M, et al. Korean hand acupressure for motion sickness in prehospital trauma care: a prospective, randomized, double-blinded trial in a geriatric population. *Anesthesia & Analgesia*. 2004;98(1):220-3.
3. Yang C, Choi S, Kim J, Ryu Y, Lim Y, Kim M, et al. The Effects of Acupuncture Stimulation for Brain Activation and Alcohol Abstinence Self-Efficacy: Functional MRI Study. *Evidence-Based Complementary and Alternative Medicine*. 2017;2017:1-7.