



Reply to Letter: The Effect of Biofeedback Therapy on ICIQ-SF Scores and Urodynamic Parameters

Omer Bayrak¹, Ilker Seckiner¹

¹University of Gaziantep, School of Medicine, Department of Urology, Gaziantep, Turkey

ARTICLE INFO

Article type:

Letter to Editor

Article history:

Received: 25 Jan 2012

Revised: 03 Feb 2012

Accepted: 14 Feb 2012

Keywords:

Therapeutics

Pelvic Floor

Please cite this paper as:

Bayrak O, Seckiner I. Reply to Letter: The Effect of Biofeedback Therapy on ICIQ-SF Scores and Urodynamic Parameters. *Nephro-Urol Mon.* 2012;4(3): 593. DOI: 10.5812/numonthly.4270

Copyright © 2012 Kowsar Corp. All rights reserved.

Dear Editor,

We appreciate having the opportunity to reply to the letter regarding our article entitled, "The Effect of Biofeedback Therapy on ICIQ-SF Scores and Urodynamic Parameters in Patients with Stress Urinary Incontinence (SUI)".

Biofeedback and pelvic floor muscle exercises are advised, especially for the treatment of stress incontinence and mixed incontinence, because they are low-cost and do not have any potential side-effects (1). However, biofeedback therapy alone, without pelvic base muscle exercises, is not enough for the treatment of SUI. We have determined that pelvic floor muscle exercises, repeated at home make a very positive and important contribution to treatment, especially when biofeedback therapy is applied in the hospital as well as. We emphasize that all patients should continue these exercises at home after their therapy at the hospital has ended, so that their current healing status can be maintained and possibly even increased.

Seckiner *et al.* found that higher maximal detrusor pressure levels were moderately correlated with higher ICIQ-SF scores. They concluded from their findings that the ICIQ-SF questionnaire may provide predictive information regarding urodynamic study results in an overactive bladder (2). In our study, patients were hesitant, because although urodynamic testing is not a difficult treatment, it is an invasive examination and has a risk of infection. Because of these reasons we think that a urodynamic test is not necessary for all patients who are planning to do biofeedback therapy. However, if surgical treatment is planned, urodynamic studies can be useful.

Financial Disclosure

None declared.

References

1. Burns PA, Pranikoff K, Nochajski TH, Hadley EC, Levy KJ, Ory MG. A comparison of effectiveness of biofeedback and pelvic floor muscle exercise treatment of stress incontinence in older community-dwelling women. *J Gerontol.* 1993;48:167-74.
2. Seckiner I, Yesilli C, Mungan NA, Aykanat A, Akduman B. Correlations between the ICIQ-SF score and urodynamic findings. *Neurourol Urodyn.* 2007;26(4):492-4.

* Corresponding author: Omer Bayrak, University of Gaziantep, School of Medicine, Department of Urology, 27310 Gaziantep, Turkey. Tel: +90-5326428800, Fax: +90-3423603998, E-mail: Dromerbayrak@yahoo.com