



# Comparison of Sleeve Gastrectomy and SASI Bypass Surgery Outcomes in Patients with Obesity After Seven Months in Shiraz

Kamran Bagheri Lankarani  <sup>1</sup>, Seyed Ali Hosseini  <sup>2</sup>, Alireza Moaref  <sup>3</sup>, Hajar Khazraei  <sup>4</sup>, Seyed Vahid Hosseini  <sup>4</sup>, <sup>\*</sup>, Abdulrazzaq Kalaf Hassan <sup>5</sup>

<sup>1</sup> Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>2</sup> Department of Internal Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>3</sup> Department of Cardiology, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>4</sup> Colorectal Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>5</sup> Iraq Medical College, University of Al-Ameed College of Medicine, Karbala, Iraq

<sup>\*</sup>Corresponding author: Colorectal Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. Email: colorectal2@sums.ac.ir

Received 2024 January 13; Revised 2024 June 26; Accepted 2024 June 30.

## Abstract

**Background:** Although bariatric surgery has been introduced as a therapeutic option for patients with obesity, there is still debate on the choice of procedure.

**Objectives:** This study aimed to compare two types of bariatric surgeries in patients with obesity: Sleeve gastrectomy (SG) and single anastomosis sleeve ileal (SASI) bypass.

**Methods:** This observational prospective study compares patients with obesity who received either of the two types of bariatric surgeries at Ghadir or Shahid Faghihi hospitals in Shiraz from October 2019 to November 2020. Metabolic profiles, shear wave liver elastography (fibroscan), and cardiac evaluations (echocardiography) were performed at baseline and then seven to eleven months after the surgery.

**Results:** Forty-five patients with obesity who had undergone SG and SASI bypass entered this study. Fasting plasma glucose (FPG) and triglycerides (TG) decreased during the follow-up in both groups ( $P = 0.032$ ,  $P < 0.001$ , respectively). The fibrosis score decreased significantly from 6.45 (4.55) before surgery to 5.40 (3.60) after surgery, and the cardiac ejection fraction increased significantly from 61.5% (12.5%) before surgery to 65.0% (8.5%) after surgery following the SASI bypass compared to the SG ( $P = 0.034$ ,  $P = 0.008$ , respectively).

**Conclusions:** Despite the lack of difference in weight reduction, SASI bypass, compared to sleeve gastrectomy, may result in a more rapid improvement in cardiac function and liver fibrosis.

**Keywords:** Sleeve Gastrectomy, SASI Bypass, Obesity, Metabolic Syndrome, Liver Fibrosis

## 1. Background

Obesity is a growing pandemic attributed to high energy intake, low physical activity, and genetic factors. Other contributors include endocrine diseases, mental illnesses, and the consumption of certain medications. Obesity is associated with metabolic syndrome (MetS) and atherosclerosis, including insulin resistance, type 2 diabetes mellitus (T2DM), cardiac disease, hypertension (HTN), dyslipidemia, and liver disease (1).

Adipose tissue is a dynamic metabolic, biologically active endocrine tissue that secretes leptin and adiponectin (2). Low adiponectin concentrations and

elevated leptin levels are associated with obesity, insulin resistance, and MetS (3-5).

In the current pandemic of obesity, the demand for its treatment is on the rise. Non-surgical treatments include dietary interventions, increasing physical activity and exercise, and other lifestyle measures, which result in sustainable weight loss in only 5 - 8% of obese individuals. The results for pharmacologic treatment of obesity are not much better, with major concerns about their side effects (6, 7). Consequently, there is a growing demand for bariatric surgeries worldwide in those who fail to respond to non-surgical treatments (8, 9). However, the acceptance rate for

bariatric surgery among eligible patients is still low, and there are concerns about weight loss maintenance after surgery, which should exceed 10% for a sustainable improvement in quality of life (10).

Surgical procedures are divided into two general categories. The first is restrictive surgery, such as sleeve gastrectomy (SG), in which a large part of the stomach (about 85%) is removed vertically. Previous studies have shown that SG is an effective measure for weight reduction (9, 11-13). The other type of bariatric surgery is a malabsorptive procedure, which has also been shown to be effective, despite more safety concerns (12).

Single anastomosis sleeve ileal (SASI) bypass surgery is a relatively new bariatric surgery method in which sleeve gastrectomy is followed by a side-to-side gastro-ileal anastomosis (8). This surgery is considered a mixed procedure with both restrictive and malabsorptive components. Single anastomosis sleeve ileal bypass surgery is regarded as safer, with less malabsorption morbidity compared to classical bypass surgery, due to the preserved gastroduodenal pathway. Other benefits of SASI bypass surgery include the absence of prostheses or excluded segments, easy feasibility, and the ability to perform upper endoscopic procedures, including endoscopic retrograde cholangiopancreatography (ERCP), after surgery (6).

There are some concerns about increased bile reflux with this procedure, but there is limited comparative data on the occurrence of this complication and its clinical significance in patients who have undergone the SASI operation. Whether this procedure could result in better outcomes for patients with obesity is a matter of ongoing debate.

## 2. Objectives

This study was designed to compare the short-term outcomes of the SG and SASI bypass procedures.

## 3. Patients and Methods

Patients with morbid obesity [body mass index (BMI) > 40] were referred for bariatric surgery to two referral centers in Shiraz (Ghadir or Shahid Faghihi hospitals) affiliated with Shiraz University of Medical Sciences (Shiraz, Iran) from October 2019 to November 2020. The patients underwent either SG or SASI bypass based on their preferences after discussions with a multidisciplinary team that informed them of the known pros and cons of each operation. Both groups received standard care from the same multidisciplinary team, including a surgeon, dietitian, psychiatrist, and internist, coordinated by a trained nurse.

The inclusion criteria were patients with a BMI > 40 who came to our center for obesity treatment and were deemed candidates for surgery by a commission consisting of a surgeon, nutritionist, and psychologist, after obtaining informed consent from patients aged 18 to 60 years old. The exclusion criteria included patients with a BMI > 55, a history of severe liver diseases such as hepatitis, a history of liver surgeries, heart disorders such as heart failure, arrhythmias, myocardial infarction, or stroke, and those with a history of previous laparoscopic surgery in the abdomen.

Sixty-five patients willingly entered the study, but 20 patients withdrew during the follow-up due to adherence to restrictive protocols related to the coronavirus disease 2019 (COVID-19) pandemic. Therefore, 45 patients completed the study. These 45 patients were categorized into two groups: 12 in the SASI bypass group and 33 in the SG group. Convenience sampling was used, and the sample size was calculated using MedCalc software, considering a two-to-one ratio for the groups. The formula for determining the sample size for comparing two means is given below (14). Using this formula, and considering a type 1 error of 5% and a power of 80%, the mean and standard deviation were  $95.7 \pm 5.8$  and  $89 \pm 7.1$ , respectively. Volume formula:

$$n_1 = \frac{\left( z_{\frac{\alpha}{2}-1} + z_{\beta-1} \right)^2 \left( \frac{1}{r} \times (\sigma_1^2 + \sigma_2^2) \right) \left( \frac{1}{r} + 1 \right)}{\Delta^2}$$

$$n_2 = r \times n_1$$

$n_1$  and  $n_2$  are the sample size in each group;  $r$  = ratio of sample size ( $r = n_1/n_2$ );  $\alpha$  and  $\beta$  are type 1 error and the power respectively;  $\sigma_1$  and  $\sigma_2$  are the standard deviation of each group;  $\Delta$  is the difference of the group means.

Informed consent was obtained from all the patients. The method of SASI bypass surgery is described in a previous study (6), where the first one-third of the length of the intestine from the first ligament was anastomosed to the stomach. All procedures were performed by a single surgeon, using the same operation room materials and technicians. Data were gathered before and seven to eleven months after the surgery. The weight and BMI of the patients were measured before and seven months after surgery. The patients were followed clinically by the multidisciplinary team. Typically, 50% of bariatric surgeries at our center are sleeve gastrectomies, 25% are SASI bypasses, and 25% are classic bypass surgeries. Weight loss was assessed by the difference between

preoperative and postoperative weights. Postoperative complications were not reported in this study.

### 3.1. Biochemical Measurements

Blood samples were collected after overnight fasting (10 - 14 h). All serum samples, collected before and seven to eleven months after surgery, were prepared in a single laboratory using standard methods and stored at -70°C until analysis. Fasting plasma glucose (FPG) was measured using the glucose oxidase method. Plasma lipid profile and liver enzyme levels [alanine aminotransferase (ALT) and aspartate aminotransferase (AST)] were measured using enzymatic methods. Serum leptin, ghrelin, adiponectin, thyroid hormones, and insulin were measured using enzyme-linked immunosorbent assay (ELISA). To monitor the accuracy and precision of biochemical tests, we performed a quality control assay. HbA1C was measured using the electrophoresis method. Uric acid, total and direct bilirubin, albumin, alkaline phosphatase, calcium, and phosphorus were measured using standard biochemical methods.

### 3.2. FibroScan® and Echocardiography Studies

After at least four hours of fasting and avoiding physical activity, all patients were evaluated for the stage of fibrosis and fat content of the liver by a single operator using FibroScan® (ECHOSENS, 502, France). The grade of fibrosis was reported according to the manufacturer's instructions, ranging from F0 to F4. The controlled attenuation parameter (CAP) score, indicative of fatty changes in the liver, was also reported from S0 to S3 as a measure of liver steatosis.

All patients underwent a complete echocardiographic study using the EE Vivid-G Echo machine (Siemens, Germany). The anteroposterior diameter and biplane volume of the left atrium (LA) were measured in all patients. Left ventricular end-systolic diameter (LV ESD) and left ventricular end-diastolic diameter (LV EDD) were calculated in the parasternal long-axis view. As defined by the guidelines of the American Society of Echocardiography (ASE), the bipolar left ventricle ejection fraction (LVEF) was calculated using the Simpson method, and the diastolic function of the right ventricle (RV) was measured using pulse wave Doppler of the mitral valve (MV) flow and tissue Doppler study of the MV annulus.

All cardiac valves were evaluated in the transthoracic echocardiography (TTE) exam, and moderate to severe valvular stenosis or regurgitation was considered significant valvular heart disease. Systolic pulmonary

artery pressure (SPAP) or mean pulmonary artery pressure (mean PAP) was measured by tricuspid regurgitation (TR) gradient or pulmonary acceleration time, respectively. The apical four-chamber view was used to evaluate the sizes of the right atrium (RA) and RV, as well as RV systolic function.

### 3.3. Statistical Analysis

Data were analyzed using the statistical package for social sciences (SPSS) for Windows, version 22 (Chicago, IL, USA). Data were expressed as median (IQR). The Mann-Whitney test and Wilcoxon test were used to compare each parameter of the groups before and after the surgery or between surgeries.

### 3.4. Ethical Considerations

The study protocol was approved by the Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.MED.REC.1399.555).

## 4. Results

### 4.1. Characteristics of the Patients

The patients' ages ranged from 18 to 57 years, with a median (IQR) of 37.0 (14.0) years, which was not significantly different between the two groups ( $P=0.877$ , Table 1). In our study, the weight and BMI of the patients significantly decreased in both groups seven months after surgery. The median weight was 117.5 (30.5) kg before surgery and decreased to 85.0 (24.7) kg after surgery ( $P < 0.001$ ). Specifically, the median weight for the SG group was 112.8 (24.5) kg before surgery and decreased to 85.0 (27.5) kg after surgery ( $P < 0.001$ ). For the SASI bypass group, the median weight was 135.0 (38.5) kg before surgery and decreased to 91.5 (16.0) kg after surgery ( $P = 0.002$ ). However, the decrease was not significantly different between the two types of surgery ( $P = 0.227$ , Table 1).

The percent change in BMI was a 29.8% decrease after the surgeries ( $P < 0.001$ ). A 29.44% decrease was seen after SG, and a 30.79% decrease was observed after SASI bypass. Both SG and SASI bypass procedures significantly decreased the BMI, but the decrease was not significantly different between the two types of surgery ( $P = 0.238$ , Table 1).

### 4.2. Biochemical Measurements

The two groups were not similar at baseline, but the differences were not statistically significant (Table 2  $P >$

**Table 1.** Demographic Characteristics of Patients in the Two Groups Before and After 7 Months<sup>a</sup>

| Parameters                 | Sleeve Gastrectomy (n=33) | SASI Bypass (n=12) | P-Value <sup>b</sup> | Values       |
|----------------------------|---------------------------|--------------------|----------------------|--------------|
| Age (y)                    | 37.0 (14.5)               | 37.0 (9.5)         | 0.877                | 37.0 (14.0)  |
| Sex                        |                           |                    |                      |              |
| Male                       | 12 (36.4%)                | 4 (33.3%)          | 0.571                | 16 (35.6%)   |
| Female                     | 21 (63.6%)                | 8 (66.7%)          |                      | 29 (64.4%)   |
| Weight before surgery (kg) | 112.8 (24.5)              | 135.0 (38.5)       | 0.010                | 117.5 (30.5) |
| Weight decrease (kg)       | 36.0 (18.0)               | 42.5 (25.2)        | 0.227                | 36.0 (19.0)  |
| % loss in body weight      | 32.1% (14.8)              | 29.9% (13.4)       | 0.843                | 29.9% (14.7) |
| BMI decrease               | 13.6 (7.0)                | 15.1 (7.5)         | 0.238                | 13.7 (6.5)   |

Abbreviations: BMI, body mass index; SASI, single anastomosis sleeve ileal.

<sup>a</sup> Values are expressed as median (IQR) or No. (%) unless otherwise indicated.

<sup>b</sup> Mann-Whitney test and chi-square test.

0.05). The mean high-density lipoprotein-cholesterol (HDL-C) level (mg/dL) increased significantly after SG ( $P < 0.001$ ), but the rise was not significant after SASI bypass. low-density lipoprotein (LDL) levels decreased following SASI bypass but increased after SG (Table 2). Liver enzymes (AST, ALT, ALKP) and creatinine significantly decreased after SG ( $P = 0.005$ ), but did not decrease after SASI bypass. The albumin level increased significantly after SG ( $P = 0.010$ ) but did not change after SASI bypass. The decrease in BMI was inversely correlated with ALT and leptin levels after SG ( $r = -0.371$ ,  $P = 0.040$ ;  $r = -0.383$ ,  $P = 0.031$ , respectively), but this correlation was not seen following SASI bypass.

Ghrelin and leptin levels decreased significantly after both types of surgeries, but there were no significant differences between the two groups ( $P = 0.571$ ,  $P = 0.414$ , respectively). Adiponectin levels increased substantially after both surgeries, but the difference was not significant between them ( $P = 0.598$ , Table 3). HbA1C decreased significantly after both types of surgery, but the decrease was similar between the groups ( $P = 0.422$ , Table 1). The drop in FPG level following both SASI bypass and SG was not significant ( $P = 0.055$  and  $P = 0.130$ , respectively). Insulin levels and homeostasis model assessment-estimated insulin resistance (HOMA-IR) decreased after SASI bypass, but this did not reach statistical significance ( $P = 0.859$  and  $P = 0.477$ , respectively). However, these parameters increased significantly after SG ( $P = 0.022$  and  $P = 0.017$ , respectively).

The blood urea nitrogen (BUN) concentration did not change significantly in either group. Uric acid decreased substantially in both groups, but the decrease was not significantly different between the groups ( $P = 0.147$ ). Although total bilirubin and direct bilirubin levels increased after both types of surgeries, these changes

remained within normal limits. Electrolytes and thyroid hormones also changed within normal ranges after the surgeries (Tables 2, 3).

#### 4.3. FibroScan® and Echocardiography Studies

The CAP score decreased significantly after both types of surgeries, with no significant difference between the groups ( $P = 0.772$ ) (Table 4). The fibrosis score decreased significantly only after SASI bypass ( $P = 0.034$ ) (Table 4). Tables 5 and 6 show the fibroscan grades and steatosis stages in the two groups before and after the surgeries.

#### 4.4. Echocardiography Study

The LV diastolic function, RV function, RV size, RA size, and pulmonary artery pressure were within the normal range in all patients before and after the surgeries. There was no significant valvular heart disease or pulmonary hypertension before or after the surgeries.

Before the surgery, two patients in the sleeve group had grade 1 heart failure, and three in the SASI bypass group had stage 1 heart failure. However, after the surgery, all patients showed improvement with ejection fraction (EF)  $> 55\%$ . The EF percentage increased significantly after SASI bypass ( $P = 0.008$ ) but not after SG ( $P = 0.620$ ), and the difference between the two groups was significant ( $P = 0.045$ , Table 4). The decrease in BMI after SG correlated with the rise in EF ( $r = 0.416$ ,  $P = 0.031$ ), indicating that a greater decline in BMI led to a larger increase in EF percentage.

Before and after the surgery, the LA volume in all patients was normal (i.e., less than 34 mL per square meter) and decreased non-significantly after both surgeries (SASI bypass:  $P = 0.109$ , SG:  $P = 0.808$ ). The LV

**Table 3.** Hormone Levels Before and Seven Months After the Surgeries

| Parameters and Time        | Groups                      |                      |                |                      |
|----------------------------|-----------------------------|----------------------|----------------|----------------------|
|                            | Sleeve Gastrectomy (n = 33) |                      | SASI (n = 12)  |                      |
|                            | Median (IQR)                | P-Value <sup>b</sup> | Mean ± SD      | P-Value <sup>b</sup> |
| <b>Adiponectin (μg/mL)</b> |                             |                      |                | 0.004                |
| Before surgery             | 3.36 (3.11)                 | < 0.001              | 3.33 ± 2.82    | 0.885                |
| After surgery              | 5.23 (4.04)                 |                      | 5.74 ± 3.41    | 0.598                |
| <b>Leptin (ng/mL)</b>      |                             |                      |                | 0.003                |
| Before surgery             | 43.62 (51.18)               | < 0.001              | 74.07 ± 62.06  | 0.082                |
| After surgery              | 18.90 (27.15)               |                      | 28.66 ± 24.00  | 0.414                |
| <b>Ghrelin (μg/mL)</b>     |                             |                      |                | 0.099                |
| Before surgery             | 0.91 (0.45)                 | 0.010                | 0.88 ± 0.30    | 0.885                |
| After surgery              | 0.55 (0.55)                 |                      | 0.75 ± 0.62    | 0.571                |
| <b>TSH (mIU/L)</b>         |                             |                      |                | 0.814                |
| Before surgery             | 2.59 (2.06)                 | 0.232                | 2.86 ± 5.97    | 0.947                |
| After surgery              | 2.40 (1.64)                 |                      | 2.50 ± 2.52    | 0.828                |
| <b>T3 (ng/dL)</b>          |                             |                      |                | 0.168                |
| Before surgery             | 148.00 (30.50)              | 0.020                | 140.00 ± 16.00 | 0.686                |
| After surgery              | 121.50 (33.25)              |                      | 110.5 ± 122.53 | 0.478                |
| <b>T4 (μg/dL)</b>          |                             |                      |                | 0.508                |
| Before surgery             | 8.57 (1.64)                 | 0.977                | 7.96 ± 3.80    | 0.487                |
| After surgery              | 8.70 (2.10)                 |                      | 7.50 ± 3.10    | 0.229                |

Abbreviations: SASI, single anastomosis sleeve ileal; TSH, thyroid stimulating hormone.

<sup>a</sup> Mann-Whitney test was used to assess significant differences between the two surgery groups.

<sup>b</sup> Wilcoxon test was used to assess significant changes within each group following surgery.

EDD in the two groups before and after the surgeries was within normal ranges for all patients.

## 5. Discussion

The rational goal of bariatric surgeries should not only be to decrease weight but also to ensure the patient's quality of life and wellbeing, including the improvement of vital organ function and alleviating the adverse effects of obesity on the heart, liver, kidneys, and other organs. In this study, BMI decreased significantly with both SG and SASI bypass operations, with a trend toward more weight reduction with the latter. These results are similar to the study by Emile et al., which showed that the percentage of excess weight loss (EWL%) at six months postoperatively was similar between the two groups (15).

In this study, it was shown that insulin levels and HOMA-IR decreased after SASI bypass and increased after SG. This is also in line with Emile et al.'s study results, which showed that the improvement in T2DM after SASI bypass was better compared to SG (15). Our results indicated that HbA1C, a long-term predictor of glucose control, decreased significantly and to a similar degree in both groups, as did the TG level. Emile et al. also

reported a better improvement in the metabolic profile in the long term after SASI bypass. They indicated that the EWL% reached 90% at one year and the remission rates of hypertension, hypercholesterolemia, and hypertriglyceridemia were 86%, 100%, and 97%, respectively (15). In another study, Salama et al. showed significant decreases in the plasma levels of FPG, insulin, and LDL and a significant increase in HDL plasma levels following SASI bypass (8).

In this study, adiponectin levels increased significantly, and leptin and ghrelin concentrations decreased considerably after both surgeries ( $P < 0.001$  for both). Similar results were reported by Buzga et al. (4). Ghrelin and leptin levels decreased significantly after both types of surgeries, but there were no significant differences between the two groups ( $P = 0.571$ ,  $P = 0.414$ ). Adiponectin levels increased substantially after both surgeries, but the difference was not significant between them ( $P = 0.598$ , Table 3).

In our study, after SG, the creatinine level significantly decreased ( $P = 0.005$ ). The mean concentrations of liver enzymes decreased but were within the normal range in both groups before and after surgeries, although the decrease was more prominent

**Table 4.** FibroScan® and Echocardiographic Parameters Before and Seven Months After the Surgeries

| Parameters and Time            | Groups                      |                      |               |                      | P-Value <sup>a</sup> |  |
|--------------------------------|-----------------------------|----------------------|---------------|----------------------|----------------------|--|
|                                | Sleeve Gastrectomy (n = 33) |                      | SASI (n = 12) |                      |                      |  |
|                                | Mean ± SD                   | P-Value <sup>b</sup> | Mean ± SD     | P-Value <sup>b</sup> |                      |  |
| EF (%)                         |                             | 0.620                |               | 0.008                |                      |  |
| Before surgery                 | 63.0% ± 10.7%               |                      | 59.0% ± 12.0% |                      | 0.047                |  |
| After surgery                  | 64.0% ± 7.0%                |                      | 66.0% ± 8.5%  |                      | 0.415                |  |
| LA volume (mL/m <sup>2</sup> ) |                             | 0.808                |               | 0.109                |                      |  |
| Before surgery                 | 45.0 ± 17.0                 |                      | 43.0 ± 14.0   |                      | 0.568                |  |
| After surgery                  | 48.0 ± 13.0                 |                      | 45.0 ± 26.5   |                      | 0.767                |  |
| LVEDD (cm)                     |                             | 0.224                |               | 0.345                |                      |  |
| Before surgery                 | 4.6 ± 0.9                   |                      | 4.9 ± 1.1     |                      | 0.231                |  |
| After surgery                  | 4.7 ± 0.6                   |                      | 5.2 ± 0.7     |                      | 0.142                |  |
| FibroScan score (kPa)          |                             | 0.131                |               | 0.034                |                      |  |
| Before surgery                 | 6.1 ± 5.1                   |                      | 6.9 ± 4.2     |                      | 0.176                |  |
| After surgery                  | 5.5 ± 4.4                   |                      | 5.5 ± 4.0     |                      | 0.445                |  |
| CAP score (dB/m)               |                             | < 0.001              |               | 0.041                |                      |  |
| Before surgery                 | 329.0 ± 75.0                |                      | 318.5 ± 129.5 |                      | 0.839                |  |
| After surgery                  | 249.5 ± 113.2               |                      | 251.0 ± 76.7  |                      | 0.772                |  |
| Steatoses (%)                  |                             | < 0.001              |               | 0.327                |                      |  |
| Before surgery                 | 79.0% ± 23.0%               |                      | 69.0% ± 63.5% |                      | 0.083                |  |
| After surgery                  | 31.0% ± 72.0%               |                      | 29.0% ± 56.0% |                      | 0.689                |  |

Abbreviations: SASI, single anastomosis sleeve ileal; EF, ejection fraction; LA, left atrium; LVEDD, left ventricular end-diastolic diameter; CAP, controlled attenuation parameter.

<sup>a</sup> Mann-Whitney test was used to assess significant differences between the two surgery groups.

<sup>b</sup> Wilcoxon test was used to assess significant changes within each group following surgery.

**Table 5.** FibroScan® Grades in the Two Groups Before and After the Surgeries

| Operation Type      | FibroScan Grade, Frequency (%) |           |       | P-Value <sup>a</sup> |
|---------------------|--------------------------------|-----------|-------|----------------------|
|                     | F0 - F1                        | F2 - F3   | F4    |                      |
| Sleeve gastrectomy  |                                |           |       | 0.161                |
| Before surgery      | 19 (57.6)                      | 12 (36.4) | 2 (6) |                      |
| After surgery       | 24 (72.8)                      | 8 (24.2)  | 1 (3) |                      |
| SASI bypass surgery |                                |           |       | 0.095                |
| Before surgery      | 6 (50)                         | 6 (50)    | 0     |                      |
| After surgery       | 9 (75)                         | 3 (25)    | 0     |                      |

Abbreviation: SASI, single anastomosis sleeve ileal.

<sup>a</sup>Pearson chi-square test.

following SG ( $P < 0.05$ ). The reduction in BMI after SG was inversely correlated with ALT and leptin levels.

The CAP score, an indicator of fatty changes in the liver, decreased significantly after both types of surgeries, with no significant difference between the groups ( $P = 0.772$ ) (Tables 4, 6). The liver fibrosis score decreased significantly only after SASI bypass surgery ( $P = 0.034$ ).

The EF% increased significantly after SASI bypass surgery ( $P = 0.008$ ). The contraction power of the heart also improved due to weight loss, which is a sign of metabolic syndrome improvement.

This study, with a short-term follow-up, revealed comparable efficacy of SG and SASI bypass surgeries in weight loss and several metabolic indicators while showing more rapid improvement of liver fibrosis and cardiac function with SASI bypass. The significance of

**Table 6.** Steatosis Stages in the Two Groups Before and After the Surgeries

| Operation Type             | Steatosis Stage, Frequency (%) |          |          |           | P-Value <sup>a</sup> |
|----------------------------|--------------------------------|----------|----------|-----------|----------------------|
|                            | S0                             | S1       | S2       | S3        |                      |
| <b>Sleeve gastrectomy</b>  |                                |          |          |           |                      |
| Before surgery             | 3 (9.1)                        | 1 (3)    | 6 (18.2) | 23 (69.7) |                      |
| After surgery              | 14 (42.5)                      | 5 (15.1) | 4 (12.1) | 10 (30.3) |                      |
| <b>SASI bypass surgery</b> |                                |          |          |           |                      |
| Before surgery             | 2 (16.7)                       | 0        | 2 (16.7) | 8 (66.6)  |                      |
| After surgery              | 3 (25)                         | 5 (41.7) | 1 (8.3)  | 3 (25)    |                      |

Abbreviation: SASI, single anastomosis sleeve ileal.

<sup>a</sup> Pearson chi-square test.

this advantage of SASI bypass needs to be demonstrated in long-term studies.

### 5.1. Limitations

This study reports on a small series from a single center, comparing heterogeneous groups (12 SASI vs. 33 SG). There was no matching and no randomization in our groups because the patients chose the type of surgery. The generalization of our findings needs to be confirmed in larger multicenter studies.

### 5.2. Conclusions

In conclusion, our data revealed that both SASI bypass and SG are effective and safe in the treatment of morbid obesity. The more rapid improvement of cardiac function and liver fibrosis observed with SASI bypass could be an advantage in certain circumstances. However, this finding needs to be confirmed in larger, randomized long-term studies.

### Acknowledgements

We thank the Clinical Research Development Center in Namazi Hospital for their assistance in data analysis.

### Footnotes

**Authors' Contribution:** S. V. H. contributed to the conception of the work, conducted the study, revised the draft, approved the final version of the manuscript, and agreed with all aspects of the work. S. A. H. and A. M. contributed to the conception of the work, drafted and revised the draft, approved the final version of the manuscript, and agreed with all aspects of the work. H. Kh. contributed to the conception of the work, conducted the study, revised the draft, approved the

final version of the manuscript, and agreed with all aspects of the work. K. B. L. contributed to the conception of the work, revised the draft, approved the final version of the manuscript, and agreed with all aspects of the work. A. K. H. contributed to the conception of the work, revised the draft, approved the final version of the manuscript, and agreed with all aspects of the work.

**Conflict of Interests Statement:** Kamran Bagheri Lankarani is one of the boardman in Shiraz E-Medical Journal.

**Data Availability:** The datasets generated and analyzed during the current study are not publicly available due to the ethics protocol, but they are available from the corresponding author upon reasonable request.

**Ethical Approval:** All methods of the study protocol were carried out in accordance with the Ethics Committee of Shiraz University of Medical Sciences in accordance with the Declaration of Helsinki (IR.SUMS.MED.REC.1399.555).

**Funding/Support:** The study was extracted from Seyed Ali Hosseini thesis of internal medicine field (no. 20941) in Shiraz University of Medical Sciences.

**Informed Consent:** Informed consent was obtained from all patients.

### References

- Bluher M. Adipose tissue dysfunction in obesity. *Exp Clin Endocrinol Diabetes*. 2009;117(6):241-50. [PubMed ID: 19358089]. <https://doi.org/10.1055/s-0029-1192044>.
- Ricci R, Bevilacqua F. The potential role of leptin and adiponectin in obesity: a comparative review. *Vet J*. 2012;191(3):292-8. [PubMed ID: 21592831]. <https://doi.org/10.1016/j.tvjl.2011.04.009>.
- Dong M, Ren J. What fans the fire: insights into mechanisms of leptin in metabolic syndrome-associated heart diseases. *Curr Pharm Des*. 2014;20(4):652-8. [PubMed ID: 23688014]. <https://doi.org/10.2174/138161282004140213160930>.

4. Buzga M, Zavadilova V, Holeczi P, Svagera Z, Svorc P, Foltys A, et al. Dietary intake and ghrelin and leptin changes after sleeve gastrectomy. *Wideochir Inne Tech Maloinwazyjne*. 2014;9(4):554-61. [PubMed ID: 25561993]. [PubMed Central ID: PMC4280419]. <https://doi.org/10.5114/wiitm.2014.45437>.
5. Paluszakiewicz R, Kalinowski P, Wroblewski T, Bartoszewicz Z, Bialobrzeska-Paluszakiewicz J, Ziarkiewicz-Wroblewska B, et al. Prospective randomized clinical trial of laparoscopic sleeve gastrectomy versus open Roux-en-Y gastric bypass for the management of patients with morbid obesity. *Wideochir Inne Tech Maloinwazyjne*. 2012;7(4):225-32. [PubMed ID: 23362420]. [PubMed Central ID: PMC3557743]. <https://doi.org/10.5114/wiitm.2012.32384>.
6. Yanovski SZ, Yanovski JA. Long-term drug treatment for obesity: a systematic and clinical review. *JAMA*. 2014;311(1):74-86. [PubMed ID: 24231879]. [PubMed Central ID: PMC3928674]. <https://doi.org/10.1001/jama.2013.281361>.
7. No Author Listed. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. *Circulation*. 2002;106(25):3143. <https://doi.org/10.1161/circ.106.25.3143>.
8. Salama TMS, Sabry K, Ghamrini YE. Single Anastomosis Sleeve Ileal Bypass: New Step in the Evolution of Bariatric Surgeries. *J Invest Surg*. 2017;30(5):291-6. [PubMed ID: 27768400]. <https://doi.org/10.1080/08941939.2016.1241841>.
9. Baltasar A, Serra C, Perez N, Bou R, Bengoechea M, Ferri L. Laparoscopic sleeve gastrectomy: a multi-purpose bariatric operation. *Obes Surg*. 2005;15(8):1124-8. [PubMed ID: 16197783]. <https://doi.org/10.1381/0960892055002248>.
10. De Luca M, Angrisani L, Himpens J, Busetto L, Scopinaro N, Weiner R, et al. Indications for Surgery for Obesity and Weight-Related Diseases: Position Statements from the International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO). *Obes Surg*. 2016;26(8):1659-96. [PubMed ID: 27412673]. [PubMed Central ID: PMC6037181]. <https://doi.org/10.1007/s11695-016-2271-4>.
11. Hung J, McQuillan BM, Thompson PL, Beilby JP. Circulating adiponectin levels associate with inflammatory markers, insulin resistance and metabolic syndrome independent of obesity. *Int J Obes (Lond)*. 2008;32(5):772-9. [PubMed ID: 18253163]. <https://doi.org/10.1038/sj.ijo.0803793>.
12. Esteghamati A, Khalilzadeh O, Anvari M, Rashidi A, Mokhtari M, Nakhjavani M. Association of serum leptin levels with homeostasis model assessment-estimated insulin resistance and metabolic syndrome: the key role of central obesity. *Metab Syndr Relat Disord*. 2009;7(5):447-52. [PubMed ID: 19419266]. <https://doi.org/10.1089/met.2008.0100>.
13. Sandeep S, Gokulakrishnan K, Velmurugan K, Deepa M, Mohan V. Visceral & subcutaneous abdominal fat in relation to insulin resistance & metabolic syndrome in non-diabetic south Indians. *Indian J Med Res*. 2010;131:629-35. [PubMed ID: 20516533].
14. Elyasinia F, Jalali SM, Zarini S, Sadeghian E, Sorush A, Pirouz A. The Effect of Laparoscopic Sleeve Gastrectomy and Gastric Bypass Surgery on Non-Alcoholic Steatohepatitis in Iranian Patients with Obesity. *Middle East J Dig Dis*. 2021;13(3):200-7. [PubMed ID: 36606220]. [PubMed Central ID: PMC9489465]. <https://doi.org/10.34172/mejdd.2021.226>.
15. Emile SH, Madyan A, Mahdy T, Elshobaky A, Elbanna HG, Abdel-Razik MA. Single anastomosis sleeve ileal (SASI) bypass versus sleeve gastrectomy: a case-matched multicenter study. *Surg Endosc*. 2021;35(2):652-60. [PubMed ID: 32072282]. <https://doi.org/10.1007/s00464-020-07430-w>.

**Table 2.** Laboratory Data of the Patients in the Two Groups

| Parameters and Time             | Sleeve Gastrectomy (n=33) <sup>a</sup> | P-Value <sup>b</sup> | SASI Bypass (n=12) <sup>a</sup> | P-Value <sup>b</sup> | P-Value <sup>c</sup> |
|---------------------------------|--|----------------------|---------------------------------|----------------------|----------------------|
| <b>Cholesterol (mg/dL)</b>      |  | 0.009                |                                 | 0.347                |                      |
| Before surgery                  | 180.0 (31.0)                           |                      | 168.0 (44.2)                    |                      | 0.323                |
| After surgery                   | 192.0 (63.0)                           |                      | 179.5 (52.0)                    |                      | 0.228                |
| <b>LDL-C (mg/dL)</b>            |  | 0.159                |                                 | 0.583                |                      |
| Before surgery                  | 117.0 (20.0)                           |                      | 104.5 (26.8)                    |                      | 0.329                |
| After surgery                   | 117.0 (28.0)                           |                      | 108.0 (65.8)                    |                      | 0.255                |
| <b>HDL-C (mg/dL)</b>            |  | < 0.001              |                                 | 0.136                |                      |
| Before surgery                  | 47.0 (13.0)                            |                      | 47.0 (13.5)                     |                      | 0.924                |
| After surgery                   | 62.0 (16.0)                            |                      | 59.0 (20.5)                     |                      | 0.233                |
| <b>Triglyceride (mg/dL)</b>     |  | 0.001                |                                 | 0.038                |                      |
| Before surgery                  | 157.0 (83.0)                           |                      | 142.5 (76.0)                    |                      | 0.588                |
| After surgery                   | 112.0 (64.0)                           |                      | 93.0 (26.7)                     |                      | 0.184                |
| <b>FPG (mg/dL)</b>              |  | 0.130                |                                 | 0.055                |                      |
| Before surgery                  | 100.0 (27.0)                           |                      | 108.0 (26.7)                    |                      | 0.174                |
| After surgery                   | 98.0 (13.0)                            |                      | 99.5 (15.5)                     |                      | 0.797                |
| <b>HbA1C (%)</b>                |  | < 0.001              |                                 | 0.003                |                      |
| Before surgery                  | 5.8 (0.5)                              |                      | 5.7 (1.1)                       |                      | 0.682                |
| After surgery                   | 4.9 (0.8)                              |                      | 5.0 (0.7)                       |                      | 0.422                |
| <b>Insulin (mIU/L)</b>          |  | 0.022                |                                 | 0.859                |                      |
| Before surgery                  | 4.7 (6.0)                              |                      | 7.8 (6.4)                       |                      | 0.073                |
| After surgery                   | 6.5 (11.6)                             |                      | 6.3 (13.1)                      |                      | 0.713                |
| <b>HOMA-IR</b>                  |  | 0.017                |                                 | 0.477                |                      |
| Before surgery                  | 1.1 (1.4)                              |                      | 2.4 (1.9)                       |                      | 0.022                |
| After surgery                   | 1.6 (2.9)                              |                      | 1.5 (3.1)                       |                      | 0.746                |
| <b>Uric acid (mg/dL)</b>        |  | 0.004                |                                 | 0.022                |                      |
| Before surgery                  | 5.5 (1.9)                              |                      | 6.8 (1.8)                       |                      | 0.137                |
| After surgery                   | 4.8 (1.3)                              |                      | 5.3 (0.5)                       |                      | 0.147                |
| <b>Total bilirubin (mg/dL)</b>  |  | < 0.001              |                                 | 0.005                |                      |
| Before surgery                  | 0.6 (0.4)                              |                      | 0.5 (0.5)                       |                      | 0.664                |
| After surgery                   | 0.8 (0.4)                              |                      | 0.7 (0.5)                       |                      | 0.542                |
| <b>Direct bilirubin (mg/dL)</b> |  | < 0.001              |                                 | 0.003                |                      |
| Before surgery                  | 0.1 (0.1)                              |                      | 0.1 (0.1)                       |                      | 0.818                |
| After surgery                   | 0.3 (0.2)                              |                      | 0.3 (0.2)                       |                      | 0.946                |
| <b>Albumin (g/dL)</b>           |  | 0.001                |                                 | 1.000                |                      |
| Before surgery                  | 4.2 (0.4)                              |                      | 4.3 (0.3)                       |                      | 0.661                |
| After surgery                   | 4.5 (0.5)                              |                      | 4.2 (0.5)                       |                      | 0.047                |
| <b>AST (U/L)</b>                |  | 0.001                |                                 | 0.388                |                      |
| Before surgery                  | 30.5 (11.7)                            |                      | 33.5 (14.5)                     |                      | 0.303                |
| After surgery                   | 24.0 (9.0)                             |                      | 31.5 (12.2)                     |                      | 0.004                |
| <b>ALT (U/L)</b>                |  | 0.001                |                                 | 0.248                |                      |
| Before surgery                  | 33.5 (22.5)                            |                      | 37.5 (26.2)                     |                      | 0.562                |
| After surgery                   | 20.0 (16.0)                            |                      | 32.5 (19.2)                     |                      | 0.027                |
| <b>ALP (U/L)</b>                |  | 0.001                |                                 | 0.209                |                      |
| Before surgery                  | 182.0 (69.5)                           |                      | 184.0 (35.5)                    |                      | 0.626                |
| After surgery                   | 150.5 (56.0)                           |                      | 158.5 (38.7)                    |                      | 0.358                |
| <b>Ca (mg/dL)</b>               |  | 0.845                |                                 | 0.592                |                      |
| Before surgery                  | 9.5 (0.8)                              |                      | 9.6 (0.7)                       |                      | 0.616                |
| After surgery                   | 9.5 (0.5)                              |                      | 9.5 (0.9)                       |                      | 0.812                |
| <b>Phosphorus (mg/dL)</b>       |  | 0.673                |                                 | 0.049                |                      |
| Before surgery                  | 3.6 (0.6)                              |                      | 3.2 (1.1)                       |                      | 0.083                |

| Parameters and Time                 | Sleeve Gastrectomy (n=33) <sup>a</sup> | P-Value <sup>b</sup> | SASI Bypass (n=12) <sup>a</sup> | P-Value <sup>b</sup> | P-Value <sup>c</sup> |
|-------------------------------------|--|----------------------|---------------------------------|----------------------|----------------------|
| After surgery                       | 3.5 (1.0)                              |                      | 3.8 (0.7)                       |                      | 0.460                |
| <b>WBC (×10<sup>9</sup>/L)</b>      |  | < 0.001              |                                 | 0.002                |                      |
| Before surgery                      | 8.3 (3.7)                              |                      | 8.2 (3.1)                       |                      | 0.772                |
| After surgery                       | 6.6 (2.1)                              |                      | 6.2 (2.6)                       |                      | 0.588                |
| <b>Hemoglobin (g/dL)</b>            |  | 0.341                |                                 | 0.724                |                      |
| Before surgery                      | 13.9 (2.1)                             |                      | 13.5 (1.8)                      |                      | 0.493                |
| After surgery                       | 14.1 (2.7)                             |                      | 14.0 (1.3)                      |                      | 0.882                |
| <b>MCV (fL)</b>                     |  | < 0.001              |                                 | 0.019                |                      |
| Before surgery                      | 81.9 (6.7)                             |                      | 80.9 (8.6)                      |                      | 0.885                |
| After surgery                       | 84.6 (7.0)                             |                      | 85.9 (5.1)                      |                      | 0.498                |
| <b>Platelet (×10<sup>9</sup>/L)</b> |  | 0.001                |                                 | 0.045                |                      |
| Before surgery                      | 263.5 (101.0)                          |                      | 275.5 (93.7)                    |                      | 0.895                |
| After surgery                       | 250.0 (86.0)                           |                      | 239.0 (112.5)                   |                      | 0.735                |

Abbreviations: SASI, single anastomosis sleeve ileal; BMI, body mass index; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; FPG, fasting plasma glucose; AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; Ca, calcium; WBC, white blood cells; MCV, mean corpuscular volume; HOMA-IR, homeostasis model assessment-estimated insulin resistance.

<sup>a</sup> Values are expressed as median (IQR).

<sup>b</sup> Wilcoxon test was used to assess significant changes within each group following surgery.

<sup>c</sup> Mann-Whitney test was used to assess significant differences between the two surgery groups.